

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 July 2001 (05.07.2001)

PCT

(10) International Publication Number
WO 01/48245 A2

- (51) International Patent Classification⁷: **C12Q 1/68** 06516 (US). **LEACH, Martin** [GB/US]; 884 School Street, Webster, MA 01570 (US).
- (21) International Application Number: PCT/US00/35346
- (22) International Filing Date:
27 December 2000 (27.12.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/472,688 27 December 1999 (27.12.1999) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 09/472,688 (CIP)
Filed on 27 December 1999 (27.12.1999)
- (71) Applicant (for all designated States except US): **CURA-GEN CORPORATION** [US/US]; 11th floor, 555 Long Wharf Drive, Branford, CT 06511 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **SHIMKETS, Richard, A.** [US/US]; 191 Lete Street, West Haven, CT
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— Without international search report and to be republished upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NUCLEIC ACIDS CONTAINING SINGLE NUCLEOTIDE POLYMORPHISMS AND METHODS OF USE THEREOF

(57) Abstract: The invention provides nucleic acids containing single-nucleotide polymorphisms identified for transcribed human sequences, as well as methods of using the nucleic acids.

WO 01/48245 A2

NUCLEIC ACIDS CONTAINING SINGLE NUCLEOTIDE POLYMORPHISMS AND METHODS OF USE THEREOF

FIELD OF THE INVENTION

The invention relates generally to nucleic acids and polypeptides and in particular to
5 the identification of human single nucleotide polymorphisms based on at least one gene
product that was not previously described.

BACKGROUND OF THE INVENTION

Sequence polymorphism-based analysis of nucleic acid is generally based on
alterations in nucleic acid sequences between related individuals. This analysis has been
10 widely used in a variety of genetic, diagnostic, and forensic applications. For example,
polymorphism analyses are used in identity and paternity analysis, and in genetic mapping
studies.

Several different types of polymorphisms in nucleic acid have been described. One
such type of variation is a restriction fragment length polymorphism (RFLP). RFLPs can
15 create or delete a recognition sequence for a restriction endonuclease in one nucleic acid
relative to a second nucleic acid. The result of the variation is in an alteration the relative
length of restriction enzyme generated DNA fragments in the two nucleic acids.

Other polymorphisms take the form of short tandem repeats (STR) sequences, which
are also referred to as variable numbers of tandem repeat (VNTR) sequences. STR sequences
20 typically that include tandem repeats of 2, 3, or 4 nucleotide sequences that are present in a
nucleic acid from one individual but absent from a second, related individual at the
corresponding genomic location.

Other polymorphisms take the form of single nucleotide variations, termed single
nucleotide polymorphisms (SNPs), between individuals. A SNP can, in some instances, be
25 referred to as a "cSNP" to denote that the nucleotide sequence containing the SNP originates
as a cDNA.

SNPs can arise in several ways. A single nucleotide polymorphism may arise due to a
substitution of one nucleotide for another at the polymorphic site. Substitutions can be
transitions or transversions. A transition is the replacement of one purine nucleotide by

another purine nucleotide, or one pyrimidine by another pyrimidine. A transversion is the replacement of a purine by a pyrimidine, or the converse.

Single nucleotide polymorphisms can also arise from a deletion of a nucleotide or an insertion of a nucleotide relative to a reference allele. Thus, the polymorphic site is a site at which one allele bears a gap with respect to a single nucleotide in another allele. Some SNPs occur within, or near genes. One such class includes SNPs falling within regions of genes encoding for a polypeptide product. These SNPs may result in an alteration of the amino acid sequence of the polypeptide product and give rise to the expression of a defective or other variant protein. Such variant products can, in some cases result in a pathological condition, e.g., genetic disease. Examples of genes in which a polymorphism within a coding sequence gives rise to genetic disease include sickle cell anemia and cystic fibrosis. Other SNPs do not result in alteration of the polypeptide product. Of course, SNPs can also occur in noncoding regions of genes.

SNPs tend to occur with great frequency and are spaced uniformly throughout the genome. The frequency and uniformity of SNPs means that there is a greater probability that such a polymorphism will be found in close proximity to a genetic locus of interest.

SUMMARY OF THE INVENTION

The invention is based in part on the discovery of single nucleotide polymorphisms (SNPs) in regions of human DNA.

Accordingly, in one aspect, the invention provides nucleic acid sequences comprising nucleic acid segments of both publicly known and novel genes, including the polymorphic site. The segments can be DNA or RNA, and can be single- or double-stranded. Preferred segments include a biallelic polymorphic site.

The invention further provides allele-specific oligonucleotides that hybridize to a segment of a fragment shown in Table 1, column 4, or its complement. These oligonucleotides can be probes or primers. Also provided are isolated nucleic acids comprising a sequence shown in Table 1, column 4, in which the polymorphic site within the sequence is occupied by a base other than the reference bases shown in Table 1, columns 5 and 6.

The invention further provides a method of analyzing a nucleic acid from an individual. The method determines which base is present at any one of the polymorphic sites shown in Table 1. Optionally, a set of bases occupying a set of polymorphic sites shown in Table 1 is determined. This type of analysis can be performed on a number of individuals,
5 who are tested for the presence of a disease phenotype.

In another aspect, the invention provides an isolated polynucleotide which includes one or more of the SNPs described herein. The polynucleotide can be, *e.g.*, a nucleotide sequence which includes one or more of the polymorphic sequences shown in Table 1 and which includes a polymorphic sequence, or a fragment of the polymorphic sequence, as long
10 as it includes the polymorphic site. The polynucleotide may alternatively contain a nucleotide sequence which includes a sequence complementary to one or more of these sequences, or a fragment of the complementary nucleotide sequence, provided that the fragment includes a polymorphic site in the polymorphic sequence.

The polynucleotide can be, *e.g.*, DNA or RNA, and can be between about 10 and
15 about 100 nucleotides, *e.g.* 10-90, 10-75, 10-51, 10-40, or 10-30, nucleotides in length.

In preferred embodiments, the polymorphic site in the polymorphic sequence includes a nucleotide other than the nucleotide listed in Table 1, column 5 for the polymorphic sequence, *e.g.*, the polymorphic site includes the nucleotide listed in Table 1, column 6 for the polymorphic sequence.

20 In other embodiments, the complement of the polymorphic site includes a nucleotide other than the complement of the nucleotide listed in Table 1, column 5 for the complement of the polymorphic sequence, *e.g.*, the complement of the nucleotide listed in Table 1, column 6 for the polymorphic sequence.

In some embodiments, the polymorphic sequence is associated with a polypeptide
25 related to one of the protein families disclosed herein. For example, the nucleic acid may be associated with a polypeptide related to angiopoietin, 4-hydroxybutyrate dehydrogenase, or any of the other proteins identified in Table 1, column 10.

In another aspect, the invention provides an isolated allele-specific oligonucleotide that hybridizes to a first polynucleotide containing a polymorphic site. The first
30 polynucleotide can be, *e.g.*, a nucleotide sequence comprising one or more polymorphic

sequences recited in Table 1, provided that the polymorphic sequence includes a nucleotide other than the nucleotide recited in Table 1, column 5 for the polymorphic sequence.

Alternatively, the first polynucleotide can be a nucleotide sequence that is a fragment of the polymorphic sequence, provided that the fragment includes a polymorphic site in the

5 polymorphic sequence, or a complementary nucleotide sequence which includes a sequence complementary to one or more polymorphic sequences in Table 1, provided that the complementary nucleotide sequence includes a nucleotide other than the complement of the nucleotide recited in Table 1, column 5. The first polynucleotide may in addition include a nucleotide sequence that is a fragment of the complementary sequence, provided that the
10 fragment includes a polymorphic site in the polymorphic sequence.

In some embodiments, the oligonucleotide does not hybridize under stringent conditions to a second polynucleotide. The second polynucleotide can be, *e.g.*, (a) a nucleotide sequence comprising one or more polymorphic sequences in Table 1, wherein the polymorphic sequence includes the nucleotide listed in Table 1, column 5 for the
15 polymorphic sequence; (b) a nucleotide sequence that is a fragment of any of the polymorphic sequences; (c) a complementary nucleotide sequence including a sequence complementary to one or more polymorphic sequences disclosed herein in Table 1; and (d) a nucleotide sequence that is a fragment of the complementary sequence, provided that the fragment includes a polymorphic site in the polymorphic sequence.

20 The oligonucleotide can be, *e.g.*, between about 10 and about 100 bases in length. In some embodiments, the oligonucleotide is between about 10 and 75 bases, 10 and 51 bases, 10 and about 40 bases, or about 15 and 30 bases in length.

The invention also provides a method of detecting a polymorphic site in a nucleic acid. The method includes contacting the nucleic acid with an oligonucleotide that hybridizes
25 to a polymorphic sequence selected shown in Table 1, or its complement, provided that the polymorphic sequence includes a nucleotide other than the nucleotide recited in Table 1, column 5 for the polymorphic sequence, or the complement includes a nucleotide other than the complement of the nucleotide recited in Table 1, column 5. The method also includes determining whether the nucleic acid and the oligonucleotide hybridize. Hybridization of the
30 oligonucleotide to the nucleic acid sequence indicates the presence of the polymorphic site in the nucleic acid.

In preferred embodiments, the oligonucleotide does not hybridize to the polymorphic sequence when the polymorphic sequence includes the nucleotide recited in Table 1, column 5 for the polymorphic sequence, or when the complement of the polymorphic sequence includes the complement of the nucleotide recited in Table 1, column 5 for the polymorphic sequence.

The oligonucleotide can be, *e.g.*, between about 10 and about 100 bases in length. In some embodiments, the oligonucleotide is between about 10 and 75 bases, 10 and 51 bases, 10 and about 40 bases, or about 15 and 30 bases in length.

In some embodiments, the polymorphic sequence identified by the oligonucleotide is associated with a nucleic acid encoding polypeptide related to one of the protein families disclosed herein. the polymorphic sequence is associated with a polypeptide related to one of the protein families disclosed herein. For example, the nucleic acid may be associated with a polypeptide related to angiopoietin, 4-hydroxybutyrate dehydrogenase, or any of the other proteins identified in Table 1, column 10.

In a further aspect, the invention provides a method of determining the relatedness of a first and second nucleic acid. The method includes providing a first nucleic acid and a second nucleic acid and contacting the first nucleic acid and the second nucleic acid with an oligonucleotide that hybridizes to a polymorphic sequence selected disclosed in Table 1, or its complement, provided that the polymorphic sequence includes a nucleotide other than the nucleotide recited in Table 1, column 5 for the polymorphic sequence, or the complement includes a nucleotide other than the complement of the nucleotide recited in Table 1, column 5. The method also includes determining whether the first nucleic acid and the second nucleic acid hybridize to the oligonucleotide, and comparing hybridization of the first and second nucleic acids to the oligonucleotide. Hybridization of first and second nucleic acids to the nucleic acid indicates the first and second subjects are related.

In preferred embodiments, the oligonucleotide does not hybridize to the polymorphic sequence when the polymorphic sequence includes the nucleotide recited in Table 1, column 5 for the polymorphic sequence, or when the complement of the polymorphic sequence includes the complement of the nucleotide recited in Table 1, column 5 column for the polymorphic sequence.

The oligonucleotide can be, *e.g.*, between about 10 and about 100 bases in length. In some embodiments, the oligonucleotide is between about 10 and 75 bases, 10 and 51 bases, 10 and about 40 bases, or about 15 and 30 bases in length.

The method can be used in a variety of applications. For example, the first nucleic acid may be isolated from physical evidence gathered at a crime scene, and the second nucleic acid may be obtained from a person suspected of having committed the crime. Matching the two nucleic acids using the method can establish whether the physical evidence originated from the person.

In another example, the first sample may be from a human male suspected of being the father of a child and the second sample may be from a child. Establishing a match using the described method can establish whether the male is the father of the child.

In another aspect, the method includes determining if a sequence polymorphism is present in a subject, such as a human. The method includes providing a nucleic acid from the subject and contacting the nucleic acid with an oligonucleotide that hybridizes to a polymorphic sequence disclosed in Table 1, or its complement, provided that the polymorphic sequence includes a nucleotide other than the nucleotide recited in Table 1, column 5 for the polymorphic sequence, or the complement includes a nucleotide other than the complement of the nucleotide recited in Table 1, column 5. Hybridization between the nucleic acid and the oligonucleotide is then determined. Hybridization of the oligonucleotide to the nucleic acid sequence indicates the presence of the polymorphism in said subject.

In another aspect, the invention provides an isolated polypeptide comprising a polymorphic site at one or more amino acid residues, and wherein the protein is encoded by a polynucleotide including one of the polymorphic sequences in Table 1, or their complement, provided that the polymorphic sequence includes a nucleotide other than the nucleotide recited in Table 1, column 5 for the polymorphic sequence, or the complement includes a nucleotide other than the complement of the nucleotide recited in Table 1, column 5.

The polypeptide can be, *e.g.*, related to one of the protein families disclosed herein. For example, polypeptide can be related to angiopoietin, 4-hydroxybutyrate dehydrogenase, ATP-dependent RNA helicase, MHC Class I histocompatibility antigen, or phosphoglycerate kinase.

In some embodiments, the polypeptide is translated in the same open reading frame as is a wild type protein whose amino acid sequence is identical to the amino acid sequence of the polymorphic protein except at the site of the polymorphism.

In some embodiments, the polypeptide encoded by the polymorphic sequence, or its complement, includes the nucleotide listed in Table 1, column 6 for the polymorphic sequence, or the complement includes the complement of the nucleotide listed in Table 1, column 6.

The invention also provides an antibody that binds specifically to a polypeptide encoded by a polynucleotide comprising a nucleotide sequence encoded by a polynucleotide including one or more of the polymorphic sequences in Table 1, or its complement. The polymorphic sequence includes a nucleotide other than the nucleotide recited in Table 1, column 5 for the polymorphic sequence, or the complement includes a nucleotide other than the complement of the nucleotide recited in Table 1, column 5.

In some embodiments, the antibody binds specifically to a polypeptide encoded by a polymorphic sequence which includes the nucleotide listed in Table 1, column 6 for the polymorphic sequence.

Preferably, the antibody does not bind specifically to a polypeptide encoded by a polymorphic sequence which includes the nucleotide listed in Table 1, column 5 for the polymorphic sequence.

The invention further provides a method of detecting the presence of a polypeptide having one or more amino acid residue polymorphisms in a subject. The method includes providing a protein sample from the subject and contacting the sample with the above-described antibody under conditions that allow for the formation of antibody-antigen complexes. The antibody-antigen complexes are then detected. The presence of the complexes indicates the presence of the polypeptide.

The invention also provides a method of treating a subject suffering from, at risk for, or suspected of, suffering from a pathology ascribed to the presence of a sequence polymorphism in a subject, *e.g.*, a human, non-human primate, cat, dog, rat, mouse, cow, pig, goat, or rabbit. The method includes providing a subject suffering from a pathology associated with aberrant expression of a first nucleic acid comprising a polymorphic sequence

shown in Table 1, or its complement, and treating the subject by administering to the subject an effective dose of a therapeutic agent. Aberrant expression can include qualitative alterations in expression of a gene, *e.g.*, expression of a gene encoding a polypeptide having an altered amino acid sequence with respect to its wild-type counterpart. Qualitatively
5 different polypeptides can include, shorter, longer, or altered polypeptides relative to the amino acid sequence of the wild-type polypeptide. Aberrant expression can also include quantitative alterations in expression of a gene. Examples of quantitative alterations in gene expression include lower or higher levels of expression of the gene relative to its wild-type counterpart, or alterations in the temporal or tissue-specific expression pattern of a gene.
10 Finally, aberrant expression may also include a combination of qualitative and quantitative alterations in gene expression.

The therapeutic agent can include, *e.g.*, second nucleic acid comprising the polymorphic sequence, provided that the second nucleic acid comprises the nucleotide present in the wild type allele. In some embodiments, the second nucleic acid sequence
15 comprises a polymorphic sequence which includes nucleotide listed in Table 1, column 5 for the polymorphic sequence.

Alternatively, the therapeutic agent can be a polypeptide encoded by a polynucleotide comprising polymorphic sequence shown in Table 1, or by a polynucleotide comprising a nucleotide sequence that is complementary to any one of the polymorphic sequences,
20 provided that the polymorphic sequence includes the nucleotide listed in Table 1, column 6 for the polymorphic sequence.

The therapeutic agent may further include an antibody as herein described, or an oligonucleotide comprising a polymorphic sequence shown in Table 1, or by a polynucleotide comprising a nucleotide sequence that is complementary to any one the
25 polymorphic sequences, provided that the polymorphic sequence includes the nucleotide listed in Table 1, column 6 for the polymorphic sequence,

In another aspect, the invention provides an oligonucleotide array comprising one or more oligonucleotides hybridizing to a first polynucleotide at a polymorphic site encompassed therein. The first polynucleotide can be, *e.g.*, a nucleotide sequence comprising
30 one or more polymorphic sequences shown in Table 1; a nucleotide sequence that is a fragment of any of the nucleotide sequence, provided that the fragment includes a

polymorphic site in the polymorphic sequence; a complementary nucleotide sequence comprising a sequence complementary to one or more of the polymorphic sequences; or a nucleotide sequence that is a fragment of the complementary sequence, provided that the fragment includes a polymorphic site in the polymorphic sequence.

5 In preferred embodiments, the array comprises 10; 100; 1,000; 10,000; 100,000 or more oligonucleotides.

The invention also provides a kit comprising one or more of the herein-described nucleic acids. The kit can include, *e.g.*, polynucleotide which includes one or more of the SNPs described herein. The polynucleotide can be, *e.g.*, a nucleotide sequence which
10 includes one or more of the polymorphic sequences shown in Table 1, and which includes a polymorphic sequence, or a fragment of the polymorphic sequence, as long as it includes the polymorphic site. The polynucleotide may alternatively contain a nucleotide sequence which includes a sequence complementary to one or more of the sequences, or a fragment of the complementary nucleotide sequence, provided that the fragment includes a polymorphic
15 site in the polymorphic sequence.

Alternatively, or in addition, the kit can include the invention provides an isolated allele-specific oligonucleotide that hybridizes to a first polynucleotide containing a polymorphic site. The first polynucleotide can be, *e.g.*, a nucleotide sequence comprising one or more polymorphic sequences shown in Table 1, provided that the polymorphic sequence
20 includes a nucleotide other than the nucleotide recited in Table 1, column 5 for the polymorphic sequence. Alternatively, the first polynucleotide can be a nucleotide sequence that is a fragment of the polymorphic sequence, provided that the fragment includes a polymorphic site in the polymorphic sequence, or a complementary nucleotide sequence which includes a sequence complementary to one or more polymorphic sequences shown in
25 Table 1, provided that the complementary nucleotide sequence includes a nucleotide other than the complement of the nucleotide recited in Table 1, column 6. The first polynucleotide may in addition include a nucleotide sequence that is a fragment of the complementary sequence, provided that the fragment includes a polymorphic site in the polymorphic sequence.

BRIEF DESCRIPTION OF THE DRAWING

FIG. 1 illustrates an example of the way in which SNP sites were identified in the present invention.

Unless otherwise defined, all technical and scientific terms used herein have the same
5 meaning as commonly understood by one of ordinary skill in the art to which this invention
belongs. Although methods and materials similar or equivalent to those described herein can
be used in the practice or testing of the present invention, suitable methods and materials are
described below. All publications, patent applications, patents, and other references
mentioned herein are incorporated by reference in their entirety. In the case of conflict, the
10 present specification, including definitions, will control. In addition, the materials, methods,
and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the following
detailed description and claims.

DETAILED DESCRIPTION OF THE INVENTION

15 The invention provides human SNPs in sequences which are transcribed, *i.e.*, are
cSNPs. Many SNPs have been identified in genes related to polypeptides of known function.
If desired, SNPs associated with various polypeptides can be used together. For example,
SNPs can be grouped according to whether they are derived from a nucleic acid encoding a
polypeptide related to particular protein family or involved in a particular function.
20 Similarly, SNPs can be grouped according to the functions played by their gene products.
Such functions include, structural proteins, proteins from which associated with metabolic
pathways fatty acid metabolism, glycolysis, intermediary metabolism, calcium metabolism,
proteases, and amino acid metabolism, etc. Specifically, the present invention provides a
large number of human cSNP's based on at least one gene product that has not been
25 previously identified. In contrast, and as defined specifically in the following paragraph, the
cSNP's involve nucleic acid sequences that are assembled from at least one known sequence.

The present invention describes 651 distinct polymorphic sites, which are summarized
in Table 1. Raw traces underlying sequence data were drawn from public databases and from
the proprietary database of the Assignee of the present invention. The sequences were
30 obtained by calling the bases from these traces, and included assigning "Phred" quality scores

for each called base. For each allelic set, at the polynucleotide level, four or more nucleotide sequences were identified having at least partial overlap with one another.

As illustrated in FIG. 1, these four or more sequences could be clustered and assembled to make a consensus contig that included an ORF. In this way, the inventors
5 found that the assembled contigs defined associated sets of two, or possibly more than two, alleles defined by a SNP at a particular polymorphic site. In order to be confirmed as a SNP site, the nucleotide change from the consensus sequence had to occur in at least two individual sequences, and had to have a "Phred" score of 23 or higher at the site of the presumed SNP. Furthermore, in a window of 5 bases on either side of the SNP, no more than
10 50% mismatching with the consensus sequence was allowed. In the assembly leading to each of the contigs defining the allelic set, the SNP alleles occur in polynucleotides found in public databases.

It was found that the assembled contigs defined associated sets of two, or possibly more than two, alleles defined by an SNP at a particular polymorphic site. These associations
15 were not previously known.

At the level of translation of an ORF contained in the contigs, allelic sets were identified in which one allele defines a known polypeptide sequence that includes the polymorphic site and another polypeptide allele is not previously known. Then, various associations of alleles are possible. For example, it is possible that an allelic pair is defined
20 in a noncoding region of the contig containing an ORF. In such cases the inventors believe that the invention resides in the recognition of the allelic pair; this association has not heretofore been made.

Alternatively, sets of allelic contigs may exist in which the polymorphic site is within an ORF, but does not result in an amino acid change among the allelic polypeptides. Thus, in
25 another embodiment, the polymorphic site resides within an ORF and results in an amino acid change, or a frameshift, among the alleles of the allelic set. In the sets of gene products that fall within this group, at least one of the alleles at the polypeptide level is a known protein. At least one of the remaining allele or alleles in the set, carrying a variant amino acid at the polymorphic site, is a novel polypeptide not heretofore known. The invention resides
30 at least in the recognition of the polymorphic allele as being a variant of the known reference polypeptide.

Table 1 provides information concerning the allelic sequences. One of the sequences may be termed a reference polymorphic sequence, and the corresponding second sequence includes the variant SNP at the polymorphic site. Since the reference polypeptide sequence is already known, the Sequence Listing accompanying this application provides only the sequence of the polymorphic allele, while its SEQ ID NO is provided in the Table. A reference to the SEQ ID NO that corresponds to the translated amino acid sequence is also given. The Table includes thirteen columns that provide descriptive information for each cSNP, each of which occupies one row in the Table. The column headings, and a description of each, are given below.

SNPs disclosed in Table 1 were detected by aligning large numbers of sequences from genetically diverse sources of publicly available mRNA libraries (Clontech). Software designed specifically to look for multiple examples of variant bases differing from a consensus sequence was created and deployed. A criteria of a minimum of 2 occurrences of a sequence differing from the consensus in high quality sequence reads was used to identify an SNP.

The SNPs described herein may be useful in diagnostic kits, for DNA arrays on chips and for other uses that involve hybridization of the SNP.

Specific SNPs may have utility where a disease has already been associated with that gene. Examples of possible disease correlations between the claimed SNPs with members of the genes of each classification are listed below:

Amylases

Amylase is responsible for endohydrolysis of 1,4-alpha-glucosidic linkages in oligosaccharides and polysaccharides. Variations in amylase gene may be indicative of delayed maturation and of various amylase producing neoplasms and carcinomas.

Amyloid

The serum amyloid A (SAA) proteins comprise a family of vertebrate proteins that associate predominantly with high density lipoproteins (HDL). The synthesis of certain members of the family is greatly increased in inflammation. Prolonged elevation of plasma SAA levels, as in chronic inflammation, results in a pathological condition, called amyloidosis, which affects the liver, kidney and spleen and which is characterized by the

highly insoluble accumulation of SAA in these tissues. Amyloid selectively inhibits insulin-stimulated glucose utilization and glycogen deposition in muscle, while not affecting adipocyte glucose metabolism. Deposition of fibrillar amyloid proteins intraneuronally, as neurofibrillary tangles, extracellularly, as plaques and in blood vessels, is characteristic of both Alzheimer's disease and aged Down's syndrome. Amyloid deposition is also associated with type II diabetes mellitus.

Angiopoeitin

Members of the angiopoetin/fibrinogen family have been shown to stimulate the generation of new blood vessels, inhibit the generation of new blood vessels, and perform several roles in blood clotting. This generation of new blood vessels, called angiogenesis, is also an essential step in tumor growth in order for the tumor to get the blood supply it needs to expand. Variation in these genes may be predictive of any form of heart disease, numerous blood clotting disorders, stroke, hypertension and predisposition to tumor formation and metastasis. In particular, these variants may be predictive of the response to various antihypertensive drugs and chemotherapeutic and anti-tumor agents.

Apoptosis-related proteins

Active cell suicide (apoptosis) is induced by events such as growth factor withdrawal and toxins. It is controlled by regulators, which have either an inhibitory effect on programmed cell death (anti-apoptotic) or block the protective effect of inhibitors (pro-apoptotic). Many viruses have found a way of countering defensive apoptosis by encoding their own anti-apoptosis genes preventing their target-cells from dying too soon. Variants of apoptosis related genes may be useful in formulation of antiaging drugs.

Cadherin, Cyclin, Polymerase, Oncogenes, Histones, Kinases

Members of the cell division/cell cycle pathways such as cyclins, many transcription factors and kinases, DNA polymerases, histones, helicases and other oncogenes play a critical role in carcinogenesis where the uncontrolled proliferation of cells leads to tumor formation and eventually metastasis. Variation in these genes may be predictive of predisposition to any form of cancer, from increased risk of tumor formation to increased rate of metastasis. In particular, these variants may be predictive of the response to various chemotherapeutic and anti-tumor agents.

Colony-stimulating factor-related proteins

Granulocyte/macrophage colony-stimulating factors are cytokines that act in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and the monocytes-macrophages.

5 Complement-related proteins

Complement proteins are immune associated cytotoxic agents, acting in a chain reaction to exterminate target cells to that were opsonized (primed) with antibodies, by forming a membrane attack complex (MAC). The mechanism of killing is by opening pores in the target cell membrane. Variations in 20 complement genes or their inhibitors are associated with many autoimmune disorders. Modified serum levels of complement products cause edemas of various tissues, lupus (SLE), vasculitis, glomerulonephritis, renal failure, hemolytic anemia, thrombocytopenia, and arthritis. They interfere with mechanisms of ADCC (antibody dependent cell cytotoxicity), severely impair immune competence and reduce phagocytic ability. Variants of complement genes may also be indicative of type I diabetes mellitus, meningitis neurological disorders such as Nemaline myopathy, Neonatal hypotonia, muscular disorders such as congenital myopathy and other diseases.

Cytochrome

The respiratory chain is a key biochemical pathway which is essential to all aerobic cells. There are five different cytochromes involved in the chain. These are heme bound proteins which serve as electron carriers. Modifications in these genes may be predictive of ataxia areflexia, dementia and myopathic and neuropathic changes in muscles. Also, association with various types of solid tumors.

Kinesins

Kinesins are tubulin molecular motors that function to transport organelles within cells and to move chromosomes along microtubules during cell division. Modifications of these genes may be indicative of neurological disorders such as Pick disease of the brain, tuberous sclerosis.

Cytokines, Interferon, Interleukin

Members of the cytokine families are known for their potent ability to stimulate cell growth and division even at low concentrations. Cytokines such as erythropoietin are cell-specific in their growth stimulation; erythropoietin is useful for the stimulation of the proliferation of erythroblasts. Variants in cytokines may be predictive for a wide variety of diseases, including cancer predisposition.

G-protein coupled receptors

G-protein coupled receptors (also called R7G) are an extensive group of hormones, neurotransmitters, odorants and light receptors which transduce extracellular signals by interaction with guanine nucleotide-binding (G) proteins. Alterations in genes coding for G-coupled proteins may be involved in and indicative of a vast number of physiological conditions. These include blood pressure regulation, renal dysfunctions, male infertility, dopamine associated cognitive, emotional, and endocrine functions, hypercalcemia, chondrodysplasia and osteoporosis, pseudohypoparathyroidism, growth retardation and dwarfism.

Thioesterases

Eukaryotic thiol proteases are a family of proteolytic enzymes which contain an active site cysteine. Catalysis proceeds through a thioester intermediate and is facilitated by a nearby histidine side chain; an asparagine completes the essential catalytic triad. Variants of thioester associated genes may be predictive of neuronal disorders and mental illnesses such as Ceroid Lipoffiscinosis, Neuronal 1, Infantile, Santavuori disease and more.

Breakdown Classifications of SNPS

The following list describes the numerical breakdown by molecule type of the SNPs described in Table 1. The key to these molecule types is as follows.

25

30

TPase_associated:	864
Guanylyl:	3
MHC:	1077
amylase:	44
amylaseinhib:	1
amyloid:	96
apoptosis:	91

	apoptosisinhib:	29
	apoptosisrecep:	14
	biotindep:	29
	cadhenn:	415
5	calcium_channel:	85
	carboxylase:	4
	cathepsin:	336
	cathepsininhib:	41
	chloride_channel:	90
10	collagen:	1542
	complement:	222
	complementinhib:	21
	complementrecept:	10
	csf:	31
15	csf recept:	37
	cyclin:	65
	cyto45O:	136
	cytochrome:	659
	deaminase:	44
20	dehydrogenase:	1235
	desaturase:	9
	dna_rna_bind:	1309
	dna_rna_bind_inhib:	16
	dynein:	108
25	elastase:	134
	elastaseinhib:	6
	eph:	487
	esterase:	258
	esteraseinhib:	3
30	fgf:	34
	fgf receptor:	12
	gaba:	45
	glucoamylase:	106
	glucuronidase:	14
35	glycoprotein:	3176
	helicase:	333
	histone:	272
	homeobox:	431
	hydrolase:	187
40	hydroxysteroid:	84
	hypoxanthine:	4
	immunoglob:	1106
	immunoglob_recept:	19
	interferon:	322
45	interleukin:	88
	interleukinrecept:	126
	isomerase:	404
	isomeraseinhibitor:	45
	isomerasereceptor:	4
50	kinase:	1684

	kinase inhibitor:	187
	kinase receptor:	233
	kinesin:	86
	laminin:	196
5	lipase:	63
	metallothionein:	62
	misc_channel:	215
	ngf:	30
	nucl_recpt:	339
10	nuclease:	298
	oncogene:	783
	oxidase:	128
	oxygenase:	14
	peptidase:	150
15	peroxidase:	115
	phosphatase:	668
	phosphataseinhib:	71
	phosphorylase:	84
	polymerase:	489
20	potassium_channel:	43
	prostaglandin:	55
	protease:	954
	proteaseinhib:	271
	reductase:	243
25	ribosomal prot:	1040
	struct:	3128
	sulfotransferase:	42
	synthase:	893
	tgf:	117
30	tgfreceptor:	41
	thioesterase:	3
	thiolase:	38
	tm7:	453
	tnf:	151
35	tnfreceptor:	36
	traffic:	22
	transcriptfactor:	1139
	transferase:	291
	transport:	900
40	tubulin:	334
	ubiquitin:	229
	water_channel:	18
	unclassified:	10567

The key to the molecule type is as follows:

	Abbrev:	Title:
5	amylase	amylase protein
	amylaseinhib	amylase inhibitor
	amyloid	amyloid protein
	apoptosis	apoptosis associated protein
	apoptosisinhib	apoptosis inhibitors
10	apoptosisrecep	apoptosis receptors
	ATPase_associated	ATPase associated protein
	biotindep	biotin dependent enzyme/protein
	cadherin	cadherin protein
	calcium_channel	calcium channel protein
15	carboxylase	carboxylase protein
	cathepsin	cathepsin/carboxypeptidases
	cathepsininhib	cathepsin/carboxypeptidase inhibitor
	chloride_channel	chloride channel protein
	collagen	collagen
20	complement	complement protein
	complementrecept	complement receptor protein
	complementinhib	complement inhibitor
	csf	colony stimulating factor
	csfrecept	colony stimulating factor receptor
25	cyclin	cyclin protein
	cyto450	cytochrome p450 protein
	cytochrome	cytochrome related protein
	deaminase	deaminase
	dehydrogenase	dehydrogenase
30	desaturase	desaturase
	dna_ma_bind	DNA/RNA binding protein/factor
	dna_ma_inhib	DNA/RNA binding protein/factor inhibitor
	dynein	dynein
35	elastase	elastase
	elastaseinhib	elastase inhibitor
	eph	EPH family of tyrosine kinases
	esterase	esterase
	esteraseinhib	esterase inhibitor
40	fgf	fibroblast growth factor
	fgfreceptor	fibroblast growth factor receptor
	gaba	GABA receptor
	glucoamylase	glucoamylase
	glucoronidase	glucoronidase
45	glycoprotein	glycoprotein
	Guanylyl	guanylylate cyclase
	helicase	helicase
	histone	histone
	HOM	homologous

	homeobox	homeobox protein
	hydrolase	hydrolase
	hydroxysteroid	hydroxysteroid associated protein
	hypoxanthine	hypoxanthine associated protein
5	immunoglob	immunoglobulin
	immunoglobrecept	immunoglobulin receptor
	interferon	interferon
	interleukin	interleukin
	interleukinrecept	interleukin receptor
10	isomerase	isomerase
	isomeraseinhibitor	isomerase inhibitor
	isomerasereceptor	isomerase receptor
	kinase	kinase
	kinaseinhibitor	kinase inhibitor
15	kinasereceptor	kinase receptor
	kinesin	kinesin
	laminin	laminin associated protein
	lipase	lipase
	metallothionein	metallothionein
20	MHC	major histocompatibility complex
	misc_channel	miscellaneous channel
	ngf	nerve growth factor
	nuci_recpt	nuclear receptor
	nuclease	nuclease
25	oncogene	oncogene associated protein
	oxidase	oxidase
	oxygenase	oxygenase
	peptidase	peptidase
	peroxidase	peroxidase
30	phosphatase	phosphatase
	phosphataseinhib	phosphatase inhibitor
	phosphorylase	phosphorylase
	PIR	PIR DATABASE (release 56, 29-OCT-1998)
35	polymerase	polymerase
	potassium_channel	potassium channel protein
	prostaglandin	prostaglandin
	protease	protease
	proteaseinhib	protease inhibitor
40	reductase	reductase
	ribosomalprot	ribosomal associated protein
	RTR	EMBLDATABASE translated entries not to be incorporated into SWISS- PROT (20-JUL-1998)
45	SIM	similar
	SPTR	EMBL DATABASE translated entries to be incorporated into SWISS-PROT (20- JUL-1998)
	struct	structural associated protein
50	sulfotransferase	sulfotransferase

	SWP	SWISS-PROT DATABASE (release 18-OCT-1998)
	SWPN	SWISS-PROT Update (release 11-NOV-98)
5	synthase	synthase
	tgf	transforming growth factor
	tgfreceptor	transforming growth factor receptor
	thioesterase	thioesterase
	thiolase	thiolase
10	tm7	seven transmembrane domain G-protein coupled receptor
	tnf	necrosis factor receptor
	traffic	tumor necrosis factor
	tnfreceptor	tumor trafficking associated protein
15	TRN	EMBL DATABASE translated entries update (20-JUL-1998)
	transcriptfactor	transcription factor
	transferase	transferase
	transport	transport protein
20	tubulin	tubulin
	ubiquitin	ubiquitin
	unclassified	Protein not categorized into one of the aforementioned protein families
	water channel	water channel protein
25		

Table 1

A compilation of polymorphisms is listed in Table 1. Table 1 includes thirteen columns that provide descriptive information for each cSNP, each of which occupies one row in the Table. The column headings, and an explanation for each, are given below.

30 The first column of the table lists the names assigned to the fragments in which the polymorphisms occur. The fragments are all human genomic fragments. The sequence of one allelic form of each of the fragments (arbitrarily referred to as the prototypical or reference form) has been previously published. These sequences are listed at <http://www-genome.wi.mit.edu/> (all STS's sequence tag sites)); <http://shgc.stanford.edu> (Stanford STS's); and <http://www.tigr.org/> (TIGR STS's). The web sites also list primers for

35 amplification of the fragments, and the genomic location of the fragments. Some fragments are expressed sequence tags, and some are random genomic fragments. All information in the web sites concerning the fragments listed in the table is incorporated by reference in its entirety for all purposes.

The second column lists the position in the fragment in which a polymorphic site has been found. Positions are numbered consecutively with the first base of the fragment sequence listed as in one of the above databases being assigned the number one. The third column lists the base occupying the polymorphic site in the sequence in the data base. This base is arbitrarily designated the reference or prototypical form, but it is not necessarily the most frequently occurring form. The fourth column in the table lists the alternative base(s) at the polymorphic site. The fifth column of the table lists a 5' (upstream or forward) primer that hybridizes with the 5' end of the DNA sequence to be amplified. The sixth column of the table lists a 3' (downstream or reverse) primer that hybridizes with the complement of the 3' end of the sequence to be amplified. The seventh column of the table lists a number of bases of sequence on either side of the polymorphic site in each fragment. The indicated sequences can either be DNA or RNA. In the latter, the T's shown in the table are replaced by U's. The base occupying the polymorphic site is indicated in EUT'AC-IUB ambiguity code.

"SEQ ID" provides the cross-references to the two nucleotide SEQ ID NOS: for the cognate pair, which are numbered consecutively, and, as explained below, amino acid SEQ ID NOS: as well, in the Sequence Listing of the application.

Each sequence entry in the Sequence Listing also includes a cross-reference to the CuraGen sequence ID, under the label "Accession number". The first pair of SEQ ID NOS: given in the first column of each row of the Table is the SEQ ID NO: identifying the nucleic acid sequence for the polymorphism. If a polymorphism carries an entry for the amino acid portion of the row, a third SEQ ID NO: appears in parentheses in the column "Amino acid before" (see below) for the reference amino acid sequence, and a fourth SEQ ID NO: appears in parentheses in the column "Amino acid after" (see below) for the polymorphic amino acid sequence. The latter SEQ ID NOS: refer to amino acid sequences giving the cognate reference and polymorphic amino acid sequences that are the translation of the nucleotide polymorphism. If a polymorphism carries no entry for the protein portion of the row, only one pair SEQ ID NOS: is provided, in the first column.

"CuraGen sequence ID" provides CuraGen Corporation's accession number.

"Base pos. of SNP" gives the numerical position of the nucleotide in the nucleic acid at which the cSNP is found, as identified in this invention.

“Polymorphic sequence” provides a 51-base sequence with the polymorphic site at the 26th base in the sequence, as well as 25 bases from the reference sequence on the 5’ side and the 3’ side of the polymorphic site. The designation at the polymorphic site is enclosed in square brackets, and provides first, the reference nucleotide; second, a “slash (/)”; and
5 third, the polymorphic nucleotide. In certain cases the polymorphism is an insertion or a deletion. In that case, the position that is “unfilled” (i.e., the reference or the polymorphic position) is indicated by the word “gap”.

“Base before” provides the nucleotide present in the reference sequence at the position at which the polymorphism is found.

10 “Base after” provides the altered nucleotide at the position of the polymorphism.

“Amino acid before” provides the amino acid in the reference protein, if the polymorphism occurs in a coding region. This column also includes the SEQ ID NO: in parentheses for the translated reference amino acid sequence if the polymorphism occurs in a coding region.

15 “Amino acid after” provides the amino acid in the polymorphic protein, if the polymorphism occurs in a coding region. This column also includes the SEQ ID NO in parentheses for the translated polymorphic amino acid sequence if the polymorphism occurs in a coding region.

“Type of change” provides information on the nature of the polymorphism.

20 “SILENT-NONCODING” is used if the polymorphism occurs in a noncoding region of a nucleic acid. “SILENT-CODING” is used if the polymorphism occurs in a coding region of a nucleic acid of a nucleic acid and results in no change of amino acid in the translated polymorphic protein. “CONSERVATIVE” is used if the polymorphism occurs in a coding region of a nucleic acid and provides a change in which the altered amino acid falls in the
25 same class as the reference amino acid. The classes are: 1) Aliphatic: Gly, Ala, Val, Leu, Ile; 2) Aromatic: Phe, Tyr, Trp; 3) Sulfur-containing: Cys, Met; 4) Aliphatic OH: Ser, Thr; 5) Basic: Lys, Arg, His; 6) Acidic: Asp, Glu, Asn, Gln; 7) Pro falls in none of the other classes; and 8) End defines a termination codon.

"NONCONSERVATIVE" is used if the polymorphism occurs in a coding region of a nucleic acid and provides a change in which the altered amino acid falls in a different class than the reference amino acid.

5 "FRAMESHIFT" relates to an insertion or a deletion. If the frameshift occurs in a coding region, the Table provides the translation of the frameshifted codons 3' to the polymorphic site.

"Protein classification of CuraGen gene" provides a generic class into which the protein is classified. Multiple classes of proteins were identified as listed above in the discussion of Table 1.

10 "Name of protein identified following a BLASTX analysis of the CuraGen sequence" provides the database reference for the protein found to resemble the novel reference-polymorphism cognate pair most closely.

"Similarity (pvalue) following a BLASTX analysis" provides the pvalue, a statistical measure from the BLASTX analysis that the polymorphic sequence is similar to, and
15 therefore an allele of, the reference, or wild-type, sequence. In the present application, a cutoff of $pvalue > 1 \times 10^{-50}$ (entered, for example, as 1.0E-50 in the Table) is used to establish that the reference-polymorphic cognate pairs are novel. A $pvalue < 1 \times 10^{-50}$ defines proteins considered to be already known.

"Map location" provides any information available at the time of filing related to
20 localization of a gene on a chromosome.

The polymorphisms are arranged in Table 1 in the following order:

SEQ ID NOS: 1-422 are nucleotide sequences for SNPs that are silent.

SEQ ID NOS: 423-480 are nucleotide sequences for SNPs that lead to conservative amino acid changes.

25 SEQ ID NOS: 481-619 are nucleotide sequences for SNPs that lead to nonconservative amino acid changes.

SEQ ID NOS: 620-651 are nucleotide sequences for SNPs that involve a gap. With respect to the reference or wild-type sequence at the position of the polymorphism, the allelic

cSNP introduces an additional nucleotide (an insertion) or deletes a nucleotide (a deletion). An SNP that involves a gap generates a frame shift.

Also presented in the sequence listing filed herewith are predicted amino acid sequences encoded by the polymorphic sequences shown in Table 1.

5 SEQ ID NOs: 652-709 are the amino acid sequences centered at the polymorphic amino acid residue for the protein products provided by SNPs that lead to conservative amino acid changes. 7 or 8 amino acids on either side of the polymorphic site are shown. The order in which these sequences appear mirrors the order of presentation of the cognate nucleotide sequences, and is set forth in the Table.

10 SEQ ID NOs: 710-848 are the amino acid sequences centered at the polymorphic amino acid residue for the protein products provided by SNPs that lead to nonconservative amino acid changes. 7 or 8 amino acids on either side of the polymorphic site are shown. The order in which these sequences appear mirrors the order of presentation of the cognate nucleotide sequences, and is set forth in the Table.

15 SEQ ID NOs: 849-880 are the amino acid sequences centered at the polymorphic amino acid residue for the protein products provided by SNPs that lead to frameshift-induced amino acid changes. 7 or 8 amino acids on either side of the polymorphic site are shown. The order in which these sequences appear mirrors the order of presentation of the cognate nucleotide sequences, and is set forth in the Table.

20 Provided herein are compositions which include, or are capable of detecting, nucleic acid sequences having these polymorphisms, as well as methods of using nucleic acids.

Identification of Individuals Carrying SNPs

Individuals carrying polymorphic alleles of the invention may be detected at either the DNA, the RNA, or the protein level using a variety of techniques that are well known in the art. Strategies for identification and detection are described in *e.g.*, EP 730,663, EP 717,113, and PCT US97/02102. The present methods usually employ pre-characterized polymorphisms. That is, the genotyping location and nature of polymorphic forms present at a site have already been determined. The availability of this information allows sets of probes to be designed for specific identification of the known polymorphic forms.

25

Many of the methods described below require amplification of DNA from target samples. This can be accomplished by e.g., PCR. (1989), B. for detecting polymorphisms. See generally PCR Technology: Principles and Applications for DNA Amplification (ed. H.A. Erlich, Freeman Press, NY, NY, 1992); PCR Protocols: A Guide to Methods and Applications (eds. Innis, et al., Academic Press, San Diego, CA, 1990); Mattila et al.,
5 Nucleic Acids Res. 19, 4967 (1991); Eckert et al., PCR Methods and Applications 1, 17 (1991); PCR (eds. McPherson et al., IRL Press, Oxford); and U.S. Patent 4,683,202.

The phrase "recombinant protein" or "recombinantly produced protein" refers to a peptide or protein produced using non-native cells that do not have an endogenous copy of
10 DNA able to express the protein. In particular, as used herein, a recombinantly produced protein relates to the gene product of a polymorphic allele, i.e., a "polymorphic protein" containing an altered amino acid at the site of translation of the nucleotide polymorphism. The cells produce the protein because they have been genetically altered by the introduction of the appropriate nucleic acid sequence. The recombinant protein will not be found in
15 association with proteins and other subcellular components normally associated with the cells producing the protein. The terms "protein" and "polypeptide" are used interchangeably herein.

The phrase "substantially purified" or "isolated" when referring to a nucleic acid, peptide or protein, means that the chemical composition is in a milieu containing fewer, or
20 preferably, essentially none, of other cellular components with which it is naturally associated. Thus, the phrase "isolated" or "substantially pure" refers to nucleic acid preparations that lack at least one protein or nucleic acid normally associated with the nucleic acid in a host cell. It is preferably in a homogeneous state although it can be in either a dry or aqueous solution. Purity and homogeneity are typically determined using analytical
25 chemistry techniques such as gel electrophoresis or high performance liquid chromatography. Generally, a substantially purified or isolated nucleic acid or protein will comprise more than 80% of all macromolecular species present in the preparation. Preferably, the nucleic acid or protein is purified to represent greater than 90% of all macromolecular species present. More preferably the nucleic acid or protein is purified to greater than 95%, and most preferably the
30 nucleic acid or protein is purified to essential homogeneity, wherein other macromolecular species are not detected by conventional analytical procedures.

The genomic DNA used for the diagnosis may be obtained from any nucleated cells of the body, such as those present in peripheral blood, urine, saliva, buccal samples, surgical specimen, and autopsy specimens. The DNA may be used directly or may be amplified enzymatically in vitro through use of PCR (Saiki et al. Science 239:487-491 (1988)) or other
5 in vitro amplification methods such as the ligase chain reaction (LCR) (Wu and Wallace Genomics 4:560-569 (1989)), strand displacement amplification (SDA) (Walker et al. Proc. Natl. Acad. Sci. U.S.A. 89:392-396 (1992)), self-sustained sequence replication (3SR) (Fahy et al. PCR Methods P&J 1:25-33 (1992)), prior to mutation analysis.

The method for preparing nucleic acids in a form that is suitable for mutation
10 detection is well known in the art. A "nucleic acid" is a deoxyribonucleotide or ribonucleotide polymer in either single- or double-stranded form, including known analogs of natural nucleotides unless otherwise indicated. The term "nucleic acids", as used herein, refers to either DNA or RNA. "Nucleic acid sequence" or "polynucleotide sequence" refers to a single-stranded sequence of deoxyribonucleotide or ribonucleotide bases read from the 5'
15 end to the 3' end. The direction of 5' to 3' addition of nascent RNA transcripts is referred to as the transcription direction; sequence regions on the DNA strand having the same sequence as the RNA and which are beyond the 5' end of the RNA transcript in the 5' direction are referred to as "upstream sequences"; sequence regions on the DNA strand having the same sequence as the RNA and which are beyond the 3' end of the RNA transcript in the 3'
20 direction are referred to as "downstream sequences". The term includes both self-replicating plasmids, infectious polymers of DNA or RNA and nonfunctional DNA or RNA. The complement of any nucleic acid sequence of the invention is understood to be included in the definition of that sequence. "Nucleic acid probes" may be DNA or RNA fragments.

The detection of polymorphisms in specific DNA sequences, can be accomplished by
25 a variety of methods including, but not limited to, restriction-fragment-length-polymorphism detection based on allele-specific restriction-endonuclease cleavage (Kan and Dozy Lancet ii:910-912 (1978)), hybridization with allele-specific oligonucleotide probes (Wallace et al. Nucl. Acids Res. 6:3543-3557 (1978)), including immobilized oligonucleotides (Saiki et al. Proc. Natl. Acad. Sci. USA, 86:6230-6234 (1969)) or oligonucleotide arrays (Maskos and
30 Southern Nucl. Acids Res 21:2269-2270 (1993)), allele-specific PCR (Newton et al. Nucl Acids Res 17:2503-2516 (1989)), mismatch-repair detection (MRD) (Faham and Cox Genome Res 5:474-482 (1995)), binding of MutS protein (Wagner et al. Nucl Acids Res 23:3944-3948 (1995)), denaturing-gradient gel electrophoresis (DGGE) (Fisher and Lerman et

- al. *Proc. Natl. Acad. Sci. U.S.A.* 80:1579-1583 (1983)), single-strand-conformation-polymorphism detection (Orita et al. *Genomics* 5:874-879 (1983)), RNAase cleavage at mismatched base-pairs (Myers et al. *Science* 230:1242 (1985)), chemical (Cotton et al. *Proc. Natl. w Sci. U.S.A.*, 8Z4397-4401 (1988)) or enzymatic (Youil et al. *Proc. Natl. Acad. Sci. U.S.A.* 92:87-91 (1995)) cleavage of heteroduplex DNA, methods based on allele specific primer extension (Syvanen et al. *Genomics* 8:684-692 (1990)), genetic bit analysis (GBA) (Nikiforov et al. &I *Acids* 22:4167-4175 (1994)), the oligonucleotide-ligation assay (OLA) (Landegren et al. *Science* 241:1077 (1988)), the allele-specific ligation chain reaction (LCR) (Barrany *Proc. Natl. Acad. Sci. U.S.A.* 88:189-193 (1991)), gap-LCR (Abravaya et al. *Nucl Acids Res* 23:675-682 (1995)), radioactive and/or fluorescent DNA sequencing using standard procedures well known in the art, and peptide nucleic acid (PNA) assays (Orum et al., *Nucl. Acids Res*, 21:5332-5356 (1993); Thiede et al., *Nucl. Acids Res.* 24:983-984 (1996)).

“Specific hybridization” or “selective hybridization” refers to the binding, or duplexing, of a nucleic acid molecule only to a second particular nucleotide sequence to which the nucleic acid is complementary, under suitably stringent conditions when that sequence is present in a complex mixture (e.g., total cellular DNA or RNA). “Stringent conditions” are conditions under which a probe will hybridize to its target subsequence, but to no other sequences. Stringent conditions are sequence-dependent and are different in different circumstances. Longer sequences hybridize specifically at higher temperatures than shorter ones. Generally, stringent conditions are selected such that the temperature is about 5°C lower than the thermal melting point (T_m) for the specific sequence to which hybridization is intended to occur at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength, pH, and nucleic acid concentration) at which 50% of the target sequence hybridizes to the complementary probe at equilibrium. Typically, stringent conditions include a salt concentration of at least about 0.01 to about 1.0 M Na ion concentration (or other salts), at pH 7.0 to 8.3. The temperature is at least about 30°C for short probes (e.g., 10 to 50 nucleotides). Stringent conditions can also be achieved with the addition of destabilizing agents such as formamide. For example, conditions of 5X SSPE (750 mM NaCl, 50 mM NaPhosphate, 5 mM EDTA, pH 7.4) and a temperature of 25-30°C are suitable for allele-specific probe hybridizations.

“Complementary” or “target” nucleic acid sequences refer to those nucleic acid sequences which selectively hybridize to a nucleic acid probe. Proper annealing conditions

depend, for example, upon a probe's length, base composition, and the number of mismatches and their position on the probe, and must often be determined empirically. For discussions of nucleic acid probe design and annealing conditions, see, for example, Sambrook et al., or Current Protocols in Molecular Biology, F. Ausubel *et al.*, ed., Greene
5 Publishing and Wiley-Interscience, New York (1987).

A perfectly matched probe has a sequence perfectly complementary to a particular target sequence. The test probe is typically perfectly complementary to a portion of the target sequence. A "polymorphic" marker or site is the locus at which a sequence difference occurs with respect to a reference sequence. Polymorphic markers include restriction fragment
10 length polymorphisms, variable number of tandem repeats (VNTR's), hypervariable regions, minisatellites, dinucleotide repeats, trinucleotide repeats, tetranucleotide repeats, simple sequence repeats, and insertion elements such as Alu. The reference allelic form may be, for example, the most abundant form in a population, or the first allelic form to be identified, and other allelic forms are designated as alternative, variant or polymorphic alleles. The allelic
15 form occurring most frequently in a selected population is sometimes referred to as the "wild type" form, and herein may also be referred to as the "reference" form. Diploid organisms may be homozygous or heterozygous for allelic forms. A diallelic polymorphism has two distinguishable forms (i.e., base sequences), and a triallelic polymorphism has three such forms.

20 As use herein an "oligonucleotide" is a single-stranded nucleic acid ranging in length from 2 to about 60 bases. Oligonucleotides are often synthetic but can also be produced from naturally occurring polynucleotides. A probe is an oligonucleotide capable of binding to a target nucleic acid of complementary sequence through one or more types of chemical bonds, usually through complementary base pairing via hydrogen bond formation. Oligonucleotides
25 probes are often between 5 and 60 bases, and, in specific embodiments, may be between 10-40, or 15-30 bases long. An oligonucleotide probe may include natural (i.e. A, G, C, or T) or modified bases (7-deazaguanosine, inosine, etc.). In addition, the bases in an oligonucleotide probe may be joined by a linkage other than a phosphodiester bond, such as a phosphoramidite linkage or a phosphorothioate linkage, or they may be peptide nucleic acids
30 in which the constituent bases are joined by peptide bonds rather than by phosphodiester bonds, so long as it does not interfere with hybridization. Examples of an oligonucleotide are shown in Table 1. Oligonucleotides can be all of a nucleic acid segment as represented in column 4 of Table 1; a nucleic acid sequence which comprises a nucleic acid segment

represented in column 4 of Table 1 and additional nucleic acids (present at either or both ends of a nucleic acid segment of column 4); or a portion (fragment) of a nucleic acid segment represented in column 4 of the table which includes a polymorphic site. Preferred polymorphic sites of the invention include segments of DNA or their complements, which include any one of the polymorphic sites shown in the Table. The segments can be between 5 and 250 bases, and, in specific embodiments are between 5-10, 5-20, 10-20, 10-50, 20-50 or 10-100 bases. The polymorphic site can occur within any position of the segment. The segments can be from any of the allelic forms of the DNA shown in the Table.

As used herein, the term "primer" refers to a single-stranded oligonucleotide which acts as a point of initiation of template-directed DNA synthesis under appropriate conditions (e.g., in the presence of four different nucleoside triphosphates and a polymerization agent, such as DNA polymerase, RNA polymerase or reverse transcriptase) in an appropriate buffer and at a suitable temperature. The appropriate length of a primer depends on the intended use of the primer, but typically ranges from 15 to 30 nucleotides. Short primer molecules generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. A primer need not be perfectly complementary to the exact sequence of the template, but should be sufficiently complementary to hybridize with it. The term "primer site" refers to the sequence of the target DNA to which a primer hybridizes. The term "primer pair" refers to a set of primers including a 5' (upstream) primer that hybridizes with the 5' end of the DNA sequence to be amplified and a 3' (downstream) primer that hybridizes with the complement of the 3' end of the sequence to be amplified.

DNA fragments can be prepared, for example, by digesting plasmid DNA, or by use of PCR. Oligonucleotides for use as primers or probes are chemically synthesized by methods known in the field of the chemical synthesis of polynucleotides, including by way of non-limiting example the phosphoramidite method described by Beaucage and Carruthers, Tetrahedron Lett 22:1859-1862 (1981) and the triester method provided by Matteucci, et al., J. Am. Chem. Soc., 103:3185 (1981) both incorporated herein by reference. These syntheses may employ an automated synthesizer, as described in Needham-VanDevanter, D.R., et al., Nucleic Acids Res. 12:6159-6168 (1984). Purification of oligonucleotides may be carried out by either native acrylamide gel electrophoresis or by anion-exchange HPLC as described in Pearson, J.D. and Regnier, F.E., J. Chrom., 255:137-149 (1983). A double stranded fragment may then be obtained, if desired, by annealing appropriate complementary single strands together under suitable conditions or by synthesizing the complementary strand

using a DNA polymerase with an appropriate primer sequence. Where a specific sequence for a nucleic acid probe is given, it is understood that the complementary strand is also identified and included. The complementary strand will work equally well in situations where the target is a double-stranded nucleic acid.

5 The sequence of the synthetic oligonucleotide or of any nucleic acid fragment can be obtained using either the dideoxy chain termination method or the Maxam-Gilbert method (see Sambrook et al. Molecular Cloning - a Laboratory Manual (2nd Ed.), Vols. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1989), which is incorporated herein by reference. This manual is hereinafter referred to as "Sambrook et al."

10 ; Zyskind et al., (1988)). Recombinant DNA Laboratory Manual, (Acad. Press, New York). Oligonucleotides useful in diagnostic assays are typically at least 8 consecutive nucleotides in length, and may range upwards of 18 nucleotides in length to greater than 100 or more consecutive nucleotides.

 Another aspect of the invention pertains to isolated antisense nucleic acid molecules

15 that are hybridizable to or complementary to the nucleic acid molecule comprising the SNP-containing nucleotide sequences of the invention, or fragments, analogs or derivatives thereof. An "antisense" nucleic acid comprises a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein, e.g., complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. In specific

20 aspects, antisense nucleic acid molecules are provided that comprise a sequence complementary to at least about 10, about 25, about 50, or about 60 nucleotides or an entire SNP coding strand, or to only a portion thereof.

 In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a polymorphic nucleotide sequence of the invention. The term

25 "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid. In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence of the invention. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated

30 regions).

Given the coding strand sequences disclosed herein, antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick or Hoogsteen base pairing. For example, the antisense nucleic acid molecule can generally be complementary to the entire coding region of an mRNA, but more preferably as embodied herein, it is an
5 oligonucleotide that is antisense to only a portion of the coding or noncoding region of the mRNA. An antisense oligonucleotide can range in length between about 5 and about 60 nucleotides, preferably between about 10 and about 45 nucleotides, more preferably between about 15 and 40 nucleotides, and still more preferably between about 15 and 30 in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis or
10 enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives and acridine
15 substituted nucleotides can be used.

Examples of modified nucleotides that can be used to generate the antisense nucleic acid include: 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine,
20 inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (*v*), wybutoxosine,
25 pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (*v*), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)*w*, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense
30 orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or

genomic DNA encoding a polymorphic protein to thereby inhibit expression of the protein, e.g., by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementary to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule that binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site.

Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, e.g., by linking the antisense nucleic acid molecules to peptides or antibodies that bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids Res* 15: 6625-6641).

The antisense nucleic acid molecule can also comprise a 2'-O-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res* 15: 6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) *FEBS Lett* 215: 327-330).

The following terms are used to describe the sequence relationships between two or more nucleic acids or polynucleotides: "reference sequence", "comparison window", "sequence identity", "percentage of sequence identity", and "substantial identity". A "reference sequence" is a defined sequence used as a basis for a sequence comparison; a reference sequence may be a subset of a larger sequence, for example, as a segment of a full-length cDNA or gene sequence given in a sequence listing, or may comprise a complete cDNA or gene sequence. Optimal alignment of sequences for aligning a comparison window may, for example, be conducted by the local homology algorithm of Smith and Waterman (*Adv. Appl. Math.* 2482 (1981)), by the homology alignment algorithm of Needleman and Wunsch (*J. Mol. Biol.* 48:443 (1970)), by the search for similarity method of Pearson and

Lipman Proc. Natl. Acad. Sci. U.S.A. 852444 (1988), or by computerized implementations of these algorithms (for example, GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Dr., Madison, WI).

5 Techniques for nucleic acid manipulation of the nucleic acid sequences harboring the cSNP's of the invention, such as subcloning nucleic acid sequences encoding polypeptides into expression vectors, labeling probes, DNA hybridization, and the like, are described generally in Sambrook et al., The phrase "nucleic acid sequence encoding" refers to a nucleic acid which directs the expression of a specific protein, peptide or amino acid sequence. The
10 nucleic acid sequences include both the DNA strand sequence that is transcribed into RNA and the RNA sequence that is translated into protein, peptide or amino acid sequence. The nucleic acid sequences include both the full length nucleic acid sequences disclosed herein as well as non-full length sequences derived from the full length protein. It being further understood that the sequence includes the degenerate codons of the native sequence or
15 sequences which may be introduced to provide codon preference in a specific host cell. Consequently, the principles of probe selection and array design can readily be extended to analyze more complex polymorphisms (see EP 730,663). For example, to characterize a triallelic SNP polymorphism, three groups of probes can be designed tiled on the three polymorphic forms as described above. As a further example, to analyze a diallelic
20 polymorphism involving a deletion of a nucleotide, one can tile a first group of probes based on the undeleted polymorphic form as the reference sequence and a second group of probes based on the deleted form as the reference sequence.

For assay of genomic DNA, virtually any biological convenient tissue samples include whole blood, semen, saliva, tears, urine, fecal material, sweat, buccal, skin and hair
25 can be used. Genomic DNA is typically amplified before analysis. Amplification is usually effected by PCR using primers flanking a suitable fragment e.g., of 50-500 nucleotides containing the locus of the polymorphism to be analyzed. Target is usually labeled in the course of amplification. The amplification product can be RNA or DNA, single stranded or double stranded. If double stranded, the amplification product is typically denatured before
30 application to an array. If genomic DNA is analyzed without amplification, it may be desirable to remove RNA from the sample before applying it to the array. Such can be accomplished by digestion with DNase-free RNAase.

DETECTION OF POLYMORPHISMS IN A NUCLEIC ACID SAMPLE

The SNPs disclosed herein can be used to determine which forms of a characterized polymorphism are present in individuals under analysis.

The design and use of allele-specific probes for analyzing polymorphisms is described by e.g., Saiki et al., Nature 324, 163-166 (1986); Dattagupta, EP 235,726, Saiki, WO 89/11548. Allele-specific probes can be designed that hybridize to a segment of target DNA from one individual but do not hybridize to the corresponding segment from another individual due to the presence of different polymorphic forms in the respective segments from the two individuals. Hybridization conditions should be sufficiently stringent that there is a significant difference in hybridization intensity between alleles, and preferably an essentially binary response, whereby a probe hybridizes to only one of the alleles. Some probes are designed to hybridize to a segment of target DNA such that the polymorphic site aligns with a central position (e.g., in a 15-mer at the 7 position; in a 16-mer, at either the 7, 8 or 9 position) of the probe. This design of probe achieves good discrimination in hybridization between different allelic forms.

Allele-specific probes are often used in pairs, one member of a pair showing a perfect match to a reference form of a target sequence and the other member showing a perfect match to a variant form. Several pairs of probes can then be immobilized on the same support for simultaneous analysis of multiple polymorphisms within the same target sequence.

The polymorphisms can also be identified by hybridization to nucleic acid arrays, some examples of which are described in published PCT application WO 95/11995. WO 95/11995 also describes subarrays that are optimized for detection of a variant form of a precharacterized polymorphism. Such a subarray contains probes designed to be complementary to a second reference sequence, which is an allelic variant of the first reference sequence. The second group of probes is designed by the same principles, except that the probes exhibit complementarity to the second reference sequence. The inclusion of a second group (or further groups) can be particularly useful for analyzing short subsequences of the primary reference sequence in which multiple mutations are expected to occur within a short distance commensurate with the length of the probes (e.g., two or more mutations within 9 to 21 bases).

An allele-specific primer hybridizes to a site on target DNA overlapping a polymorphism and only primes amplification of an allelic form to which the primer exhibits perfect complementarity. See Gibbs, *Nucleic Acid Res.* 17 2427-2448 (1989). This primer is used in conjunction with a second primer which hybridizes at a distal site.

- 5 Amplification proceeds from the two-primers, resulting in a detectable product which indicates the particular allelic form is present. A control is usually performed with a second pair of primers, one of which shows a single base mismatch at the polymorphic site and the other of which exhibits perfect complementarity to a distal site. The single-base mismatch prevents amplification and no detectable product is formed. The method works best when
- 10 the mismatch is included in the 3'-most position of the oligonucleotide aligned with the polymorphism because this position is most destabilizing to elongation from the primer (see, e.g., WO 93/22456).

- Amplification products generated using the polymerase chain reaction can be analyzed by the use of denaturing gradient gel electrophoresis. Different alleles can be
- 15 identified based on the different sequence-dependent melting properties and electrophoretic migration of DNA in solution. Erlich, ed., *PCR Technology, Principles and Applications for DNA Amplification*, (W.H. Freeman and Co New York, 1992, Chapter 7).

- Alleles of target sequences can be differentiated using single-strand conformation polymorphism analysis, which identifies base differences by alteration in electrophoretic
- 20 migration of single stranded PCR products, as described in Orita et al., *Proc. Nat. Acad. Sci.* 86, 2766-2770 (1989). Amplified PCR products can be generated and heated or otherwise denatured, to form single stranded amplification products. Single-stranded nucleic acids may refold or form secondary structures which are partially dependent on the base sequence. The different electrophoretic mobilities of single-stranded amplification
- 25 products can be related to base-sequence differences between alleles of target sequences.

- The genotype of an individual with respect to a pathology suspected of being caused by a genetic polymorphism may be assessed by association analysis. Phenotypic traits suitable for association analysis include diseases that have known but hitherto unmapped genetic components (e.g., agammaglobulinemia, diabetes insipidus, Lesch-Nyhan syndrome,
- 30 muscular dystrophy, Wiskott-Aldrich syndrome, Fabry's disease, familial hypercholesterolemia, polycystic kidney disease, hereditary spherocytosis, von Willebrand's

disease, tuberous sclerosis, hereditary hemorrhagic telangiectasia, familial colonic polyposis, Ehlers-Danlos syndrome, osteogenesis imperfecta, and acute intermittent porphyria).

Phenotypic traits also include symptoms of, or susceptibility to, multifactorial diseases of which a component is or may be genetic, such as autoimmune diseases, inflammation, cancer, system, diseases of the nervous and infection by pathogenic microorganisms. Some examples of autoimmune diseases include rheumatoid arthritis, multiple sclerosis, diabetes (insulin-dependent and non-independent), systemic lupus erythematosus and Graves disease. Some examples of cancers include cancers of the bladder, brain, breast, colon, esophagus, kidney, oral cavity, ovary, pancreas, prostate, skin, stomach, leukemia, liver, lung, and uterus. Phenotypic traits also include characteristics such as longevity, appearance (e.g., baldness, obesity), strength, speed, endurance, fertility, and susceptibility or receptivity to particular drugs or therapeutic treatments.

Such correlations can be exploited in several ways. In the case of a strong correlation between a polymorphic form and a disease for which treatment is available, detection of the polymorphic form set in a human or animal patient may justify immediate administration of treatment, or at least the institution of regular monitoring of the patient. Detection of a polymorphic form correlated with serious disease in a couple contemplating a family may also be valuable to the couple in their reproductive decisions. For example, the female partner might elect to undergo in vitro fertilization to avoid the possibility of transmitting such a polymorphism from her husband to her offspring. In the case of a weaker, but still statistically significant correlation between a polymorphic set and human disease, immediate therapeutic intervention or monitoring may not be justified. Nevertheless, the patient can be motivated to begin simple life-style changes (e.g., diet, exercise) that can be accomplished at little cost to the patient but confer potential benefits in reducing the risk of conditions to which the patient may have increased susceptibility by virtue of variant alleles. After determining polymorphic form(s) present in an individual at one or more polymorphic sites, this information can be used in a number of methods.

Determination of which polymorphic forms occupy a set of polymorphic sites in an individual identifies a set of polymorphic forms that distinguishes the individual. See generally National Research Council, *The Evaluation of Forensic DNA Evidence* (Eds. Pollard et al., National Academy Press, DC, 1996). Since the polymorphic sites are within a 50,000 bp region in the human genome, the probability of recombination between these

polymorphic sites is low. That low probability means the haplotype (the set of all 10 polymorphic sites) set forth in this application should be inherited without change for at least several generations. The more sites that are analyzed the lower the probability that the set of polymorphic forms in one individual is the same as that in an unrelated individual.

5 Preferably, if multiple sites are analyzed, the sites are unlinked. Thus, polymorphisms of the invention are often used in conjunction with polymorphisms in distal genes. Preferred polymorphisms for use in forensics are diallelic because the population frequencies of two polymorphic forms can usually be determined with greater accuracy than those of multiple polymorphic forms at multi-allelic loci.

10 The capacity to identify a distinguishing or unique set of forensic markers in an individual is useful for forensic analysis. For example, one can determine whether a blood sample from a suspect matches a blood or other tissue sample from a crime scene by determining whether the set of polymorphic forms occupying selected polymorphic sites is the same in the suspect and the sample. If the set of polymorphic markers does not match
15 between a suspect and a sample, it can be concluded (barring experimental error) that the suspect was not the source of the sample. If the set of markers does match, one can conclude that the DNA from the suspect is consistent with that found at the crime scene. If frequencies of the polymorphic forms at the loci tested have been determined (e.g., by analysis of a suitable population of individuals), one can perform a statistical analysis to determine the
20 probability that a match of suspect and crime scene sample would occur by chance.

$p(ID)$ is the probability that two random individuals have the same polymorphic or allelic form at a given polymorphic site. In diallelic loci, four genotypes are possible: AA, AB, BA, and BB. If alleles A and B occur in a haploid genome of the organism with frequencies x and y , the probability of each genotype in a diploid organism are (see WO
25 95/12607):

$$\text{Homozygote: } p(AA)=x^2$$

$$\text{Homozygote: } p(BB)=y^2=(1-x)^2$$

$$\text{Single Heterozygote: } p(AB)=p(BA)=xy=x(1-x)$$

$$\text{Both Heterozygotes: } p(AB+BA)=2xy=2x(1-x)$$

The probability of identity at one locus (i.e. the probability that two individuals, picked at random from a population will have identical polymorphic forms at a given locus) is given by the equation:

$$p(ID) = (x^2)^2 + (2xy)^2 + (y^2)^2.$$

- 5 These calculations can be extended for any number of polymorphic forms at a given locus. For example, the probability of identity $p(ID)$ for a 3-allele system where the alleles have the frequencies in the population of x , y and z , respectively, is equal to the sum of the squares of the genotype frequencies:

$$p(ID) = x^4 + (2xy)^2 + (2yz)^2 + (2xz)^2 + z^4 + y^4$$

- 10 In a locus of n alleles, the appropriate binomial expansion is used to calculate $p(ID)$ and $p(exc)$.

The cumulative probability of identity ($cum p(ID)$) for each of multiple unlinked loci is determined by multiplying the probabilities provided by each locus:

$$cum p(ID) = p(ID1)p(ID2)p(ID3) \dots p(IDn)$$

- 15 The cumulative probability of non-identity for n loci (i.e. the probability that two random individuals will be different at 1 or more loci) is given by the equation:

$$cum p(nonID) = 1 - cum p(ID).$$

- If several polymorphic loci are tested, the cumulative probability of non-identity for random individuals becomes very high (e.g., one billion to one). Such probabilities can be taken into
20 account together with other evidence in determining the guilt or innocence of the suspect.

- The object of paternity testing is usually to determine whether a male is the father of a child. In most cases, the mother of the child is known and thus, the mother's contribution to the child's genotype can be traced. Paternity testing investigates whether the part of the child's genotype not attributable to the mother is consistent with that of the putative father. Paternity
25 testing can be performed by analyzing sets of polymorphisms in the putative father and the child.

If the set of polymorphisms in the child attributable to the father does not match the putative father, it can be concluded, barring experimental error, that the putative father is not the real father. If the set of polymorphisms in the child attributable to the father does match the set of polymorphisms of the putative father, a statistical calculation can be performed to determine the probability of coincidental match.

The probability of parentage exclusion (representing the probability that a random male will have a polymorphic form at a given polymorphic site that makes him incompatible as the father) is given by the equation (see WO 95/12607):

$$p(exc) = xy(1-xy)$$

where x and y are the population frequencies of alleles A and B of a diallelic polymorphic site. (At a triallelic site $p(exc) = xy(1-xy) + yz(1-yz) + xz(1-xz) + 3xyz(1-xyz)$), where x, y and z are the respective population frequencies of alleles A, B and C). The probability of non-exclusion is:

$$p(non-exc) = 1 - p(exc)$$

The cumulative probability of non-exclusion (representing the value obtained when n loci are used) is thus:

$$cum p(non-exc) = p(non-exc1)p(non-exc2)p(non-exc3) \dots p(non-excn)$$

The cumulative probability of exclusion for n loci (representing the probability that a random male will be excluded) is:

$$cum p(exc) = 1 - cum p(non-exc).$$

If several polymorphic loci are included in the analysis, the cumulative probability of exclusion of a random male is very high. This probability can be taken into account in assessing the liability of a putative father whose polymorphic marker set matches the child's polymorphic marker set attributable to his/her father.

The polymorphisms of the invention may contribute to the phenotype of an organism in different ways. Some polymorphisms occur within a protein coding sequence and contribute to phenotype by affecting protein structure. The effect may be neutral, beneficial or detrimental, or both beneficial and detrimental, depending on the circumstances. For

example, a heterozygous sickle cell mutation confers resistance to malaria, but a homozygous sickle cell mutation is usually lethal. Other polymorphisms occur in noncoding regions but may exert phenotypic effects indirectly via influence on replication, transcription, and translation. A single polymorphism may affect more than one phenotypic trait. Likewise, a
5 single phenotypic trait may be affected by polymorphisms in different genes. Further, some polymorphisms predispose an individual to a distinct mutation that is causally related to a certain phenotype.

Phenotypic traits include diseases that have known but hitherto unmapped genetic components. Phenotypic traits also include symptoms of, or susceptibility to, multifactorial
10 diseases of which a component is or may be genetic, such as autoimmune diseases, inflammation, cancer, diseases of the nervous system, and infection by pathogenic microorganisms. Some examples of autoimmune diseases include rheumatoid arthritis, multiple sclerosis, diabetes (insulin-dependent and non-independent), systemic lupus erythematosus and Graves disease. Some examples of cancers include cancers of the bladder,
15 brain, breast, colon, esophagus, kidney, leukemia, liver, lung, oral cavity, ovary, pancreas, prostate, skin, stomach and uterus. Phenotypic traits also include characteristics such as longevity, appearance (e.g., baldness, obesity), strength, speed, endurance, fertility, and susceptibility or receptivity to particular drugs or therapeutic treatments.

Correlation is performed for a population of individuals who have been tested for the
20 presence or absence of a phenotypic trait of interest and for polymorphic markers sets. To perform such analysis, the presence or absence of a set of polymorphisms (i.e. a polymorphic set) is determined for a set of the individuals, some of whom exhibit a particular trait, and some of which exhibit lack of the trait. The alleles of each polymorphism of the set are then reviewed to determine whether the presence or absence of a particular allele is associated
25 with the trait of interest. Correlation can be performed by standard statistical methods such as a χ^2 -squared test and statistically significant correlations between polymorphic form(s) and phenotypic characteristics are noted. For example, it might be found that the presence of allele A1 at polymorphism A correlates with heart disease. As a further example, it might be found that the combined presence of allele A1 at polymorphism A and allele B1 at
30 polymorphism B correlates with increased milk production of a farm animal.

Such correlations can be exploited in several ways. In the case of a strong correlation between a set of one or more polymorphic forms and a disease for which treatment is

available, detection of the polymorphic form set in a human or animal patient may justify immediate administration of treatment, or at least the institution of regular monitoring of the patient. Detection of a polymorphic form correlated with serious disease in a couple contemplating a family may also be valuable to the couple in their reproductive decisions.

5 For example, the female partner might elect to undergo in vitro fertilization to avoid the possibility of transmitting such a polymorphism from her husband to her offspring. In the case of a weaker, but still statistically significant correlation between a polymorphic set and human disease, immediate therapeutic intervention or monitoring may not be justified. Nevertheless, the patient can be motivated to begin simple life-style changes (e.g., diet,
10 exercise) that can be accomplished at little cost to the patient but confer potential benefits in reducing the risk of conditions to which the patient may have increased susceptibility by virtue of variant alleles. Identification of a polymorphic set in a patient correlated with enhanced receptiveness to one of several treatment regimes for a disease indicates that this treatment regime should be followed.

15 For animals and plants, correlations between characteristics and phenotype are useful for breeding for desired characteristics. For example, Beitz et al., U.S. Pat. No. 5,292,639 discuss use of bovine mitochondrial polymorphisms in a breeding program to improve milk production in cows. To evaluate the effect of mtDNA D-loop sequence polymorphism on milk production, each cow was assigned a value of 1 if variant or 0 if wild type with respect
20 to a prototypical mitochondrial DNA sequence at each of 17 locations considered.

The previous section concerns identifying correlations between phenotypic traits and polymorphisms that directly or indirectly contribute to those traits. The present section describes identification of a physical linkage between a genetic locus associated with a trait of interest and polymorphic markers that are not associated with the trait, but are in physical
25 proximity with the genetic locus responsible for the trait and co-segregate with it. Such analysis is useful for mapping a genetic locus associated with a phenotypic trait to a chromosomal position, and thereby cloning gene(s) responsible for the trait. See Lander et al., *Proc. Natl. Acad. Sci. (USA)* 83, 7353-7357 (1986); Lander et al., *Proc. Natl. Acad. Sci. (USA)* 84, 2363-2367 (1987); Donis-Keller et al., *Cell* 51, 319-337 (1987); Lander et al.,
30 *Genetics* 121, 185-199 (1989)). Genes localized by linkage can be cloned by a process known as directional cloning. See Wainwright, *Med. J. Australia* 159, 170-174 (1993); Collins, *Nature Genetics* 1, 3-6 (1992) (each of which is incorporated by reference in its entirety for all purposes).

Linkage studies are typically performed on members of a family. Available members of the family are characterized for the presence or absence of a phenotypic trait and for a set of polymorphic markers. The distribution of polymorphic markers in an informative meiosis is then analyzed to determine which polymorphic markers co-segregate with a phenotypic trait. See, e.g., Kerem et al., *Science* 245, 1073-1080 (1989); Monaco et al., *Nature* 316, 842 (1985); Yamoka et al., *Neurology* 40, 222-226 (1990); Rossiter et al., *FASEB Journal* 5, 21-27 (1991).

Linkage is analyzed by calculation of LOD (log of the odds) values. A lod value is the relative likelihood of obtaining observed segregation data for a marker and a genetic locus when the two are located at a recombination fraction θ , versus the situation in which the two are not linked, and thus segregating independently (Thompson & Thompson, *Genetics in Medicine* (5th ed, W.B. Saunders Company, Philadelphia, 1991); Strachan, "Mapping the human genome" in *The Human Genome* (BIOS Scientific Publishers Ltd, Oxford), Chapter 4). A series of likelihood ratios are calculated at various recombination fractions (θ), ranging from $\theta = 0.0$ (coincident loci) to $\theta = 0.50$ (unlinked). Thus, the likelihood at a given value of θ is: probability of data if loci linked at θ to probability of data if loci unlinked. The computed likelihood is usually expressed as the \log_{10} of this ratio (i.e., a lod score). For example, a lod score of 3 indicates 1000:1 odds against an apparent observed linkage being a coincidence. The use of logarithms allows data collected from different families to be combined by simple addition. Computer programs are available for the calculation of lod scores for differing values of θ (e.g., LIPED, MLINK (Lathrop, *Proc. Nat. Acad. Sci. (USA)* 81, 3443-3446 (1984)). For any particular lod score, a recombination fraction may be determined from mathematical tables. See Smith et al., *Mathematical tables for research workers in human genetics* (Churchill, London, 1961); Smith, *Ann. Hum. Genet.* 32, 127-150 (1968). The value of θ at which the lod score is the highest is considered to be the best estimate of the recombination fraction.

Positive lod score values suggest that the two loci are linked, whereas negative values suggest that linkage is less likely (at that value of θ) than the possibility that the two loci are unlinked. By convention, a combined lod score of + 3 or greater (equivalent to greater than 1000:1 odds in favor of linkage) is considered definitive evidence that two loci are linked. Similarly, by convention, a negative lod score of -2 or less is taken as definitive evidence against linkage of the two loci being compared. Negative linkage data are useful in

excluding a chromosome or a segment thereof from consideration. The search focuses on the remaining non-excluded chromosomal locations.

The invention further provides transgenic nonhuman animals capable of expressing an exogenous variant gene and/or having one or both alleles of an endogenous variant gene inactivated. Expression of an exogenous variant gene is usually achieved by operably
5 linking the gene to a promoter and optionally an enhancer, and microinjecting the construct into a zygote. See Hogan et al., "Manipulating the Mouse Embryo, A Laboratory Manual," Cold Spring Harbor Laboratory. (1989). Inactivation of endogenous variant genes can be achieved by forming a transgene in which a cloned variant gene is inactivated by insertion of
10 a positive selection marker. See Capecchi, Science 244, 1288-1292 The transgene is then introduced into an embryonic stem cell, where it undergoes homologous recombination with an endogenous variant gene. Mice and other rodents are preferred animals. Such animals provide useful drug screening systems.

The invention further provides methods for assessing the pharmacogenomic
15 susceptibility of a subject harboring a single nucleotide polymorphism to a particular pharmaceutical compound, or to a class of such compounds. Genetic polymorphism in drug-metabolizing enzymes, drug transporters, receptors for pharmaceutical agents, and other drug targets have been correlated with individual differences based on distinction in the efficacy and toxicity of the pharmaceutical agent administered to a subject. Pharmacogenomic
20 characterization of a subjects susceptibility to a drug enhances the ability to tailor a dosing regimen to the particular genetic constitution of the subject, thereby enhancing and optimizing the therapeutic effectiveness of the therapy.

In cases in which a cSNP leads to a polymorphic protein that is ascribed to be the cause of a pathological condition, method of treating such a condition includes administering
25 to a subject experiencing the pathology the wild type cognate of the polymorphic protein. Once administered in an effective dosing regimen, the wild type cognate provides complementation or remediation of the defect due to the polymorphic protein. The subject's condition is ameliorated by this protein therapy.

A subject suspected of suffering from a pathology ascribable to a polymorphic protein
30 that arises from a cSNP is to be diagnosed using any of a variety of diagnostic methods capable of identifying the presence of the cSNP in the nucleic acid, or of the cognate

polymorphic protein, in a suitable clinical sample taken from the subject. Once the presence of the cSNP has been ascertained, and the pathology is correctable by administering a normal or wild-type gene, the subject is treated with a pharmaceutical composition that includes a nucleic acid that harbors the correcting wild-type gene, or a fragment containing a correcting sequence of the wild-type gene. Non-limiting examples of ways in which such a nucleic acid may be administered include incorporating the wild-type gene in a viral vector, such as an adenovirus or adeno associated virus, and administration of a naked DNA in a pharmaceutical composition that promotes intracellular uptake of the administered nucleic acid. Once the nucleic acid that includes the gene coding for the wild-type allele of the polymorphism is incorporated within a cell of the subject, it will initiate *de novo* biosynthesis of the wild-type gene product. If the nucleic acid is further incorporated into the genome of the subject, the treatment will have long-term effects, providing *de novo* synthesis of the wild-type protein for a prolonged duration. The synthesis of the wild-type protein in the cells of the subject will contribute to a therapeutic enhancement of the clinical condition of the subject.

A subject suffering from a pathology ascribed to a SNP may be treated so as to correct the genetic defect. (See Kren et al., Proc. Natl. Acad. Sci. USA 96:10349-10354 (1999)). Such a subject is identified by any method that can detect the polymorphism in a sample drawn from the subject. Such a genetic defect may be permanently corrected by administering to such a subject a nucleic acid fragment incorporating a repair sequence that supplies the wild-type nucleotide at the position of the SNP. This site-specific repair sequence encompasses an RNA/DNA oligonucleotide which operates to promote endogenous repair of a subject's genomic DNA. Upon administration in an appropriate vehicle, such as a complex with polyethylenimine or encapsulated in anionic liposomes, a genetic defect leading to an inborn pathology may be overcome, as the chimeric oligonucleotides induces incorporation of the wild-type sequence into the subject's genome. Upon incorporation, the wild-type gene product is expressed, and the replacement is propagated, thereby engendering a permanent repair.

The invention further provides kits comprising at least one allele-specific oligonucleotide as described above. Often, the kits contain one or more pairs of allele-specific oligonucleotides hybridizing to different forms of a polymorphism. In some kits, the allele-specific oligonucleotides are provided immobilized to a substrate. For example, the same substrate can comprise allele-specific oligonucleotide probes for detecting at least 10, 100, 1000 or all of the polymorphisms shown in the Table. Optional additional

components of the kit include, for example, restriction enzymes, reverse-transcriptase or polymerase, the substrate nucleoside triphosphates, means used to label (for example, an avidin-enzyme conjugate and enzyme substrate and chromogen if the label is biotin), and the appropriate buffers for reverse transcription, PCR, or hybridization reactions. Usually, the
5 kit also contains instructions for carrying out the hybridizing methods.

Several aspects of the present invention rely on having available the polymorphic proteins encoded by the nucleic acids comprising a SNP of the inventions. There are various methods of isolating these nucleic acid sequences. For example, DNA is isolated from a genomic or cDNA library using labeled oligonucleotide probes having sequences
10 complementary to the sequences disclosed herein.

Such probes can be used directly in hybridization assays. Alternatively probes can be designed for use in amplification techniques such as PCR.

To prepare a cDNA library, mRNA is isolated from tissue such as heart or pancreas, preferably a tissue wherein expression of the gene or gene family is likely to occur. cDNA is
15 prepared from the mRNA and ligated into a recombinant vector. The vector is transfected into a recombinant host for propagation, screening and cloning. Methods for making and screening cDNA libraries are well known, See Gubler, U. and Hoffman, B.J. *Gene* 25:263-269 (1983) and Sambrook et al.

For a genomic library, for example, the DNA is extracted from tissue and either
20 mechanically sheared or enzymatically digested to yield fragments of about 12-20 kb. The fragments are then separated by gradient centrifugation from undesired sizes and are constructed in bacteriophage lambda vectors. These vectors and phage are packaged *in vitro*, as described in Sambrook, et al. Recombinant phage are analyzed by plaque hybridization as described in Benton and Davis, Science 196:180-182 (1977). Colony hybridization is carried
25 out as generally described in M. Grunstein et al. *Proc. Natl. Acad. Sci. USA.* 72:3961-3965 (1975). DNA of interest is identified in either cDNA or genomic libraries by its ability to hybridize with nucleic acid probes, for example on Southern blots, and these DNA regions are isolated by standard methods familiar to those of skill in the art. See Sambrook, et al.

In PCR techniques, oligonucleotide primers complementary to the two 3' borders of
30 the DNA region to be amplified are synthesized. The polymerase chain reaction is then carried out using the two primers. See PCR Protocols: a Guide to Methods and Applications

(Innis, M, Gelfand, D., Sninsky, J. and White, T., eds.), Academic Press, San Diego (1990).

Primers can be selected to amplify the entire regions encoding a full-length sequence of interest or to amplify smaller DNA segments as desired. PCR can be used in a variety of protocols to isolate cDNA's encoding a sequence of interest. In these protocols, appropriate
5 primers and probes for amplifying DNA encoding a sequence of interest are generated from analysis of the DNA sequences listed herein. Once such regions are PCR-amplified, they can be sequenced and oligonucleotide probes can be prepared from the sequence.

Once DNA encoding a sequence comprising a cSNP is isolated and cloned, one can express the encoded polymorphic proteins in a variety of recombinantly engineered cells. It
10 is expected that those of skill in the art are knowledgeable in the numerous expression systems available for expression of DNA encoding a sequence of interest. No attempt to describe in detail the various methods known for the expression of proteins in prokaryotes or eukaryotes is made here.

In brief summary, the expression of natural or synthetic nucleic acids encoding a
15 sequence of interest will typically be achieved by operably linking the DNA or cDNA to a promoter (which is either constitutive or inducible), followed by incorporation into an expression vector. The vectors can be suitable for replication and integration in either prokaryotes or eukaryotes. Typical expression vectors contain, initiation sequences, transcription and translation terminators, and promoters useful for regulation of the
20 expression of a polynucleotide sequence of interest. To obtain high level expression of a cloned gene, it is desirable to construct expression plasmids which contain, at the minimum, a strong promoter to direct transcription, a ribosome binding site for translational initiation, and a transcription/translation terminator. The expression vectors may also comprise generic expression cassettes containing at least one independent terminator sequence, sequences
25 permitting replication of the plasmid in both eukaryotes and prokaryotes, i.e., shuttle vectors, and selection markers for both prokaryotic and eukaryotic systems. See Sambrook et al.

A variety of prokaryotic expression systems may be used to express the polymorphic proteins of the invention. Examples include *E. coli*, *Bacillus*, *Streptomyces*, and the like.

It is preferred to construct expression plasmids which contain, at the minimum, a
30 strong promoter to direct transcription, a ribosome binding site for translational initiation, and a transcription/translation terminator. Examples of regulatory regions suitable for this

purpose in *E. coli* are the promoter and operator region of the *E. coli* tryptophan biosynthetic pathway as described by Yanofsky, C., J. Bacterial. 158:1018-1024 (1984) and the leftward promoter of phage lambda (P_L) as described by A, I. and Hagen, D., Ann. Rev. Genet. 14:399-445 (1980). The inclusion of selection markers in DNA vectors transformed in *E.*
5 *coli* is also useful. Examples of such markers include genes specifying resistance to ampicillin, tetracycline, or chloramphenicol. See Sambrook et al. for details concerning selection markers for use in *E. coli*.

To enhance proper folding of the expressed recombinant protein, during purification from *E. coli*, the expressed protein may first be denatured and then renatured. This can be
10 accomplished by solubilizing the bacterially produced proteins in a chaotropic agent such as guanidine HCl and reducing all the cysteine residues with a reducing agent such as beta-mercaptoethanol. The protein is then renatured, either by slow dialysis or by gel filtration. See U.S. Patent No. 4,511,503. Detection of the expressed antigen is achieved by methods known in the art as radioimmunoassay, or Western blotting techniques or
15 immunoprecipitation. Purification from *E. coli* can be achieved following procedures such as those described in U.S. Patent No. 4,511,503.

Any of a variety of eukaryotic expression systems such as yeast, insect cell lines, bird, fish, and mammalian cells, may also be used to express a polymorphic protein of the invention. As explained briefly below, a nucleotide sequence harboring a cSNP may be
20 expressed in these eukaryotic systems. Synthesis of heterologous proteins in yeast is well known. Methods in Yeast Genetics, Sherman, F., et al., Cold Spring Harbor Laboratory, (1982) is a well recognized work describing the various methods available to produce the protein in yeast. Suitable vectors usually have expression control sequences, such as promoters, including 3-phosphoglycerate kinase or other glycolytic enzymes, and an origin of
25 replication, termination sequences and the like as desired. For instance, suitable vectors are described in the literature (Botstein, et al., Gene 8:17-24 (1979); Broach, et al., Gene 8:121-133 (1979)).

Two procedures are used in transforming yeast cells. In one case, yeast cells are first converted into protoplasts using zymolyase, lyticase or glucanase, followed by addition of
30 DNA and polyethylene glycol (PEG). The PEG-treated protoplasts are then regenerated in a 3% agar medium under selective conditions. Details of this procedure are given in the papers by J.D. Beggs, Nature (London) 275:104-109 (1978); and Hinnen, A., et al., Proc. Natl.

Acad. Sci. USA, 75:1929-1933 (1978). The second procedure does not involve removal of the cell wall. Instead the cells are treated with lithium chloride or acetate and PEG and put on selective plates (Ito, H., et al., J. Bact, 153:163-168 (1983)). cells and applying standard protein isolation techniques to the lysates:.

5 The purification process can be monitored by using Western blot techniques or radioimmunoassay or other standard techniques. The sequences encoding the proteins of the invention can also be ligated to various immunoassay expression vectors for use in transforming cell cultures of, for instance, mammalian, insect, bird or fish origin. Illustrative of cell cultures useful for the production of the polypeptides are mammalian cells.

10 Mammalian cell systems often will be in the form of monolayers of cells although mammalian cell suspensions may also be used. A number of suitable host cell lines capable of expressing intact proteins have been developed in the art, and include the HEK293, BHK21, and CHO cell lines, and various human cells such as COS cell lines, HeLa cells, myeloma cell lines, Jurkat cells, etc. Expression vectors for these cells can include
15 expression control sequences, such as an origin of replication, a promoter (e.g., the CMV promoter, a HSV *tk* promoter or *pgk* (phosphoglycerate kinase) promoter), an enhancer (Queen et al. Immunol. Rev. 89:49 (1986)) and necessary processing information sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites (e.g., an SV40 large T Ag poly A addition site), and transcriptional terminator sequences.

20 Other animal cells are available, for instance, from the American Type Culture Collection Catalogue of Cell Lines and Hybridomas (7th edition, (1992)). Appropriate vectors for expressing the proteins of the invention in insect cells are usually derived from baculovirus. Insect cell lines include mosquito larvae, silkworm, armyworm, moth and *Drosophila* cell lines such as a Schneider cell line (See Schneider J. Embryol. Exp. Morphol.,
25 27:353-365 (1987). As indicated above, the vector, e.g., a plasmid, which is used to transform the host cell, preferably contains DNA sequences to initiate transcription and sequences to control the translation of the protein. These sequences are referred to as expression control sequences. As with yeast, when higher animal host cells are employed, polyadenylation or transcription terminator sequences from known mammalian genes need to
30 be incorporated into the vector. An example of a terminator sequence is the polyadenylation sequence from the bovine growth hormone gene. Sequences for accurate splicing of the transcript may also be included. An example of a splicing sequence is the VP1 intron from SV40 (Sprague, J. et al., J. Virol. 45: 773-781 (1983)). Additionally, gene sequences to

control replication in the host cell may be Saveria-Campo, M., 1985, "Bovine Papilloma virus DNA a Eukaryotic Cloning Vector" in DNA Cloning Vol. II a Practical Approach Ed. D.M. Glover, IRL Press, Arlington, Virginia pp. 213-238. The host cells are competent or rendered competent for transformation by various means. There are several well-known
5 methods of introducing DNA into animal cells. These include: calcium phosphate precipitation, fusion of the recipient cells with bacterial protoplasts containing the DNA, treatment of the recipient cells with liposomes containing the DNA, DEAE dextran, electroporation and micro-injection of the DNA directly into the cells.

The transformed cells are cultured by means well known in the art (Biochemical
10 Methods in Cell Culture and Virology, Kuchler, R.J., Dowden, Hutchinson and Ross, Inc., (1977)). The expressed polypeptides are isolated from cells grown as suspensions or as monolayers. The latter are recovered by well known mechanical, chemical or enzymatic means.

General methods of expressing recombinant proteins are also known and are
15 exemplified in R. Kaufman, Methods in Enzymology 185, 537-566 (1990). As defined herein "operably linked" refers to linkage of a promoter upstream from a DNA sequence such that the promoter mediates transcription of the DNA sequence. Specifically, "operably linked" means that the isolated polynucleotide of the invention and an expression control sequence are situated within a vector or cell in such a way that the gene encoding the protein
20 is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression sequence. The term "vector", refers to viral expression systems, autonomous self-replicating circular DNA (plasmids), and includes both expression and nonexpression plasmids.

The term "gene" as used herein is intended to refer to a nucleic acid sequence which
25 encodes a polypeptide. This definition includes various sequence polymorphisms, mutations, and/or sequence variants wherein such alterations do not affect the function of the gene product. The term "gene" is intended to include not only coding sequences but also regulatory regions such as promoters, enhancers, termination regions and similar untranslated nucleotide sequences. The term further includes all introns and other DNA sequences spliced
30 from the mRNA transcript, along with variants resulting from alternative splice sites.

A number of types of cells may act as suitable host cells for expression of the protein. Mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A43 1 cells, human Co10205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains
5 derived from in vitro culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL- 60, U937, HaK or Jurkat cells. Alternatively, it may be possible to produce the protein in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Potentially suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida* or any yeast strain capable of expressing heterologous
10 proteins. Potentially suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins. If the protein is made in yeast or bacteria, it may be necessary to modify the protein produced therein, for example by phosphorylation or glycosylation of the appropriate sites, in order to obtain the functional protein.

15 The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, e.g., Invitrogen, San Diego, California, U.S.A. (the MaxBac© kit), and such methods are well known in the art, as
20 described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed." The protein of the invention may be prepared by culturing transformed host cells under culture conditions suitable to express the recombinant protein.

25 The polymorphic protein of the invention may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a nucleotide sequence encoding the protein. The protein may also be produced by known conventional chemical synthesis. Methods for constructing the proteins of the present invention by synthetic
30 means are known to those skilled in the art.

The polymorphic proteins produced by recombinant DNA technology may be purified by techniques commonly employed to isolate or purify recombinant proteins. Recombinantly

produced proteins can be directly expressed or expressed as a fusion protein. The protein is then purified by a combination of cell lysis (e.g., sonication) and affinity chromatography. For fusion products, subsequent digestion of the fusion protein with an appropriate proteolytic enzyme releases the desired polypeptide. The polypeptides of this invention may
5 be purified to substantial purity by standard techniques well known in the art, including selective precipitation with such substances as ammonium sulfate, column chromatography, immunopurification methods, and others. See, for instance, R. Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag: New York (1982), incorporated herein by reference. For example, in an embodiment, antibodies may be raised to the proteins of the invention as
10 described herein. Cell membranes are isolated from a cell line expressing the recombinant protein, the protein is extracted from the membranes and immunoprecipitated. The proteins may then be further purified by standard protein chemistry techniques as described above.

The resulting expressed protein may then be purified from such culture (i.e., from culture medium or cell extracts) using known purification processes, such as gel filtration
15 and ion exchange chromatography. The purification of the protein may also include an affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-Toyopearl® or Cibacrom blue 3GA Sepharose B; one or more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity
20 chromatography. Alternatively, the protein of the invention may also be expressed in a form which will facilitate purification. For example, it may be expressed as a fusion protein, such as those of maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin (TRX). Kits for expression and purification of such fusion proteins are commercially available from New England BioLab (Beverly, MA), Pharmacia (Piscataway, NJ) and
25 InVitrogen, respectively. The protein can also be tagged with an epitope and subsequently purified by using a specific antibody directed to such epitope. One such epitope ("Flag") is commercially available from Kodak (New Haven, CT). Finally, one or more reverse-phase high performance liquid chromatography (RP- HPLC) steps employing hydrophobic RP- HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be
30 employed to further purify the protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous isolated recombinant protein. The protein thus purified is substantially free of other

mammalian proteins and is defined in accordance with the present invention as an "isolated protein."

The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, *i.e.*, molecules that contain
5 an antigen binding site that specifically binds (immunoreacts with) an antigen, such as polymorphic. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, F_{ab} and $F_{(ab)2}$ fragments, and an F_{ab} expression library. In a specific embodiment, antibodies to human polymorphic proteins are disclosed.

The phrase "specifically binds to", "immunospecifically binds to" or is "specifically
10 immunoreactive with", an antibody when referring to a protein or peptide, refers to a binding reaction which is determinative of the presence of the protein in the presence of a heterogeneous population of proteins and other biological materials. Thus, for example, under designated immunoassay conditions, the specified antibodies bind to a particular protein and do not bind in a significant amount to other proteins present in the sample.
15 Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. Of particular interest in the present invention is an antibody that binds immunospecifically to a polymorphic protein but not to its cognate wild type allelic protein, or vice versa. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example,
20 solid-phase ELISA immunoassays are routinely used to select monoclonal antibodies specifically immunoreactive with a protein. See Harlow and Lane (1988) Antibodies, a Laboratory Manual, Cold Spring Harbor Publications, New York, for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity.

25 Polyclonal and/or monoclonal antibodies that immunospecifically bind to polymorphic gene products but not to the corresponding prototypical or "wild-type" gene products are also provided. Antibodies can be made by injecting mice or other animals with the variant gene product or synthetic peptide. Monoclonal antibodies are screened as are described, for example, in Harlow & Lane, Antibodies, A Laboratory Manual, Cold Spring
30 Harbor Press, New York (1988); Goding, Monoclonal antibodies, Principles and Practice (2d ed.) Academic Press, New York (1986). Monoclonal antibodies are tested for specific

immunoreactivity with a variant gene product and lack of immunoreactivity to the corresponding prototypical gene product.

An isolated polymorphic protein, or a portion or fragment thereof, can be used as an immunogen to generate the antibody that bind the polymorphic protein using standard techniques for polyclonal and monoclonal antibody preparation. The full-length polymorphic protein can be used or, alternatively, the invention provides antigenic peptide fragments of polymorphic for use as immunogens. The antigenic peptide of a polymorphic protein of the invention comprises at least 8 amino acid residues of the amino acid sequence encompassing the polymorphic amino acid and encompasses an epitope of the polymorphic protein such that an antibody raised against the peptide forms a specific immune complex with the polymorphic protein. Preferably, the antigenic peptide comprises at least 10 amino acid residues, more preferably at least 15 amino acid residues, even more preferably at least 20 amino acid residues, and most preferably at least 30 amino acid residues. Preferred epitopes encompassed by the antigenic peptide are regions of polymorphic that are located on the surface of the protein, *e.g.*, hydrophilic regions.

For the production of polyclonal antibodies, various suitable host animals (*e.g.*, rabbit, goat, mouse or other mammal) may be immunized by injection with the polymorphic protein. An appropriate immunogenic preparation can contain, for example, recombinantly expressed polymorphic protein or a chemically synthesized polymorphic polypeptide. The preparation can further include an adjuvant. Various adjuvants used to increase the immunological response include, but are not limited to, Freund's (complete and incomplete), mineral gels (*e.g.*, aluminum hydroxide), surface active substances (*e.g.*, lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, etc.), human adjuvants such as *Bacille Calmette-Guerin* and *Corynebacterium parvum*, or similar immunostimulatory agents. If desired, the antibody molecules directed against polymorphic proteins can be isolated from the mammal (*e.g.*, from the blood) and further purified by well known techniques, such as protein A chromatography, to obtain the IgG fraction.

The term "monoclonal antibody" or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that originates from the clone of a singly hybridoma cell, and that contains only one type of antigen binding site capable of immunoreacting with a particular epitope of a polymorphic protein. A monoclonal antibody composition thus typically displays a single binding affinity for a particular polymorphic

protein with which it immunoreacts. For preparation of monoclonal antibodies directed towards a particular polymorphic protein, or derivatives, fragments, analogs or homologs thereof, any technique that provides for the production of antibody molecules by continuous cell line culture may be utilized. Such techniques include, but are not limited to, the

5 hybridoma technique (see Kohler & Milstein, 1975 *Nature* 256: 495-497); the trioma technique; the human B-cell hybridoma technique (see Kozbor, *et al.*, 1983 *Immunol Today* 4: 72) and the EBV hybridoma technique to produce human monoclonal antibodies (see Cole, *et al.*, 1985 In: MONOCLONAL ANTIBODIES AND CANCER THERAPY, Alan R. Liss, Inc., pp. 77-96). Human monoclonal antibodies may be utilized in the practice of the present

10 invention and may be produced by using human hybridomas (see Cote, *et al.*, 1983. *Proc Natl Acad Sci USA* 80: 2026-2030) or by transforming human B-cells with Epstein Barr Virus *in vitro* (see Cole, *et al.*, 1985 In: MONOCLONAL ANTIBODIES AND CANCER THERAPY, Alan R. Liss, Inc., pp. 77-96).

According to the invention, techniques can be adapted for the production of

15 single-chain antibodies specific to a polymorphic protein (see *e.g.*, U.S. Patent No. 4,946,778). In addition, methodologies can be adapted for the construction of F_{ab} expression libraries (see *e.g.*, Huse, *et al.*, 1989 *Science* 246: 1275-1281) to allow rapid and effective identification of monoclonal F_{ab} fragments with the desired specificity for a polymorphic protein or derivatives, fragments, analogs or homologs thereof. Non-human antibodies can

20 be "humanized" by techniques well known in the art. See *e.g.*, U.S. Patent No. 5,225,539. Antibody fragments that contain the idiotypes to a polymorphic protein may be produced by techniques known in the art including, but not limited to: (i) an $F_{(ab)2}$ fragment produced by pepsin digestion of an antibody molecule; (ii) an F_{ab} fragment generated by reducing the disulfide bridges of an $F_{(ab)2}$ fragment; (iii) an F_{ab} fragment generated by the treatment of the

25 antibody molecule with papain and a reducing agent and (iv) F_v fragments.

Additionally, recombinant anti-polymorphic protein antibodies, such as chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, which can be made using standard recombinant DNA techniques, are within the scope of the invention. Such chimeric and humanized monoclonal antibodies can be produced by

30 recombinant DNA techniques known in the art, for example using methods described in PCT International Application No. PCT/US86/02269; European Patent Application No. 184,187; European Patent Application No. 171,496; European Patent Application No. 173,494; PCT International Publication No. WO 86/01533; U.S. Pat. No. 4,816,567; European Patent

Application No. 125,023; Better *et al.* (1988) *Science* 240:1041-1043; Liu *et al.* (1987) *PNAS* 84:3439-3443; Liu *et al.* (1987) *J Immunol.* 139:3521-3526; Sun *et al.* (1987) *PNAS* 84:214-218; Nishimura *et al.* (1987) *Cancer Res* 47:999-1005; Wood *et al.* (1985) *Nature* 314:446-449; Shaw *et al.* (1988) *J Natl Cancer Inst* 80:1553-1559; Morrison (1985) *Science* 229:1202-1207; Oi *et al.* (1986) *BioTechniques* 4:214; U.S. Pat. No. 5,225,539; Jones *et al.* (1986) *Nature* 321:552-525; Verhoeyan *et al.* (1988) *Science* 239:1534; and Beidler *et al.* (1988) *J Immunol* 141:4053-4060.

In one embodiment, methodologies for the screening of antibodies that possess the desired specificity include, but are not limited to, enzyme-linked immunosorbent assay (ELISA) and other immunologically-mediated techniques known within the art.

Anti-polymorphic protein antibodies may be used in methods known within the art relating to the detection, quantitation and/or cellular or tissue localization of a polymorphic protein (*e.g.*, for use in measuring levels of the polymorphic protein within appropriate physiological samples, for use in diagnostic methods, for use in imaging the protein, and the like). In a given embodiment, antibodies for polymorphic proteins, or derivatives, fragments, analogs or homologs thereof, that contain the antibody-derived CDR, are utilized as pharmacologically-active compounds in therapeutic applications intended to treat a pathology in a subject that arises from the presence of the cSNP allele in the subject.

An anti-polymorphic protein antibody (*e.g.*, monoclonal antibody) can be used to isolate polymorphic proteins by a variety of immunochemical techniques, such as immunoaffinity chromatography or immunoprecipitation. An anti-polymorphic protein antibody can facilitate the purification of natural polymorphic protein from cells and of recombinantly produced polymorphic proteins expressed in host cells. Moreover, an anti-polymorphic protein antibody can be used to detect polymorphic protein (*e.g.*, in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the polymorphic protein. Anti-polymorphic antibodies can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, *e.g.*, to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (*i.e.*, physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase,

- g a lactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a
- 5 luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

Seq ID	CuraGen sequence ID	Base pos. of SNP	Polymorphic sequence	Base before	Base after	Amino acid before	Amino acid after	Type of change	Protein classification of CuraGen gene	Name of protein identified following a BLASTX analysis of the CuraGen sequence	Similarity (pValue) following a BLASTX analysis	Map location
1	cg43333349	1008	COCTGACAGGGGA GTCTGAGCCACAA /GACCCGCTCACC CGAGTGCAAGCAC G	A	G	Gln	Gln	SILENT- CODING	ATPase associat ed	Human Gene SWISSPROT- ID:P20648 POTASSIUM- TRANSPORTING ATPASE ALPHA CHAIN (EC 3.6.1.36) (PROTON PUMP) (GASTRIC H+/K+ ATPASE ALPHA SUBUNIT) - HOMO SAPIENS (HUMAN), 1035 aa.	0	19
2	cg43931765	2296	ATGGATAGTCCAT CTGGTTGGATGCIA /TGTGTAAGTGTG GCCTCGTTCAAGT	A	T	Thr	Thr	SILENT- CODING	cadherin	Human Gene SWISSPROT- ID:P18084 INTEGRIN BETA-5 SUBUNIT PRECURSOR - HOMO SAPIENS (HUMAN), 799 aa.	0	3

3	cg44130533	1832	AATACAAAGCTGA GTGGAGAGCAGTT /GIGGTGAAGAAGT ATGGCATTCCAAG T	T	G	Val	Val	SILENT- CODING	cadherin	Human Gene SWISSNEW- ID:P13591 NEURAL CELL ADHESION MOLECULE, 140 KD ISOFORM PRECURSOR (N- CAM 140) (NCAM-140) (CD56 ANTIGEN) - HOMO SAPIENS (HUMAN), 848 aa. peptide:SWISSPROT-ID:P13591 NEURAL CELL ADHESION MOLECULE, 140 KD ISOFORM PRECURSOR (N-CAM 140) (NCAM-140) (CD56 ANTIGEN) - HOMO SAPIENS (HUMAN), 848 aa.	0	11
4	cg34888922	2330	TATTGTTATTATGT ATTCTGTTTAC A/G TGTTTCTGTGTC CTGCTAAGAGAA	A	G	Thr	Thr	SILENT- CODING	cadherin	Human Gene SWISSNEW- ID:Q08554 DESMOCOLLIN 1A/1B PRECURSOR (DESMOSOMAL GLYCOPROTEIN 2/3) (DG2/DG3) - HOMO SAPIENS (HUMAN), 894 aa. peptide:SWISSPROT-ID:Q08554 DESMOCOLLIN 1A/1B PRECURSOR (DESMOSOMAL GLYCOPROTEIN 2/3) (DG2 / DG3) - HOMO SAPIENS (HUMAN), 894 aa.	0	18

5	cg4888922	815	CAAGGAGCATTTGA CCGTGAGAAATAIC /TGAACAGTTTGC GTTATATGGCTATG	C	T	Tyr	Tyr	SILENT- CODING	cadherin	Human Gene SWISSNEW- ID:Q08554 DESMOCOLLIN 1A/1B PRECURSOR (DESMOSOMAL GLYCOPROTEIN 2/3) (DG2/DG3) - HOMO SAPIENS (HUMAN), 894 aa. pcis: SWISSPROT-ID:Q08554 DESMOCOLLIN 1A/1B PRECURSOR (DESMOSOMAL GLYCOPROTEIN 2/3) (DG2 / DG3) - HOMO SAPIENS (HUMAN), 894 aa.	0	18
6	cg40310734	1172	TGGCGTCGTATTIT GGGCATTTCAGTIG/ CJGCTGTCACCTGAC GTCAACGGGGATG	G	C	Val	Val	SILENT- CODING	cadherin	Human Gene SWISSPROT- ID:P08514 PLATELET MEMBRANE GLYCOPROTEIN IIB PRECURSOR (GPIIB) (INTEGRIN ALPHA- IIB) (CD41) - HOMO SAPIENS (HUMAN), 1039 aa.	0	17 (17q21.3 2)
7	cg40310734	2243	AGGGGGCCTATGA AGCAGAGCTGGC[C /GJGTGCACCTGCC CCAGGGCGCCCCAC T	C	G	Ala	Ala	SILENT- CODING	cadherin	Human Gene SWISSPROT- ID:P08514 PLATELET MEMBRANE GLYCOPROTEIN IIB PRECURSOR (GPIIB) (INTEGRIN ALPHA- IIB) (CD41) - HOMO SAPIENS (HUMAN), 1039 aa.	0	17 (17q21.3 2)
8	cg40310734	812	GTTACTGTGAAGC GGGCTTCAGCTC[C/ GJGTGTCACCTCAG GCCGGAGAGCTGG	C	G	Ser	Ser	SILENT- CODING	cadherin	Human Gene SWISSPROT- ID:P08514 PLATELET MEMBRANE GLYCOPROTEIN IIB PRECURSOR (GPIIB) (INTEGRIN ALPHA- IIB) (CD41) - HOMO SAPIENS (HUMAN), 1039 aa.	0	17 (17q21.3 2)

9	cg43331935	1922	GTGACAAAGTACTT CATAGAGGATGGI G/TJCGCCTGGTCAT CCACAGCCTGGAC T	G	T	Gly	Gly	SILENT- CODING	cadherin	Human Gene SWISSPROT- ID:P32004 NEURAL CELL ADHESION MOLECULE L1 PRECURSOR (N-CAM L1) - HOMO SAPIENS (HUMAN), 1257 aa.	0	X
10	cg42388009	383	AAGGAGAAACAA TGAAAGAACCGAAI C/TJGAAGACGAAG ACTCTGAGGCTGA GA	C	T	Asn	Asn	SILENT- CODING	cadherin	Human Gene SWISSPROT- ID:P21815 BONE SIALOPROTEIN II PRECURSOR (BSP II) (CELL-BINDING SIALOPROTEIN) (INTEGRIN- BINDING SIALOPROTEIN) - HOMO SAPIENS (HUMAN), 317 aa.	7.00E-172	4
11	cg42388009	389	AAAACAAATGAAGA ACCGAACGAAGAI C/TJGAAGACTCTG AGGCTGAGAATAC CA	C	T	Asp	Asp	SILENT- CODING	cadherin	Human Gene SWISSPROT- ID:P21815 BONE SIALOPROTEIN II PRECURSOR (BSP II) (CELL-BINDING SIALOPROTEIN) (INTEGRIN- BINDING SIALOPROTEIN) - HOMO SAPIENS (HUMAN), 317 aa.	7.00E-172	4
12	cg44126574	1289	AGAACGGCCAGCC CCTGTGGATCCTTC/ GJGGGGATGCTCTC CTCAGGTCCTACT	C	G	Leu	Leu	SILENT- CODING	cathepsin	Human Gene SPTREMBL- ID:Q64411 PROGASTRICIN PRECURSOR (EC 3.4.23.3) (PEPSIN C) - CAVIA PORCELLUS (GUINEA PIG), 394 aa.	8.00E-155	6 (6p21.3)

13	cg43970983	3066	GGACTCCAGTGTG CAGGGCATCCAG[C /T]TACATCCTATCC TGGCGGCCACTCA	C	T	Ser	Ser	SILENT- CODING	collagen	Human Gene SWISSPROT- ID:Q02388 COLLAGEN ALPHA 1(VII) CHAIN PRECURSOR (LONG-CHAIN COLLAGEN) (LC COLLAGEN) - HOMO SAPIENS (HUMAN), 2944 aa.	0	3 (3p21.3)
14	cg44032748	245	TAAGACGGGCAGC TACACCGGCAGC[A /G]GTTACCTGCCA GCTGAGCAACTGG T	A	G	Ala	Ala	SILENT- CODING	complement	Human Gene SWISSPROT- ID:P07357 COMPLEMENT COMPONENT C8 ALPHA CHAIN PRECURSOR - HOMO SAPIENS (HUMAN), 584 aa.	0	1 (1p32)
15	cg41553795	222	TCCAGGCCAAGGC CAATTTTGATGCT/ GJCAGCAGTTTGCA GGGACCTGGCTCC	T	G	Ala	Ala	SILENT- CODING	complement	Human Gene Homologous to SWISSPROT-ID:P07360 COMPLEMENT C8 GAMMA CHAIN PRECURSOR - HOMO SAPIENS (HUMAN), 202 aa.	1.40E-104	9 (9q34.3)
16	cg43942011	1371	AGGTAGGAGGGCT TGGTCTCCAAAC[A /G]CCTATTGTTCA TTCTCCACAGTGC	A	G	Gly	Gly	SILENT- CODING	complementrecep pt	Human Gene Similar to TREMBLNEW-ID:E246058 COMPLEMENT RECEPTOR 2 - MUS MUSCULUS (MOUSE), 651 aa (fragment).	1.10E-69	1 (1q32)
17	cg21644442	1219	AACAGCCGGCAGA TGTAACCTGGTAC[A /C]GCCTTGCCAG GGTGGGCCCCCGTG A	A	C	Thr	Thr	SILENT- CODING	csf	Human Gene SWISSPROT- ID:P09603 MACROPHAGE COLONY STIMULATING FACTOR-1 PRECURSOR (CSF-1) (MCSF) - HOMO SAPIENS (HUMAN), 554 aa.	5.00E-304	1 (1p21)

18	cg41533258	597	TCCAGGCGCCGGGC AGGAGGGGTCCT[G /A]GTGCTCCCAT CTGCAGAGCTTCC	G	A	Leu	Leu	SILENT- CODING	csf	Human Gene Homologous to SWISSPROT-ID:P09919 GRANULOCYTE COLONY- STIMULATING FACTOR PRECURSOR (G-CSF) (PLURIPOTIN) - HOMO SAPIENS (HUMAN), 207 aa.	1.50E-107	17 (17q11.2)
19	cg43996714	1743	ATGTGCCCACTGC ATTGGGTGTCCIA/ GIGGAGTTGATACT GGTGGGATCACAG	A	G	Pro	Pro	SILENT- CODING	dehydrogenase	Human Gene TREMBLNEW- ID:G2979625 PYRUVATE DEHYDROGENASE COMPLEX PROTEIN X SUBUNIT PRECURSOR - HOMO SAPIENS (HUMAN), 501 aa.	1.60E-266	11
20	cg43259523	366	CAGAAATATGGAGG CACAGGAGCTTCTA /TTTTTATCCACT GTGCTCGTGATAG	A	T	Ser	Ser	SILENT- CODING	dehydrogenase	Human Gene SWISSPROT- ID:P45954 ACYL-COA DEHYDROGENASE, SHORT/BRANCHED CHAIN SPECIFIC PRECURSOR (EC 1.3.99.-) (SBCAD) (2-METHYL BRANCHED CHAIN ACYL-COA DEHYDROGENASE) (2- MEBCAD) - HOMO SAPIENS (HUMAN), 432 aa.	2.00E-229	10 (10q25)

21	cg43057018	1528	GAATAAGAAATTC AATCTGGATGCAIC /TTGGTGACCCATA CCCTGCCTTTTGA	C	T	Leu	Leu	SILENT- CODING	dehydrogenase	Human Gene SWISSNEW- ID:P08319 ALCOHOL DEHYDROGENASE CLASS II PI CHAIN (EC 1.1.1.1) - HOMO SAPIENS (HUMAN), 391 aa. lpcis:SWISSPROT-ID:P08319 ALCOHOL DEHYDROGENASE CLASS II PI CHAIN (EC 1.1.1.1) - HOMO SAPIENS (HUMAN), 391 aa.	1.30E-209	4 (4q22)
22	cg1395871	430	GTGCTCCAGAGGG GCCAGCAGGCACI A/GJGGAAAAACCG AAACCAACCAAGGA CT	A	G	Thr	Thr	SILENT- CODING	dynein	Human Gene Homologous to SPTREMBL-ID:Q92816 CYTOPLASMIC DYNEIN 3 HEAVY CHAIN - HOMO SAPIENS (HUMAN), 197 aa (fragment).	2.50E-103	
23	cg1395871	436	CAGAGGGGCCAGC AGGCACAGGAAAI A/GJACCGAAACCA CCAAGGACTTGGC TA	A	G	Lys	Lys	SILENT- CODING	dynein	Human Gene Homologous to SPTREMBL-ID:Q92816 CYTOPLASMIC DYNEIN 3 HEAVY CHAIN - HOMO SAPIENS (HUMAN), 197 aa (fragment).	2.50E-103	
24	cg1395871	542	AGCAATGGGAAAG TTTTTTAAAGGAIC/ TJTGCTTCTCTG GTGCTTGGGCTTG	C	T	Leu	Leu	SILENT- CODING	dynein	Human Gene Homologous to SPTREMBL-ID:Q92816 CYTOPLASMIC DYNEIN 3 HEAVY CHAIN - HOMO SAPIENS (HUMAN), 197 aa (fragment).	2.50E-103	

25	cg1395871	571	C	T	Phe	Phe	SILENT-CODING	dyneln	Human Gene Homologous to SP/TREMBL-ID:Q92816 CYTOPLASMIC DYNEIN 3 HEAVY CHAIN - HOMO SAPIENS (HUMAN), 197 aa (fragment).	2.50E-103	
26	cg43950268	1269	G	A	Asp	Asp	SILENT-CODING	eph	Human Gene TREMBLNEW- ID:G2865466 HEAT SHOCK PROTEIN 75 - HOMO SAPIENS (HUMAN), 649 aa.	0	16
27	cg43918531	461	C	T	Arg	Arg	SILENT-CODING	eph	Human Gene Homologous to SWISSNEW-ID:Q52500 THERMOSOME SUBUNIT (HEAT-SHOCK PROTEIN) - PYROCOCCLUS KODAKARAENSIS, 546 aa. pels:SWISSPROT-ID:Q52500 THERMOSOME SUBUNIT (HEAT-SHOCK PROTEIN) - PYROCOCCLUS SP. (STRAIN KOD1), 546 aa.	1.00E-104	5

28	cg43957743	1146	CAAGTTCCTCAAT AAAGTGGCAGTTTC /TTTCAGGTTCTACT GGCTCCACTTCTC	C	T	Glu	Glu	SILENT- CODING	esterase	Human Gene SWISSNEW- ID:Q15166 SERUM PARAOXONASE/ARYLESTERA SE 3 (EC 3.1.1.2) (EC 3.1.8.1) (PON 3) (SERUM ARYLDIARYLPHOSPHATASE 3) (A-ESTERASE 3) (AROMATIC ESTERASE 3) - HOMO SAPIENS (HUMAN), 341 aa (fragment); pcis:SWISSPROT- ID:Q15166 SERUM PARAOXONASE/ARYLESTERA SE 3 (EC 3.1.1.2) (EC 3.1.8.1) (PON 3) (SERUM ARYLDIARYLPHOSPHATASE 3) (A-ESTERASE 3) (AROMATIC ESTERASE 3) - HOMO SAPIENS (HUMAN), 341 aa (fragment).	1.90E-178	
----	------------	------	---	---	---	-----	-----	-------------------	----------	--	-----------	--

29	cg43319420	963	TCACCTCAGGAG GTGGCTGTCTGTC/ TGTCCACGACAAC TACAGAAACAACC	C	T	Cys	Cys	SILENT- CODING	esterase	Human Gene Similar to SWISSNEW-ID:Q23917 3',5'- CYCLIC-NUCLEOTIDE PHOSPHODIESTERASE REGA (EC 3.1.4.17) (PDEASE REGA) - DICTYOSTELIUM DISCOIDEUM (SLIME MOLD), 793 aa. pcis:SWISSPROT- ID:Q23917 3',5'-CYCLIC- NUCLEOTIDE PHOSPHODIESTERASE REGA (EC 3.1.4.17) (PDEASE REGA) - DICTYOSTELIUM DISCOIDEUM (SLIME MOLD), 793 aa.	3.30E-60	21
30	cg3001932	1631	TCTTCAACATCGTC TAATGGCTTTAICT TTATGTGAACATAA ACATGGCCTCCC	C	T	Tyr	Tyr	SILENT- CODING	gaba	Human Gene SWISSPROT- ID:P47870 GAMMA- AMINOBUTYRIC-ACID RECEPTOR BETA-2 SUBUNIT PRECURSOR (GABA(A) RECEPTOR) - HOMO SAPIENS (HUMAN), 474 aa.	1.90E-256	5 (5q34)
31	cg43975899	370	GGATTITGGACAG ACTCCTAGATGGIC TTTATGACAATCGC CTGAGACCAAGGAT	C	T	Gly	Gly	SILENT- CODING	gaba	Human Gene SWISSPROT- ID:P14867 GAMMA- AMINOBUTYRIC-ACID RECEPTOR ALPHA-1 SUBUNIT PRECURSOR (GABA(A) RECEPTOR) - HOMO SAPIENS (HUMAN), 456 aa.	1.30E-248	5 (5q34)

32	cg43299024	1643	GGGCCCACTTCC CCTGGACGTCCA /GJTGGAAACGACCT GGACTACATGGAC T	A	G	Gln	Gln	SILENT- CODING	glucoamylase	Human Gene TREMBLNEW- ID:G2826521 MALTASE- GLUCOAMYLASE (EC 3.2.1.20) - HOMO SAPIENS (HUMAN), 1857 aa.	7.40E-199	17 (17q25.2)
33	cg43299024	2021	TGAACGAGCCTTC CAACTTCATCAGIG /AJGGCTCTGAGGA CGGCTGCCCAAC A	G	A	Arg	Arg	SILENT- CODING	glucoamylase	Human Gene TREMBLNEW- ID:G2826521 MALTASE- GLUCOAMYLASE (EC 3.2.1.20) - HOMO SAPIENS (HUMAN), 1857 aa.	7.40E-199	17 (17q25.2)
34	cg43969076	443	AATTCCAAATGAG CTCTCCAACCAACIG/ AJTATTTTCTGCGT TTTTGATCCAGAC	G	A	Tyr	Tyr	SILENT- CODING	glucuronidase	Human Gene SWISSPROT- ID:P08236 BETA- GLUCURONIDASE PRECURSOR (EC 3.2.1.31) (BETA-G1) - HOMO SAPIENS (HUMAN), 651 aa.	0	7 (7q21.11)
35	cg43969014	325	AATTCCAGATGAG CTCTCCAACCAACIG/ AJTATTTTCTGCGT TTTTGATCCAGAC	G	A	Tyr	Tyr	SILENT- CODING	glucuronidase	Human Gene Similar to SWISSPROT-ID:P08236 BETA- GLUCURONIDASE PRECURSOR (EC 3.2.1.31) (BETA-G1) - HOMO SAPIENS (HUMAN), 651 aa.	7.40E-80	5
36	cg43065549	880	GGACCATCTCTGT GACCACACCTGCIG /AJGACGCTGTCATT GGCCACTACTCGC	G	A	Ala	Ala	SILENT- CODING	glycoprotein	Human Gene SWISSPROT- ID:P16452 ERYTHROCYTE MEMBRANE PROTEIN BAND 4.2 (P4.2) (FALLIDIN) - HOMO SAPIENS (HUMAN), 690 aa.	0	15 (15q15)

37	cg43063549	991	ACCCCTGGAATAG AGAGGATGCTGT /GJTTCCTGAAGAA TGAGGGCTCAGCGC A	T	G	Val	Val	SILENT- CODING	glycoprotein	Human Gene SWISSPROT- ID:P16452 ERYTHROCYTE MEMBRANE PROTEIN BAND 4.2 (P4.2) (PALLIDIN) - HOMO SAPIENS (HUMAN), 690 aa.	0	15 (15q15)
38	cg44004239	1141	TGACGTCATCCAT GTCCAATGTCCAIC/ TJACCATGGCCCCC CCAAAATGCTCTC	C	T	Val	Val	SILENT- CODING	glycoprotein	Human Gene SWISSPROT- ID:Q12889 OVIDUCT-SPECIFIC GLYCOPROTEIN PRECURSOR (OVIDUCTAL GLYCOPROTEIN) (OVIDUCTIN) (ESTROGEN- DEPENDENT OVIDUCT PROTEIN) - HOMO SAPIENS (HUMAN), 678 aa.	0	
39	cg44004239	1846	GAAGGGATATAAC TGAAGCAATAAAIC /TJTTTCACGGTIG GCAAATGTGGACA	C	T	Lys	Lys	SILENT- CODING	glycoprotein	Human Gene SWISSPROT- ID:Q12889 OVIDUCT-SPECIFIC GLYCOPROTEIN PRECURSOR (OVIDUCTAL GLYCOPROTEIN) (OVIDUCTIN) (ESTROGEN- DEPENDENT OVIDUCT PROTEIN) - HOMO SAPIENS (HUMAN), 678 aa.	0	
40	cg43957605	1677	AGGACTGTTTTTCA TTCAGCTTCAGIAC JGTGATCCCATGG GCTCTTCTGTGA	A	C	Thr	Thr	SILENT- CODING	glycoprotein	Human Gene SWISSPROT- ID:Q00013 55 KD ERYTHROCYTE MEMBRANE PROTEIN (P55) - HOMO SAPIENS (HUMAN), 466 aa.	3.10E-249	X (Xq28)

41	cg40915005	1229	ATGTCCTCAGGATTC TACCCAAAGCCIC/ TGTGTGGGTGATG TGGATGCGGGGTG	C	T	Pro	Pro	SILENT- CODING	glycoprotein	Human Gene SWISSNEW- ID:P06126 T-CELL SURFACE GLYCOPROTEIN CD1A PRECURSOR (CD1A ANTIGEN) (T-CELL SURFACE ANTIGEN T6/LEU-6) (HTA1 THYMOCYTE ANTIGEN) - HOMO SAPIENS (HUMAN), 327 aa.lpcis:SWISSPROT-ID:P06126 T-CELL SURFACE GLYCOPROTEIN CD1A PRECURSOR (CD1A ANTIGEN) (T-CELL SURFACE ANTIGEN T6/LEU-6) (HTA1 THYMOCYTE ANTIGEN) - HOMO SAPIENS (HUMAN), 327 aa.	2.00E-183	1 (1q21)
42	cg40356255	1210	TGGCAATAATAGT GCCTTCCTTGCTIC/ TCTTTGCTATGC CTTGCATTATGGT	C	T	Leu	Leu	SILENT- CODING	glycoprotein	Human Gene SWISSNEW- ID:P29016 T-CELL SURFACE GLYCOPROTEIN CD1B PRECURSOR (CD1B ANTIGEN) - HOMO SAPIENS (HUMAN), 333 aa.lpcis:SWISSPROT- ID:P29016 T-CELL SURFACE GLYCOPROTEIN CD1B PRECURSOR (CD1B ANTIGEN) - HOMO SAPIENS (HUMAN), 333 aa.	6.70E-183	1 (1q21)

43	cg44004667	1183	CTGTGATATCTACA TCTGGGCGCCCTCT TTGGCCGGGACTTG TGGGTCCTTCT	C	T	Leu	Leu	SILENT- CODING	glycoprotein	Human Gene Homologous to SWISSNEW-ID:P01732 T-CELL SURFACE GLYCOPROTEIN CD8 ALPHA CHAIN PRECURSOR (T-LYMPHOCYTE DIFFERENTIATION ANTIGEN T8/LEU-2) - HOMO SAPIENS (HUMAN), 235 aa. Jccls: SWISSPROT-ID: P01732 T-CELL SURFACE GLYCOPROTEIN CD8 ALPHA CHAIN PRECURSOR (T- LYMPHOCYTE DIFFERENTIATION ANTIGEN T8/LEU-2) - HOMO SAPIENS (HUMAN), 235 aa.	7.60E-127	
44	cg43068999	544	AGGTCCTGCGACA GGTTACTTTGTG/ AAGAAGCTCAGCCC AAGATTGTCCTGG	G	A	Val	Val	SILENT- CODING	glycoprotein	Human Gene Homologous to SWISSPROT-ID: P02743 SERUM AMYLOID P-COMPONENT PRECURSOR (SAP) (9.5S ALPHA-1-GLYCOPROTEIN) - HOMO SAPIENS (HUMAN), 223 aa.	1.60E-119	1 (1q21)
45	cg41568631	1242	ATGGCCAGTGTCTG GGTCCTTGCTGGC/ AATGACCACACACA GTGCTGGCGCTGCC	C	A	Gly	Gly	SILENT- CODING	glycoprotein	Human Gene Similar to SWISSPROT-ID: P16452 ERYTHROCYTE MEMBRANE PROTEIN BAND 4.2 (P4.2) (PALLIDIN) - HOMO SAPIENS (HUMAN), 690 aa.	9.90E-70	14 (14q11.2)

46	cg41568631	1545	GCTCTGTGGAGTC CATCAAGAAATGG[C /G]CTGGTCTACATG AAGTACGACACGC	C	G	Gly	Gly	SILENT- CODING	glycoprotein	Human Gene Similar to SWISSPROT-ID:P16452 ERYTHROCYTE MEMBRANE PROTEIN BAND 4.2 (P4.2) (PALLIDIN) - HOMO SAPIENS (HUMAN), 690 aa.	9.90E-70	14 (14q11.2)
47	cg41603916	361	GCATCCAGTGGGT AGGGGACCCCTCG[C /T]TGGAAGGATGG CTCCATTGTCATAC	C	T	Arg	Arg	SILENT- CODING	glycoprotein	Human Gene Similar to SPTREMBL-ID:Q91406 IP1=CNS MYELIN P0-LIKE GLYCOPROTEIN - UNKNOWN, 202 aa.	3.00E-52	1 (1q22)
48	cg41603916	409	TACACAACTAGA CTACAGTGACAA[T /C]GGCACGTTCACT TGTGACGTCAAAA	T	C	Asn	Asn	SILENT- CODING	glycoprotein	Human Gene Similar to SPTREMBL-ID:Q91406 IP1=CNS MYELIN P0-LIKE GLYCOPROTEIN - UNKNOWN, 202 aa.	3.00E-52	1 (1q22)
49	cg34317662	465	AGTCCCTTCTCCGT GGCACCTACGC[G/ C]TATGGTTTGTAG AAGCCCTCTGCCA	G	C	Ala	Ala	SILENT- CODING	helicase	Human Gene Homologous to SWISSPROT-ID:Q12099 PROBABLE ATP-DEPENDENT RNA HELICASE PAL1 - SACCHAROMYCES CEREVISIAE (BAKER'S YEAST), 399 aa.	3.60E-120	17
50	cg43983917	1353	AGTCTTACTTTGCC ATTAAACCACAA[C/ T]CCCGACGCCAAG GACTTGAAGCAGC	C	T	Asn	Asn	SILENT- CODING	homeobox	Human Gene SWISSPROT- ID:P50458 HOMEBOX PROTEIN LH-2 - HOMO SAPIENS (HUMAN), 423 aa.	4.30E-216	

51	cg43983917	1359	AC TTGCCA TTAA C CACAACCCG A[C/ TGCCAAGGACTTG AAGCAGCTCGGC	C	T	Asp	SILENT- CODING	homeobox	Human Gene SWISSPROT- ID:P50458 HOMEBOX PROTEIN LH-2 - HOMO SAPIENS (HUMAN), 423 aa.	4.30E-216	
52	cg42730678	979	TGGAGCGAGCGTG GATCCAGTTGCGIG /TJGCGGGTTGTTT GGGTCAAGTTGCT	G	T	Ala	SILENT- CODING	homeobox	Human Gene SWISSPROT- ID:P28356 HOMEBOX PROTEIN HOX-D9 (HOX-4C) (HOX-5.2) - HOMO SAPIENS (HUMAN), 342 aa.	2.60E-188	2
53	cg42714160	689	GTTACCAGACGCT GGAGCTGGAGAA[G/AJGAGTTTCACT ACAATCGCTACCT GA	G	A	Lys	SILENT- CODING	homeobox	Human Gene Homologous to SWISSPROT-ID:P17509 HOMEBOX PROTEIN HOX-B6 (HOX-2B) (HOX-2.2) (HU-2) - HOMO SAPIENS (HUMAN), 224 aa.	1.10E-123	
54	cg43959084	810	TCAGGTAGCGATT GTAGTGAATTCIT/ C/TTCCTCAGCTCC AGGCTCTGGTAGC	T	C	Lys	SILENT- CODING	homeobox	Human Gene Homologous to SWISSPROT-ID:P09629 HOMEBOX PROTEIN HOX-B7 (HOX-2C) (HHO.C1) - HOMO SAPIENS (HUMAN), 217 aa.	1.30E-113	
55	cg42359655	1124	GGGAAGCATTTGC CAATCAGTCCAGIA /GJGCGGAAAGGGA TGCCITCTCTGCAGG	A	G	Arg	SILENT- CODING	hydrolase	Human Gene SWISSPROT- ID:P09848 LACTASE- PHLORIZIN HYDROLASE PRECURSOR (EC 3.2.1.108) (EC 3.2.1.62) (LACTASE- GLYCOSYLCEAMIDASE) - HOMO SAPIENS (HUMAN), 1927 aa.	0	2 (2q21)

56	cg42359655	2468	ACAGCCAGCGGTT TGGCCTGCACCA[C /TJGTCAACTTCAGC GACAGCAGCAAGT	C	T	His	His	SILENT- CODING	hydrolase	Human Gene SWISSPROT- ID:P09848 LACTASE- PHLORIZIN HYDROLASE PRECURSOR (EC 3.2.1.108) (EC 3.2.1.62) (LACTASE- GLYCOSYL CERAMIDASE) - HOMO SAPIENS (HUMAN), 1927 aa.	0	2 (2q21)
57	cg42359655	4340	ATCTGGTCACCCGTG CAGAACCTGGG[C/ TJGTGCCCACTAC CGTTTTTCCATCT	C	T	Gly	Gly	SILENT- CODING	hydrolase	Human Gene SWISSPROT- ID:P09848 LACTASE- PHLORIZIN HYDROLASE PRECURSOR (EC 3.2.1.108) (EC 3.2.1.62) (LACTASE- GLYCOSYL CERAMIDASE) - HOMO SAPIENS (HUMAN), 1927 aa.	0	2 (2q21)
58	cg43998672	1329	TGGTGTGGGCCTT GGTGAACCTAG[C /AJACGCGGCTAAT GTCTCCTGGTTTGG	C	A	Val	Val	SILENT- CODING	hydroxysteroid	Human Gene SPTREMBL- ID:Q13194 11-BETA- HYDROXYSTEROID DEHYDROGENASE TYPE 2 - HOMO SAPIENS (HUMAN), 405 aa.	2,00E-220 (16q22)	16 (16q22)
59	cg43922672	1689	GGAAGCTGACTCC AGAGGCCATGCC[C /TJGACCTCAACTCC TCCACTGACTCTG	C	T	Pro	Pro	SILENT- CODING	interleukin	Human Gene TREMBLNEW- ID:G2114410 INTERLEUKIN-16 - HOMO SAPIENS (HUMAN), 631 aa.	0	15

60	cg42908571	630	GTAGTGAGGAACA AGCCAGAGCTGTG /C/CAGATGAGTAC AAAAGTCTTGATC C	G	C	Val	Val	SILENT- CODING	interleukin	Human Gene Homologous to SWISSPROT-ID:P05231 INTERLEUKIN-6 PRECURSOR (IL-6) (B-CELL STIMULATORY FACTOR 2) (BSF-2) (INTERFERON BETA-2) (HYBRIDOMA GROWTH FACTOR) - HOMO SAPIENS (HUMAN), 212 aa.	3.40E-108	7 (7p21)
61	cg43942050	181	AGTTGGAAGTGAA TGGATCGCAGCA(C /T/TCACTGACCTGT GCITTTGAGGACC	C	T	His	His	SILENT- CODING	interleukinrecept	Human Gene SWISSNEW- ID:P16871 INTERLEUKIN-7 RECEPTOR ALPHA CHAIN PRECURSOR (IL-7R-ALPHA) (CDW127) (CD127 ANTIGEN) - HOMO SAPIENS (HUMAN), 459 aa.lpcis:SWISSPROT-ID:P16871 INTERLEUKIN-7 RECEPTOR ALPHA CHAIN PRECURSOR (IL-7R-ALPHA) (CDW127) (CD127 ANTIGEN) - HOMO SAPIENS (HUMAN), 459 aa.	3.10E-249	5 (5p13)

62	cg43145505	1249	TAAATATTCGAGA CAITGACAAGATTC TTTATGTTTCAACA GGTATCTACCATG	C	T	Ile	Ile	SILENT- CODING	kinase	Human Gene SWISSNEW- ID:P42336 PHOSPHATIDYLINOSITOL 3- KINASE CATALYTIC SUBUNIT, ALPHA ISOFORM (EC 2.7.1.137) (PI3-KINASE P110 SUBUNIT ALPHA) (PTDINS-3- KINASE P110) (PI3K) - HOMO SAPIENS (HUMAN), 1068 aa.lpcis:SWISSPROT-ID:P42336 PHOSPHATIDYLINOSITOL 3- KINASE CATALYTIC SUBUNIT, ALPHA ISOFORM (EC 2.7.1.137) (PI3-KINASE P110 SUBUNIT ALPHA) (PTDINS-3- KINASE P110) (PI3K) - HOMO SAPIENS (HUMAN), 1068 aa.	0	3
63	cg43918241	1693	AGATCTTTGAGGA AGGGGAATCTGAJC TTGATGAGTTTGAC ATGGATGAGAATC	C	T	Asp	Asp	SILENT- CODING	kinase	Human Gene SPTRMBL- ID:Q63553 SNF1-RELATED KINASE - RATTUS NORVEGICUS (RAT), 746 aa.	0	3
64	cg43090990	1438	TTCTGAGGCACAT GTTTGTACATTC/ TTCAGACCAAGGA AAACCTCTTTTITG	C	T	Phe	Phe	SILENT- CODING	kinase	Human Gene SWISSPROT- ID:Q04759 PROTEIN KINASE C, THETA TYPE (EC 2.7.1.-) (NPKC-THETA) - HOMO SAPIENS (HUMAN), 706 aa.	0	10

65	cg43969763	2339	TTAGTATCATTCAC TGATGATCTAAATC GTCCTGAAAATATC CTCTTTGTAACC	A	G	Lys	Lys	SILENT- CODING	kinase	Human Gene SWISSPROT- ID:Q13627 SERINE/THREONINE-SPECIFIC PROTEIN KINASE MINIBRAIN HOMOLOG (EC 2.7.1.-) (HP86) (DYRK) - HOMO SAPIENS (HUMAN), 763 aa.	0	21 (21q22.1)
66	cg42879455	2062	AGGTATATACCAT CATGTACAGTTGTT/ CTGGCATGAGAAA GCAGATGAGCGTC	T	C	Cys	Cys	SILENT- CODING	kinase	Human Gene SWISSPROT- ID:Q06187 TYROSINE-PROTEIN KINASE BTK (EC 2.7.1.112) (BRUTON'S TYROSINE KINASE) (AGAMMAGLOBULINAEMIA TYROSINE KINASE) (ATK) (B CELL PROGENITOR KINASE) (BPK) - HOMO SAPIENS (HUMAN), 639 aa.	0	X (Xq21.3)
67	cg42659872	1744	TGGCTCCGGCTAC ACCAACATCATGTA /CJGGTGCTAAGC ATATCCTGAGACG C	A	C	Arg	Arg	SILENT- CODING	kinase	Human Gene SPTREMBL- ID:Q16715 PYRUVATE KINASE (EC 2.7.1.40) - HOMO SAPIENS (HUMAN), 587 aa (fragment).	9.80E-308	1 (1q21)

68	cg42506800	1323	GCTTGCCAAATTTCT CGTCTGTATGCA/C JAAGTACTTTCAAG GAGATCTGAATC	A	C	Ala	Ala	SILENT- CODING	kinase	Human Gene SWISSPROT- ID:Q16654 [PYRUVATE DEHYDROGENASE(LIPOAMID E)] KINASE ISOZYME 4 PRECURSOR (EC 2.7.1.99) (PYRUVATE DEHYDROGENASE KINASE ISOFORM 4) - HOMO SAPIENS (HUMAN), 411 aa.lpcis:SPTRMBL-ID:Q16654 PYRUVATE DEHYDROGENASE KINASE ISOFORM 4 - HOMO SAPIENS (HUMAN), 411 aa.	1.60E-220	7 (7q21.3)
69	cg43966621	526	CTGTGGAGTACAT GTAGCTGAAGAG/C /TJCGCTCAATCTTC CTCAAGGGAACAC	C	T	Arg	Arg	SILENT- CODING	kinase	Human Gene SWISSPROT- ID:Q15119 [PYRUVATE DEHYDROGENASE(LIPOAMID E)] KINASE ISOZYME 2 PRECURSOR (EC 2.7.1.99) (PYRUVATE DEHYDROGENASE KINASE ISOFORM 2) - HOMO SAPIENS (HUMAN), 407 aa.lpcis:SPTRMBL-ID:Q15119 PYRUVATE DEHYDROGENASE KINASE - HOMO SAPIENS (HUMAN), 407 aa.	3.80E-219	17
70	cg43917871	1448	ACATCATATTGGC GCTGCTGACGGG/C /TJGTACTGCCCCCT GGCATGCTAGATG	C	T	Thr	Thr	SILENT- CODING	kinase	Human Gene SWISSPROT- ID:P19138 CASEIN KINASE II, ALPHA CHAIN (CK II) (EC 2.7.1.37) - HOMO SAPIENS (HUMAN), AND BOS TAURUS (BOVINE), 391 aa.	2.00E-215	11 (20p13)

71	cg43917871	1526	CAGTGTAGAAATA GGGTGCTCCATTT/ GJGCTCTCTTGCA GTAAGCCGTGACT	T	G	Ala	Ala	SILENT- CODING	kinase	Human Gene SWISSPROT- ID:P19138 CASEIN KINASE II, ALPHA CHAIN (CK II) (EC 2.7.1.37) - HOMO SAPIENS (HUMAN), AND BOS TAURUS (BOVINE), 391 aa.	2.00E-215	11 (20p13)
72	cg44131752	912	AGCTCAATGGTGG CTCTGGTGCTCIG/ AJTCCCGAAGTGAC CTGCCTGGTTCCG	G	A	Ser	Ser	SILENT- CODING	kinase	Human Gene SPTREMBL- ID:Q15599 TYROSINE KINASE ACTIVATOR PROTEIN 1 (TKA- 1) - HOMO SAPIENS (HUMAN), 450 aa.	7.80E-173	16
73	cg43969473	1765	AATTCAGCCACT CATCTATGGCAA/T/ CJGATGTGGATTCT GTGGATGTTGCAA	T	C	Asn	Asn	SILENT- CODING	kinase	Human Gene SPTREMBL- ID:Q27467 SIMILARITY TO TYROSINE-PROTEIN KINASE - CAENORHABDITIS ELEGANS, 1280 aa.	2.10E-134	11
74	cg44023829	610	AGACCCCGCGGTC CCCTGGCCAAAGCT/ /CJGTGGAGTGCTG CCAAGGGGACTGG T	T	C	Ala	Ala	SILENT- CODING	kinasereceptor	Human Gene SWISSPROT- ID:Q04771 ACTIVIN RECEPTOR TYPE I PRECURSOR (EC 2.7.1.-) (ACTR-I) (SERINE/THREONINE-PROTEIN KINASE RECEPTOR R1) (SKR1) (ACTIVIN RECEPTOR-LIKE KINASE 2) (ALK-2) (TGF-B SUPERFAMILY RECEPTOR TYPE I) (TSR-I) - HOMO SAPIENS (HUMAN), 509 aa.	7.90E-283	2

75	cg43318277	1107	CTCAGGCTTTGCAG TCATCTGGTCCG/A JCCTAGCACTCCCT CCTCTCCTCGGC	G	A	Pro	SILENT- CODING	MHC	Human Gene SPTREMBL- ID:Q02646 MHC BINDING PROTEIN 2 - HOMO SAPIENS (HUMAN), 2500 aa.	1.20E-247	6
76	cg43966144	632	TTAACACGAGGGA GCCTGTGATGCTG/ AJGCCTGCTATGTG TGGGGCTTCTATC	G	A	Leu	SILENT- CODING	MHC	Human Gene Homologous to SWISSPROT-ID:P28068 CLASS II HISTOCOMPATIBILITY ANTIGEN, M BETA CHAIN PRECURSOR - HOMO SAPIENS (HUMAN), 263 aa.	9.10E-147	6 (6p21.3)
77	cg42686658	644	CCCCTGTGATCAAT ATCACCTGGCT/A/ GJCGCAACGGCCA AACTGTCACTGAG G	A	G	Leu	SILENT- CODING	MHC	Human Gene Homologous to SWISSPROT-ID:P06340 HLA CLASS II HISTOCOMPATIBILITY ANTIGEN, DZ ALPHA CHAIN PRECURSOR (MHC DN-ALPHA) - HOMO SAPIENS (HUMAN), 250 aa.	3.70E-134	6 (6p21.3)
78	cg42686658	857	CACCACCAAGATGC CATGGAGACCCCTG /AJGCTGTGCCCTG GGCCTGGCCATCG	G	A	Leu	SILENT- CODING	MHC	Human Gene Homologous to SWISSPROT-ID:P06340 HLA CLASS II HISTOCOMPATIBILITY ANTIGEN, DZ ALPHA CHAIN PRECURSOR (MHC DN-ALPHA) - HOMO SAPIENS (HUMAN), 250 aa.	3.70E-134	6 (6p21.3)
79	cg42686658	869	CCATGGAGACCCT GGTCTGTGCCCTG/ AJGGCCTGGCCATC GGCCTGGTGGGCT	G	A	Leu	SILENT- CODING	MHC	Human Gene Homologous to SWISSPROT-ID:P06340 HLA CLASS II HISTOCOMPATIBILITY ANTIGEN, DZ ALPHA CHAIN PRECURSOR (MHC DN-ALPHA) - HOMO SAPIENS (HUMAN), 250 aa.	3.70E-134	6 (6p21.3)

80	cg42686658	881	TGGTCTGTGCCCTG GGCCTGGCCATTC/T JGGCCTGGTGGGCT TCCTCGTGGGCA	C	T	Ile	Ile	SILENT- CODING	MHC	Human Gene Homologous to SWISSPROT-ID:P06340 HLA CLASS II HISTOCOMPATIBILITY ANTIGEN, DZ ALPHA CHAIN PRECURSOR (MHC DN-ALPHA) - HOMO SAPIENS (HUMAN), 250 aa.	3.70E-134	6 (6p21.3)
81	cg42686658	893	TGGGCTGGCCAT CGGCCTGGTGGG[C /G]TTCCTCGTGGGC ACCGTCCTCATCA	C	G	Gly	Gly	SILENT- CODING	MHC	Human Gene Homologous to SWISSPROT-ID:P06340 HLA CLASS II HISTOCOMPATIBILITY ANTIGEN, DZ ALPHA CHAIN PRECURSOR (MHC DN-ALPHA) - HOMO SAPIENS (HUMAN), 250 aa.	3.70E-134	6 (6p21.3)
82	cg42686658	905	TCGGCCTGGTGGG CTTCCTCGTGGG[C/ T]ACCGTCCTCATC ATCATGGGCACAT	C	T	Gly	Gly	SILENT- CODING	MHC	Human Gene Homologous to SWISSPROT-ID:P06340 HLA CLASS II HISTOCOMPATIBILITY ANTIGEN, DZ ALPHA CHAIN PRECURSOR (MHC DN-ALPHA) - HOMO SAPIENS (HUMAN), 250 aa.	3.70E-134	6 (6p21.3)
83	cg38337333	279	GTTTCCTCATTAGC CCTGTGACCCCA/T JGCACACGCAGGG ACCTACAGATGTC	A	T	Pro	Pro	SILENT- CODING	MHC	Human Gene Homologous to SPTREMBL-ID:Q95368 HLA CLASS I INHIBITORY NK RECEPTOR - HOMO SAPIENS (HUMAN), 455 aa.	1.80E-113	19
84	cg38337333	492	TTGACATCTACCAT CTATCCAGGGA[G/ A]GGGGAAGCCCA TGAACTTAGGCTC C	G	A	Glu	Glu	SILENT- CODING	MHC	Human Gene Homologous to SPTREMBL-ID:Q95368 HLA CLASS I INHIBITORY NK RECEPTOR - HOMO SAPIENS (HUMAN), 455 aa.	1.80E-113	19

85	cg38337333	699	CTTCTAGTAGTGG CCTTCACCCACIT/A JGAACCAAGCTTCA AAACTGGTATCG	T	A	Thr	Thr	SILENT- CODING	MHC	Human Gene Homologous to SPTREMBL-ID:Q95368 HLA CLASS I INHIBITORY NK RECEPTOR - HOMO SAPIENS (HUMAN), 455 aa.	1.80E-113	19
86	cg38337333	774	GGTACTCAGTGGC CATCATCCTCTT/C TJACCATCCTTCCC TTCTTCTCCTTC	C	T	Phe	Phe	SILENT- CODING	MHC	Human Gene Homologous to SPTREMBL-ID:Q95368 HLA CLASS I INHIBITORY NK RECEPTOR - HOMO SAPIENS (HUMAN), 455 aa.	1.80E-113	19
87	cg38337333	783	TGCCATCATCCTC TTCACCATCCTT/C JCCCTCTTCTCCT TCATCGCTGGT	T	C	Leu	Leu	SILENT- CODING	MHC	Human Gene Homologous to SPTREMBL-ID:Q95368 HLA CLASS I INHIBITORY NK RECEPTOR - HOMO SAPIENS (HUMAN), 455 aa.	1.80E-113	19
88	cg43984759	649	AGGAGCTCAAGCG TGAGGCGGAGAC/C /TJCTACGGGAGCG GGAAGGCGAGGAG T	C	T	Thr	Thr	SILENT- CODING	misc_channel	Human Gene SPTREMBL- ID:Q14193 H-DRK1 K(+) CHANNEL - HOMO SAPIENS (HUMAN), 858 aa.	0	20
89	cg39660131	990	TCATGGGCAACCT AAGGCACAAGT/G C/TJGTGCGCAACTT CACAGCGCTCAAC G	C	T	Cys	Cys	SILENT- CODING	misc_channel	Human Gene SPTREMBL- ID:Q14524 SODIUM CHANNEL ALPHA SUBUNIT - HOMO SAPIENS (HUMAN), 2016 aa.	0	3 (3p24)

90	cg44963814	717	CGGAATACCTGGC CATCACCTCTGA[A/ GTAGCAAAGAGAA CTGCACGGGCGTC C	A	G	Glu	Glu	SILENT- CODING	misc_channel	Human Gene Homologous to SWISSPROT-ID:Q07699 SODIUM CHANNEL BETA-1 SUBUNIT PRECURSOR - HOMO SAPIENS (HUMAN), 218 aa.jpels:TREMBLNEW- ID:G2804300 VOLTAGE-GATED SODIUM CHANNEL BETA-1 SUBUNIT - HOMO SAPIENS (HUMAN), 218 aa.	2.20E-113	19 (19q13.1)
91	cg21413267	870	AGAGTGGCGAGTG GGTCATCGTGGGAC /TJGCCGTGGGCAC CTACAAACACCAGG A	C	T	Asp	Asp	SILENT- CODING	misc_channel	Human Gene Similar to SPTREMBL-ID:P91197 SIMILAR TO LIGAND-GATED IONIC CHANNEL PROTEIN - CAENORHABDITIS ELEGANS, 461 aa.	7.90E-79	
92	cg21413267	909	ACAACACCAAGAA GTACGAGTGTGTC /TJGCCGAGATCTA CCCGGACATCACC T	C	T	Cys	Cys	SILENT- CODING	misc_channel	Human Gene Similar to SPTREMBL-ID:P91197 SIMILAR TO LIGAND-GATED IONIC CHANNEL PROTEIN - CAENORHABDITIS ELEGANS, 461 aa.	7.90E-79	
93	cg3000465	1160	AGAGGCTCTTCTG CAGAAACTTCC[A/ CJAAATTACTTGC ATGAAAGATCATG	A	C	Pro	Pro	SILENT- CODING	misc_channel	Human Gene Similar to SPTREMBL-ID:P91197 SIMILAR TO LIGAND-GATED IONIC CHANNEL PROTEIN - CAENORHABDITIS ELEGANS, 461 aa.	6.10E-70	8 (8p11.2)

94	cg30421838	3766	GTCTAGGATGGAG ATCCTACAAACA/C /TGTGAGTGGCA GATGCTGATTTTG	C	T	His	His	SILENT- CODING	nuc_recpt	Human Gene SWISSNEW- ID:P06401 PROGESTERONE RECEPTOR (PR) - HOMO SAPIENS (HUMAN), 933 aa. pcis:SWISSPROT-ID:P06401 PROGESTERONE RECEPTOR (PR) - HOMO SAPIENS (HUMAN), 933 aa.	0	11 (11q22)
95	cg30421838	4114	ATAAATTGCATGA TCTTGTCAAACA/C G/CITTCATCTGTAC TGCTTGAATACAT	A	G	Gln	Gln	SILENT- CODING	nuc_recpt	Human Gene SWISSNEW- ID:P06401 PROGESTERONE RECEPTOR (PR) - HOMO SAPIENS (HUMAN), 933 aa. pcis:SWISSPROT-ID:P06401 PROGESTERONE RECEPTOR (PR) - HOMO SAPIENS (HUMAN), 933 aa.	0	11 (11q22)
96	cg43947341	713	TTACGTCGCCAAA TTCCCAGGGCAC/C /GJTTCGACACGAA CTTCAGTACGGGA T	A	G	Asn	Asn	SILENT- CODING	nuclease	Human Gene Homologous to SWISSPROT-ID:P07992 DNA EXCISION REPAIR PROTEIN ERCC-1 - HOMO SAPIENS (HUMAN), 297 aa.	1.10E-115	
97	cg43939230	4226	TCCCTGTGACCCA GGCAGGTGCATG/C /GJTGACACTGGT CGTGACCTGGCCA G	A	G	Thr	Thr	SILENT- CODING	oncogene	Human Gene SPTREMBL- ID:Q99907 LATENT TRANSFORMING GROWTH FACTOR-BETA-BINDING PROTEIN-2 - HOMO SAPIENS (HUMAN), 1821 aa.	0	14 (14q24)

98	cg42674136	1447	CGGCACACAGGCC GCTCGCGGAGC/C TGTGGCCACCCC CAGCCCCTGGCCA	C	T	Ala	Ala	SILENT- CODING	oncogene	Human Gene SWISSPROT- ID:P31314 HOMEBOX PROTEIN HOX-11 (TCL-3 PROTO-ONCOGENE) - HOMO SAPIENS (HUMAN), 330 aa.	3.70E-182	10
99	cg41972699	742	AGAACTCGCGGT CTCCCACTACAT/C/ TATCAACTCGCTG CCCAACCGCCGTT	C	T	Ile	Ile	SILENT- CODING	oncogene	Human Gene Similar to SWISSPROT-ID:Q64010 PROTO- ONCOGENE C-CRK (P38) (ADAPTER MOLECULE CRK) - MUS MUSCULUS (MOUSE), 304 aa.	2.40E-84	22 (22q11)
100	cg42849556	963	CTGCAACTACCTTG AACCAGTTGAG/C/ TTGCGGATCCACC CTCAGCAGCAGCC	C	T	Leu	Leu	SILENT- CODING	oxidase	Human Gene SWISSPROT- ID:P19878 NEUTROPHIL CYTOSOL FACTOR 2 (NCF-2) (NEUTROPHIL NADPH OXIDASE FACTOR 2) (P67- PHOX) - HOMO SAPIENS (HUMAN), 526 aa.	2.80E-287	1 (1q25)
101	cg43996195	1310	CAGCATGACCTGG CACTGTACTTCG/C/ AIGGAAAGTTGGG GATTTCACCGTAGT	G	A	Pro	Pro	SILENT- CODING	phosphorylase	Human Gene SWISSPROT- ID:P00491 PURINE NUCLEOSIDE PHOSPHORYLASE (EC 2.4.2.1) (INOSINE PHOSPHORYLASE) (PNP) - HOMO SAPIENS (HUMAN), 289 aa.	2.40E-155	

102	cg43996195	1421	TTGCAACTTGAGG TCGGTGCTTAGTG/ ATGAGACAGAAG CCATTCTGCAGTGT	G	A	His	His	SILENT- CODING	phosphorylase	Human Gene SWISSPROT- ID:P00491 PURINE NUCLEOSIDE PHOSPHORYLASE (EC 2.4.2.1) (INOSINE PHOSPHORYLASE) (PNP) - HOMO SAPIENS (HUMAN), 289 aa.	2.40E-155	
103	cg43948227	372	TTTACAGTTTCTT ACTGCATCATCTA/T JATGTCAGAAATCT GTTCCTTCAGCT	A	T	Ile	Ile	SILENT- CODING	polymerase	Human Gene Similar to SWISSNEW-ID:P53999 ACTIVATED RNA POLYMERASE II TRANSCRIPTIONAL COACTIVATOR P15 (PC4) (P14) - HOMO SAPIENS (HUMAN), 126 aa. pcis:SWISSPROT- ID:P53999 ACTIVATED RNA POLYMERASE II TRANSCRIPTIONAL COACTIVATOR P15 (PC4) (P14) - HOMO SAPIENS (HUMAN), 126 aa.	5.40E-62	5

104	cg43333426	1302	AGAGCCACTACAA GGTGGACTACTCJA /GJCGTTTTCACAAAG ACCTACGAGGTGG	A	G	Ser	Ser	SILENT- CODING	potassium_chan nel	Human Gene SWISSNEW- ID:P48050 INWARD RECTIFIER (POTASSIUM CHANNEL 4 (POTASSIUM CHANNEL, INWARDLY RECTIFYING, SUBFAMILY 1, MEMBER 4) (HIPPOCAMPAL INWARD RECTIFIER) (HIR) (HRK1) (HRK2) (KIR2.3) - HOMO SAPIENS (HUMAN), 445 aa.lpcis:SWISSPROT-ID:P48050 INWARD RECTIFIER (POTASSIUM CHANNEL 4 (POTASSIUM CHANNEL, INWARDLY RECTIFYING, SUBFAMILY 1, MEMBER 4) (HIPPOCAMPAL INWARD RECTIFIER) (HIR) (HRK1) (HRK2) (KIR2.3) - HOMO SAPIENS (HUMAN), 445 aa.	4.40E-241	
-----	------------	------	--	---	---	-----	-----	-------------------	-----------------------	--	-----------	--

105	cg43051431	1683	TCACACCTGTCTG ACCGTGGAGGAT/ CTGGTTCTACGAA GTTGACTACAACA	T	C	Asp	SILENT- CODING	potassium_chan nel	Human Gene SWISSPROT- ID:P48051 G PROTEIN- ACTIVATED INWARD RECTIFIER POTASSIUM CHANNEL 2 (GIRK2) (POTASSIUM CHANNEL, INWARDLY RECTIFYING, SUBFAMILY 1, MEMBER 6) (KATP-2) (BIR1) (KIR3.2) - HOMO SAPIENS (HUMAN), 423 aa. lpcis: TREMBLNEW- ID:G1518526 INWARDLY RECTIFYING POTASSIUM CHANNEL KIR3.2 - HOMO SAPIENS (HUMAN), 423 aa.	1.60E-227	16
106	cg43920929	1081	GCAGGATCACCTG CACCTCTTGGG[C/ GIACCATGATGCTC ATCCAGCTGTCTA	C	G	Val	SILENT- CODING	proteaseinhib	Human Gene SWISSPROT- ID:P07093 GLIA DERIVED NEXIN PRECURSOR (GDN) (PROTEASE NEXIN 1) (PN-1) (PROTEASE INHIBITOR 7) - HOMO SAPIENS (HUMAN), 398 aa.	1.20E-208	2
107	cg43059041	624	AGTCAGACACCCAG CTTAGAAATGAC[C /TATGGGCAATGC CTTGTTCTTGATG	C	T	Thr	SILENT- CODING	proteaseinhib	Human Gene Similar to SWISSPROT-ID:P17475 ALPHA- 1-ANTIPROTEINASE PRECURSOR (ALPHA-1- ANTITRYPSIN) (ALPHA-1- PROTEINASE INHIBITOR) - RATTUS NORVEGICUS (RAT), 411 aa.	4.40E-83	14 (14q32.1)

108	cg40148056	1385	GGAGGACAGGCAA CTCATCACCGAA[C /TITAGTCATCAGCA AGATGAACCAAGCT	C	T	Leu	Leu	SILENT- CODING	struct	Human Gene SPTREMBL- ID:Q92777 SYNAPSIN IIB - HOMO SAPIENS (HUMAN), 478 aa.	2.90E-260	3 (3p)
109	cg42894986	1002	ACCCGTTCTTCTGC CCACCCACTGA[G/ AJGCCCCAGACCGT GACTTCTTGGTGG	G	A	Glu	Glu	SILENT- CODING	struct	Human Gene SPTREMBL- ID:Q28686 50-KDA DYSTROPHIN-ASSOCIATED GLYCOPROTEIN PRECURSOR - ORYCTOLAGUS CUNICULUS (RABBIT), 387 aa.	1.40E-180	17
110	cg43961212	2160	TCTGGAAAGCCGGA CATCCTCTGAGC[A/ GJAGTCGACTGATC CGCTGGCGAACC A	A	G	Leu	Leu	SILENT- CODING	struct	Human Gene Homologous to TREMBLNEW-ID:G1703715 PANTOPHYISIN=SYNAPTOPY SIN HOMOLOG - MUS SP, 261 aa.	2.40E-114	7
111	cg42898003	497	TCATCAGAGATTC GATCTCCTCGTC[C/ AJGTCACGTGCTCC CCGGAGGCCCTGA	C	A	Thr	Thr	SILENT- CODING	struct	Human Gene Similar to SWISSPROT-ID:P02585 TROPONIN C, SKELETAL MUSCLE - HOMO SAPIENS (HUMAN), 159 aa.	1.50E-80	20 (20q12)
112	cg43960684	788	GCCTTGAGGAGGA GGCGCGGTTCGG[C /GJGACGACACTGA GGCGGCCCATCCGC G	C	G	Arg	Arg	SILENT- CODING	struct	Human Gene Similar to SWISSPROT-ID:P02535 KERATIN, TYPE I CYTOSKELETAL 10 (CYTOKERATIN 10) (56 KD CYTOKERATIN) (KERATIN, TYPE I CYTOSKELETAL 59 KD) - MUS MUSCULUS (MOUSE), 569 aa.	8.30E-58	8

113	cg43938714	1049	TTCGGAAAGGCCA AGCAGTGAACCTTG /CJATGATGGATGC CACCAATATGCCA G	G	C	Leu	Leu	SILENT- CODING	synthase	Human Gene Similar to SPREMBL-ID:Q42761, SQUALENE SYNTHASE (EC 2.5.1.21) (FARNESYL- DIPHOSPHATE FARNESYLTRANSFERASE) (FARNESYLTRANSFERASE) (PRESQUALENE-DI- DIPHOSPHATE SYNTHASE) - GLYCRRHIZA GLABRA, 412 aa.	9.20E-83	8
114	cg43124627	901	ACACCCACAGCAG TTTGGTTTAGGTA/ TJTTATCTGTAAT GGAAAGTTCTGCG	A	T	Gly	Gly	SILENT- CODING	synthase	Human Gene Similar to SWISSNEW-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE-COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa. pcis:SWISSPROT-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE-COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa.	7.70E-79	16

115	cg43968419	906	TCTTCTCCAACAGT A CTGCCACCCGCIAT TGTCGTTGGCTGCG CCTCCAAGGCC	T	Ala	Ala	SILENT- CODING	synthase	Human Gene Similar to SWISSNEW-ID:P53556 8- AMINO-7-OXONONANOATE SYNTHASE (EC 2.3.1.47) (7- KETO-8-AMINO- PELARGONIC ACID SYNTHETASE) (7-KAP SYNTHETASE) (L-ALANINE-- PIMELYL COA LIGASE) - BACILLUS SUBTILIS, 389 aa.[pcis:SWISSPROT-ID:P53556 8-AMINO-7-OXONONANOATE SYNTHASE (EC 2.3.1.47) (7- KETO-8-AMINO- PELARGONIC ACID SYNTHETASE) (7-KAP SYNTHETASE) (L-ALANINE-- PIMELYL COA LIGASE) - BACILLUS SUBTILIS, 389 aa.	9,90E-70	
-----	------------	-----	---	---	-----	-----	-------------------	----------	--	----------	--

116	cg43064068	1484	TTGTGGTCTCTGGCC TCGCAGTTCTCTG/A JTCCCATGACCCAG AACAGCTCACCA	G	A	Leu	Leu	SILENT- CODING	synthase	Human Gene Similar to SWISSNEW-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE-COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa. pcls:SWISSPROT-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE-COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa.	7.40E-65	
117	cg43064068	1622	TCACAGGGAAAT TCAACGAGCCAAIG /AJCTCGAGACAA GGAGTGAAGATG T	G	A	Lys	Lys	SILENT- CODING	synthase	Human Gene Similar to SWISSNEW-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE-COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa. pcls:SWISSPROT-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE-COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa.	7.40E-65	

118	cg41084924	1278	TGACTCTCCCGAC CCGTCCCACCA[CT] JGGTCTCCACAGCA CTCCCGACAGCC	C	T	His	His	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P14416 D(2) DOPAMINE RECEPTOR - HOMO SAPIENS (HUMAN), 443 aa.	1.70E-241	11
119	cg41084924	1662	TCCGCAAGGCCTT CCTGAAGATCCT[C] TJCACTGCTGACTC TGCTGCCCTGCCCG	C	T	Leu	Leu	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P14416 D(2) DOPAMINE RECEPTOR - HOMO SAPIENS (HUMAN), 443 aa.	1.70E-241	11
120	cg41084924	606	TCTCTGTCGCCACA CTGGTCATGCC[C/A] JGGGTTGTCTACC TGGAGGTGGTAG	C	A	Pro	Pro	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P14416 D(2) DOPAMINE RECEPTOR - HOMO SAPIENS (HUMAN), 443 aa.	1.70E-241	11
121	cg43985000	1471	TTGCTCTTTGCTGG TTCCCTCTTCA[CT] TTAAGCCGTATATT GAAGAAAACTG	C	T	His	His	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P25101 ENDOTHELIN-1 RECEPTOR PRECURSOR (ET-A) - HOMO SAPIENS (HUMAN), 427 aa.	1.60E-236	4
122	cg43985000	1507	TATTGAAGAAAAC TGTGTATAACGAIA /GJATGGACAAGAA CCGATGTGAATT C	A	G	Glu	Glu	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P25101 ENDOTHELIN-1 RECEPTOR PRECURSOR (ET-A) - HOMO SAPIENS (HUMAN), 427 aa.	1.60E-236	4

123	cg44930578	561	ACGTGAACACCGA CACTCTACTCCAAIG/ A/GTGCTGGTGACC GCCGTGTACCTGG	G	A	Lys	Lys	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P30989 NEUROTENSIN RECEPTOR TYPE 1 (NT-R-1) (HIGH-AFFINITY LEVOCABASTINE- INSENSITIVE NEUROTENSIN RECEPTOR) (NTRH) - HOMO SAPIENS (HUMAN), 418 aa.	5.00E-217	
124	cg3003519	1263	ATTCCTTGATTGCT AGGACCCCTTTC/T TAAAAGCACCCCTGA ACATACCTACTG	C	T	Tyr	Tyr	SILENT- CODING	tm7	Human Gene SWISSNEW- ID:P32247 BOMBESIN RECEPTOR SUBTYPE-3 (BRS-3) - HOMO SAPIENS (HUMAN), 399 aa.lpcis:SWISSPROT- ID:P32247 BOMBESIN RECEPTOR SUBTYPE-3 (BRS-3) - HOMO SAPIENS (HUMAN), 399 aa.lpcis:TREMBLNEW- ID:E1240254 BOMBESIN RECEPTOR SUBTYPE-3 (UTERINE BOMBESIN RECEPTOR, BRS-3) - HOMO SAPIENS (HUMAN), 399 aa.	3.00E-212	X

125	cg3003519	711	CTTATGCTGTGATC ATTTCAGTGGG[CT JATCCTTGGAAATG CTATTCTCATCA	C	T	Gly	Gly	SILENT- CODING	tm7	Human Gene SWISSNEW- ID:P32247 BOMBESIN RECEPTOR SUBTYPE-3 (BRS-3) - HOMO SAPIENS (HUMAN), 399 aa, pcIs:SWISSPROT- ID:P32247 BOMBESIN RECEPTOR SUBTYPE-3 (BRS-3) - HOMO SAPIENS (HUMAN), 399 aa, pcIs:TREMBLNEW- ID:E1240254 BOMBESIN RECEPTOR SUBTYPE-3 (UTERINE BOMBESIN RECEPTOR, BRS-3) - HOMO SAPIENS (HUMAN), 399 aa.	3.00E-212	X
126	cg43969010	1182	TCCGAAAGAAAGTC TTGGGAGGTGTA[CT TTCAGGGAGTGTG CCAGAAAGGGGGC T	C	T	Tyr	Tyr	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P30411 B2 BRADYKININ RECEPTOR (BK-2 RECEPTOR) - HOMO SAPIENS (HUMAN), 391 aa.	9.00E-211	12 (14q32.1)
127	cg43263108	1097	AGACACCCCTTTCC CAGCTCGCCTC[CA JGGGAGGAGGGAC CCAAGGGCCCCCT	C	A	Ser	Ser	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P43119 PROSTACYCLIN RECEPTOR (PROSTANOID IP RECEPTOR) (PGI RECEPTOR) - HOMO SAPIENS (HUMAN), 386 aa.	8.30E-208	19 (19q13.3)
128	cg43263108	272	GCCCTCGGCCTTC GCGGTGCTGGT[CT GJACCGGACTGGCG GCCACCGACCTGC	C	G	Val	Val	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P43119 PROSTACYCLIN RECEPTOR (PROSTANOID IP RECEPTOR) (PGI RECEPTOR) - HOMO SAPIENS (HUMAN), 386 aa.	8.30E-208	19 (19q13.3)

129	cg43267238	1220	CCAGACTGGTCCT GGTGGTGGTGGCJA /GJGTCITTCGTCGTC TGCTGGACTCCCA	A	G	Ala	Ala	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P41145 KAPPA-TYPE OPIOID RECEPTOR (KOR-1) - HOMO SAPIENS (HUMAN), 380 aa.	2.10E-204	8 (8q11.2)
130	cg43267238	392	CAGCACTCACCAT GGAATCCCCGATJC /TJCAGATCTCCGC GGGAGCGGGGCC	C	T	Ile	Ile	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P41145 KAPPA-TYPE OPIOID RECEPTOR (KOR-1) - HOMO SAPIENS (HUMAN), 380 aa.	2.10E-204	8 (8q11.2)
131	cg43267238	413	CGATCCAGATCTTC CGCGGGAGCC[G/ TJGGCCTACCTGC GCCCCGAGCGCCT	G	T	Pro	Pro	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P41145 KAPPA-TYPE OPIOID RECEPTOR (KOR-1) - HOMO SAPIENS (HUMAN), 380 aa.	2.10E-204	8 (8q11.2)
132	cg43264978	155	TGGATCTGCACCTC TTCGACTACTC[A/C JGAGCCAGGGAAC TTCTCGGACATCA	A	C	Ser	Ser	SILENT- CODING	tm7	Human Gene TREMBLNEW- ID:G2736282 G PROTEIN COUPLED RECEPTOR - HOMO SAPIENS (HUMAN), 362 aa.	1.40E-196	
133	cg3001696	1154	CGCTGCACCTGTG CATCGCGCTGGGJC /TTACGCCAATAG CAGCCTCAACCCC G	C	T	Gly	Gly	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P41143 DELTA-TYPE OPIOID RECEPTOR (DOR-1) - HOMO SAPIENS (HUMAN), 372 aa.	2.10E-195	1 (1p36.1)

134	cg3001696	815	TGGCTGTGACCCG TCCCCGGGACGGG /TGCAGTGGTGTG CATGCTCCAGTTC	G	T	Gly	Gly	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P41143 DELTA-TYPE OPIOID RECEPTOR (DOR-1) - HOMO SAPIENS (HUMAN), 372 aa.	2.10E-195	1 (1p36.1)
135	cg42704646	407	TGGCCTTCCCGATC ACCATGCTGCTTC/G JACTGGTTTCGTGG GCAACGCACTGG	C	G	Leu	Leu	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P43115 PROSTAGLANDIN E2 RECEPTOR, EP3 SUBTYPE (PROSTANOID EP3 RECEPTOR) (PGE RECEPTOR, EP3 SUBTYPE) - HOMO SAPIENS (HUMAN), 390 aa.	3.10E-194	1 (1p31.2)
136	cg43326635	347	GGGATGCCACCTT CTGCTTCATCGTTC/ GTTCGCTGGCGGTG GCTGATGTGGCCG	C	G	Val	Val	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:P30542 ADENOSINE A1 RECEPTOR - HOMO SAPIENS (HUMAN), 326 aa.	1.10E-173	1
137	cg3003708	358	CCATCTCCTTCTGT GGCTGTCTCACIA/ GTCAGATGTATTTC GTTTCATGTTTCG	A	G	Thr	Thr	SILENT- CODING	tm7	Human Gene TREMBLNEW- ID:E1246031 OLFACTORY RECEPTOR - HOMO SAPIENS (HUMAN), 312 aa.	2.50E-160	
138	cg3003708	787	GGTGGAAAGCCCTT CTCCACCTGTGGIT/ CTCTCACCTGGCT GTGGTCTCTCTCT	T	C	Gly	Gly	SILENT- CODING	tm7	Human Gene TREMBLNEW- ID:E1246031 OLFACTORY RECEPTOR - HOMO SAPIENS (HUMAN), 312 aa.	2.50E-160	

139	cg3003708	841	ACAGCACCATCAT TGCTGTGTATTIT/ CJAAGCCTCTGTCC TCCCACTCAGCTG	T	C	Phe	Phe	SILENT- CODING	tm7	Human Gene TREMBLNEW- ID:E1246031 OLFACTORY RECEPTOR - HOMO SAPIENS (HUMAN), 312 aa.	2.50E-160	
140	cg36729339	537	ACTCTCAATGTAC TTTTTCCTCTC[CT] AACCTCTCCTCTT GGACCTCTGCT	C	T	Ser	Ser	SILENT- CODING	tm7	Human Gene SWISSPROT- ID:Q15062 OLFACTORY RECEPTOR-LIKE PROTEIN FAT11 - HOMO SAPIENS (HUMAN), 316 aa.	1.90E-153	
141	cg38841806	717	GACATCAGGCGCA CGGTGCCAACTTC /GJGCGCAATCTGCA GGCCAAAGAAAG T	C	G	Leu	Leu	SILENT- CODING	tm7	Human Gene Similar to SWISSPROT-ID:P30975 TACHYKININ-LIKE PEPTIDES RECEPTOR 99D (DTKR) - DROSOPHILA MELANOGASTER (FRUIT FLY), 519 aa.	2.10E-67	
142	cg38841806	723	AGGCGCACGGTGC CAACCTCCGCCAT/ CJCTGCAGGCCAAG AAGAAAGTTTGTGA	T	C	His	His	SILENT- CODING	tm7	Human Gene Similar to SWISSPROT-ID:P30975 TACHYKININ-LIKE PEPTIDES RECEPTOR 99D (DTKR) - DROSOPHILA MELANOGASTER (FRUIT FLY), 519 aa.	2.10E-67	
143	cg38841806	96	CAGCCTTCTCCATG CCCAGCTGGCAJG/ AJCTGGCACTGTGG GCACCAGCCTACC	G	A	Gln	Gln	SILENT- CODING	tm7	Human Gene Similar to SWISSPROT-ID:P30975 TACHYKININ-LIKE PEPTIDES RECEPTOR 99D (DTKR) - DROSOPHILA MELANOGASTER (FRUIT FLY), 519 aa.	2.10E-67	

144	cg43040273	1966	CCTGTGCTGATCTG GTCATGGGCCT[G/ AJGCAGTGTGCCCC TTTGGGGCCGCC	G	A	Leu	Leu	SILENT- CODING	tm7	Human Gene Similar to SWISSPROT-ID:Q24563 DOPAMINE RECEPTOR 2 - DROSOPHILA MELANOGASTER (FRUIT FLY), 539 aa.	2.00E-58	5 (5q32)
145	cg43040273	2237	CTTGCCCATTCAGA TGCACTGGTAC[C/ AJGGCCACCCACC AGGAAGCCATCAA	C	A	Arg	Arg	SILENT- CODING	tm7	Human Gene Similar to SWISSPROT-ID:Q24563 DOPAMINE RECEPTOR 2 - DROSOPHILA MELANOGASTER (FRUIT FLY), 539 aa.	2.00E-58	5 (5q32)
146	cg43336100	687	TTGGAAGCGTGCA TCCAGTGAGACC[A /TATGAGGCTTGA GTCITTTAGTGCCT	A	T	Pro	Pro	SILENT- CODING	tnf	Human Gene SWISSPROT- ID:P26022 PENTAXIN- RELATED PROTEIN PTX3 PRECURSOR (TUMOR NECROSIS FACTOR- INDUCIBLE PROTEIN TSG-14) - HOMO SAPIENS (HUMAN), 381 aa.	2.20E-207	3 (3q25)
147	cg21646034	376	GTGTGAGCAGAGA TGCCAGAACCA[A /GJTGACCCGAAC ACCATACATATG G	A	G	Lys	Lys	SILENT- CODING	transcriptfactor	Human Gene SWISSPROT- ID:Q06545 GA BINDING PROTEIN BETA-2 CHAIN (GABP-BETA-2 SUBUNIT) (TRANSCRIPTION FACTOR E4TF1-47) (GAPBP2) - HOMO SAPIENS (HUMAN), 347 aa.	9.00E-179	15

148	cg43916882	1608	TGGCAGCTACCAG CACACTGCCTCC[A/ G]CCGTCAATAAAG GCACTGATGGTCT	A	G	Gly	Gly	SILENT- CODING	transferase	Human Gene SWISSPROT- ID:P39656 DOLICHYL- DIPHOSPHOOLIGOSACCHARI DE-PROTEIN GLYCOSYLTRANSFERASE 48 KD SUBUNIT PRECURSOR (EC 2.4.1.119) (OLIGOSACCHARYL TRANSFERASE 48 KD SUBUNIT) (DDOST 48 KD SUBUNIT) (KIAA0115) (HAA0643) - HOMO SAPIENS (HUMAN), 456 aa.	5.30E-245	1
149	cg2537639	294	TGGCTCCCATGTGTC TGGGAGGGCAC[A/ G]TTCAACATCGAC ATCCTCAACGAGC	A	G	Thr	Thr	SILENT- CODING	transferase	Human Gene SWISSPROT- ID:P16442 FUCOSYLGLYCOPROTEIN ALPHA-N- ACETYL GALACTOSAMINYLT RANSFERASE (EC 2.4.1.40) (HISTO-BLOOD GROUP A TRANSFERASE) (A TRANSFERASE) / FUCOSYLGLYCOPROTEIN 3- ALPHA- GALACTOSYLTRANSFERASE (EC 2.4.1.37) (HISTO-BLOOD GROUP B TRANSFERASE) (B TRANSFERASE) (NAGAT) - HOMO SAPIENS (HUMAN), 354 aa.	6.50E-192	9 (9q34)

150	cg2537639	654	ACGTGGACATGGA GTTCCGGGACCAIC TJGTGGCGGTGGA GATCCTGACTCCG C	T	His	His	SILENT- CODING	transferase	Human Gene SWISSPROT- ID:P16442 FUCOSYLGLYCOPROTEIN ALPHA-N- ACETYL GALACTOSAMINYLT RANSFERASE (EC 2.4.1.40) (HISTO-BLOOD GROUP A TRANSFERASE) (A TRANSFERASE) / FUCOSYLGLYCOPROTEIN 3- ALPHA- GALACTOSYLTRANSFERASE (EC 2.4.1.37) (HISTO-BLOOD GROUP B TRANSFERASE) (B TRANSFERASE) (NAGAT) - HOMO SAPIENS (HUMAN), 354 aa.	6.50E-192	9 (9q34)
-----	-----------	-----	--	---	-----	-----	-------------------	-------------	--	-----------	----------

151	cg2537639	678	ACGTGGGCGGTGGA GATCCTGACTCC[G/ AJCTGTTGGCACC CTGCACCCCGGCT	G	A	Pro	Pro	SILENT- CODING	transferase	Human Gene SWISSPROT- ID:P16442 FUCOSYLGLYCOPROTEIN ALPHA-N- ACETYL GALACTOSAMINYLT RANSFERASE (EC 2.4.1.40) (HISTO-BLOOD GROUP A TRANSFERASE) (A TRANSFERASE) / FUCOSYLGLYCOPROTEIN 3- ALPHA- GALACTOSYLTRANSFERASE (EC 2.4.1.37) (HISTO-BLOOD GROUP B TRANSFERASE) (B TRANSFERASE) (NAGAT) - HOMO SAPIENS (HUMAN), 354 aa.	6.50E-192	9 (9q34)
-----	-----------	-----	---	---	---	-----	-----	-------------------	-------------	--	-----------	----------

152	cg2537639	768	GGCCCCAGTCCCA GGCCTACATCCG/C/ TJAAGGACGAGGG CGATTCTACTACC	C	T	Pro	Pro	SILENT- CODING	transferase	Human Gene SWISSPROT- ID: P16442 FUCOSYLGLYCOPROTEIN ALPHA-N- ACETYL GALACTOSAMINYLT RANSFERASE (EC 2.4.1.40) (HISTO-BLOOD GROUP A TRANSFERASE) (A TRANSFERASE) / FUCOSYLGLYCOPROTEIN 3- ALPHA- GALACTOSYLTRANSFERASE (EC 2.4.1.37) (HISTO-BLOOD GROUP B TRANSFERASE) (B TRANSFERASE) (NAGAT) - HOMO SAPIENS (HUMAN), 354 aa	6.50E-192	9 (9q34)
-----	-----------	-----	--	---	---	-----	-----	-------------------	-------------	--	-----------	----------

153	cg2537639	927	ACGAGAGCCACCT GAACAAGTACCTG /AJCTGCGCCACAA ACCCACCAAGGTG C	G	A	Leu	Leu	SILENT- CODING	transferase	Human Gene SWISSPROT- ID:P16442 FUCOSYLGLYCOPROTEIN ALPHA-N- ACETYLGLALACTOSAMINYLT RANSFERASE (EC 2.4.1.40) (HISTO-BLOOD GROUP A TRANSFERASE) (A TRANSFERASE) / FUCOSYLGLYCOPROTEIN 3- ALPHA- GALACTOSYLTRANSFERASE (EC 2.4.1.37) (HISTO-BLOOD GROUP B TRANSFERASE) (B TRANSFERASE) (NAGAT) - HOMO SAPIENS (HUMAN), 354 aa.	6.50E-192	9 (9q34)
154	cg44000740	732	GGGAGAGATACTGG CTCAGGAGGAAIA /CJACAGGGAACAT CACCTTATGCCAC A	A	C	Val	Val	SILENT- CODING	transferase	Human Gene Homologous to SWISSPROT-ID:P30711 GLUTATHIONE S- TRANSFERASE THETA 1 (EC 2.5.1.18) (CLASS-THETA) - HOMO SAPIENS (HUMAN), 239 aa.	1.60E-117	16

155	cg38869466	1185	ACGCAGTGGCCGT GGGCTCCCTCTG/C/ TJGCTCTTCCGCC AGTCTTCTAGGTT	C	T	Cys	Cys	SILENT- CODING	transport	Human Gene SWISSPROT- ID:P30825 HIGH-AFFINITY CATIONIC AMINO ACID TRANSPORTER-1 (CAT-1) (CAT1) (SYSTEM Y+ BASIC AMINO ACID TRANSPORTER) (ECOTROPIC RETROVIRAL LEUKEMIA RECEPTOR HOMOLOG) (ERR) (ECOTROPIC RETROVIRUS RECEPTOR HOMOLOG) - HOMO SAPIENS (HUMAN), 629 aa.	0	13
156	cg40351913	1347	CCATCGCCACGCT CCCTCTGTCTCTC/A/ G/GCCTGGGCCGTG GTCTTCTTCATCA	A	G	Ser	Ser	SILENT- CODING	transport	Human Gene SWISSPROT- ID:Q01959 SODIUM- DEPENDENT DOPAMINE TRANSPORTER (DA TRANSPORTER) (DAT) - HOMO SAPIENS (HUMAN), 620 aa.	0	5 (5p15.3)
157	cg43964039	1719	GATGGAACAGCTC CTCGGGTGTCTT/G/ A/TCACCTTTGGCTG GCTCCCCCCTGCC	G	A	Asp	Asp	SILENT- CODING	transport	Human Gene SWISSPROT- ID:P11166 GLUCOSE TRANSPORTER TYPE 1, ERYTHROCYTE/BRAIN - HOMO SAPIENS (HUMAN), 492 aa.	1.60E-259	1
158	cg43992017	1656	GCGGCTGCTGGTG GATGGGTGGCG/C/ /GJGGGTGCAGCC TCCACCCCTCCTCCC	C	G	Pro	Pro	SILENT- CODING	transport	Human Gene SPTREMBL- ID:Q14728 TETRACYCLINE TRANSPORTER-LIKE PROTEIN MRNA - HOMO SAPIENS (HUMAN), 455 aa.	4.40E-241	

159	cg43948629	1238	CGCTGTAATGGC TGTGAACATGCTC/ TJACCCAGCAGGAG GTCCCTGTCGTTA	C	T	Leu	Leu	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSNEW- ACC:Q15031 PROBABLE LEUCYL-TRNA SYNTHETASE, MITOCHONDRIAL PRECURSOR (EC 6.1.1.4) (LEUCINE-TRNA LIGASE) (LEURS) (K1AA0028) - Homo sapiens (Human), 903 aa.	0	3
160	cg43955093	2875	CATTGACTAGGGG CTGTGGGGGCATC /GJCGCCAGGTGT CCCTCCATCAGAG G	C	G	Arg	Arg	SILENT- CODING	UNCLASSIFIED	Human Gene SPTREMBL- ACC:Q16084 P130 - HOMO SAPIENS (HUMAN), 1139 aa.	0	16
161	cg43955093	3385	AGCAGGCCAAGAG AGATCTGTGGAATC /TJGCATCTTGTCC AGAATACCAGATA	C	T	Ala	Ala	SILENT- CODING	UNCLASSIFIED	Human Gene SPTREMBL- ACC:Q16084 P130 - HOMO SAPIENS (HUMAN), 1139 aa.	0	16
162	cg43035918	517	CGCTGGCATAGGA CATGGCGGGCTTG /TJCCCCCGCAGA GCTCTGGGGGCTA C	G	T	Gly	Gly	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P42694 HYPOTHETICAL PROTEIN KIAA0054 - Homo sapiens (Human), 1942 aa.	0	17
163	cg43974592	254	AAATAACAAGGCA TTGAAGAATGGCT /AJGACGAGCGGAA AGACGAAGGAAG G	T	A	Ala	Ala	SILENT- CODING	UNCLASSIFIED	Human Gene REMTREMBL- ACC:E1296438 SEQUENCE 28 FROM PATENT WO9727323 - UNIDENTIFIED, 1829 aa.	0	2 (2q34)

164	cg43956384	206	AAGACGCAACGC TGCCACCATGGA/C /TAGTAGCACCTG GAGCCCCAAGACC A	C	T	Asp	Asp	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P13866 SODIUM/GLUCOSE COTRANSPORTER 1 (NA(+)/GLUCOSE COTRANSPORTER 1) (HIGH AFFINITY SODIUM-GLUCOSE COTRANSPORTER) - Homo sapiens (Human), 664 aa.	0	22 (22q13.1)
165	cg44025634	2757	TGAAAGTATTCAA TCCAGAGGAAGAAJ A/GJCTGGAATTG CCCTTCTGTCTTA G	A	G	Lys	Lys	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P00450 CERULOPLASMIN PRECURSOR (EC 1.16.3.1) (FERROXIDASE) - Homo sapiens (Human), 1065 aa.	0	3 (3q21)
166	cg43940037	2472	GCTGGCGCACTGC TAGCCTCAGAGGTT /AJGCCAGCACCTC CTCAGCCCCCGCG C	T	A	Ala	Ala	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P41250 GLYCYL-TRNA SYNTHETASE (EC 6.1.1.14) (GLYCINE--TRNA LIGASE) (GLYRS) - Homo sapiens (Human), 685 aa.	0	7 (7p15)
167	cg44024279	481	AAAACCAGCTACC TGCCCTTCTGGAJA/ GJGAACCTTGCCAT GAGAAAGAAATTT	A	G	Glu	Glu	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P02771 ALPHA- FETOPROTEIN PRECURSOR (ALPHA-FETOGLOBULIN) (ALPHA-1- FETOPROTEIN) - Homo sapiens (Human), 609 aa.	0	

168	cg43926814	1122	CATGAGTTTGTATC CCAGCTCTTCTCT TCCCTGGCTTCT GGGCCATTCTC	C	T	Glu	Glu	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSNEW- ACC:Q13573 NUCLEAR PROTEIN SKIP (SNW1 PROTEIN) (NUCLEAR RECEPTOR COACTIVATOR NCOA-62) - Homo sapiens (Human), 536 aa.	5.00E-289	14
169	cg40918088	1778	TGGAGCTGGAAT TACTGTGTATGAJA/ GJGCCCTTAGCAGCT GCTGATGAGCTTT	A	G	Glu	Glu	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P51854 TRANSKETOLASE 2 (EC 2.2.1.1) (TK 2) (TRANSKETOLASE RELATED PROTEIN) - Homo sapiens (Human), 557 aa.	1.80E-287	X (Xq28)
170	cg43966985	1242	TCAACACCTACGT CCACTTCCAGGJG TJAAGATGAAGGG CTTCTCCCTGCTGG	G	T	Gly	Gly	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P01019 ANGIOTENSINOGEN PRECURSOR - Homo sapiens (Human), 485 aa.	3.90E-257	1 (1q42)
171	cg43924009	770	TGGCTTGACAAA TTGCTTGAAGACIA TTCGATCCATGTAA GTGGACTGTCTTG	A	T	Arg	Arg	SILENT- CODING	UNCLASSIFIED	Human Gene SPTREMBL- ACC:O43411 HYPOTHETICAL 49.3 KD PROTEIN - HOMO SAPIENS (HUMAN), 442 aa (fragment).	6.90E-239	
172	cg42913861	2186	CTGGGCAGCTGCC CTCACAGTAGTTTC/ GJCCGTAGTAGCCG GTGGGTGCTATGA	C	G	Gly	Gly	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P09529 INHIBIN BETA B CHAIN PRECURSOR (ACTIVIN BETA-B CHAIN) - Homo sapiens (Human), 407 aa.	3.00E-227	2 (2cen)

173	cg42913861	2354	GCCGAGCCTGCAC CACACACAAAGG[C C/TTCGGTGCAGCTC TTCCCTGGGTCCA	C	T	Arg	Arg	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P09529 INHIBIN BETA B CHAIN PRECURSOR (ACTIVIN BETA-B CHAIN) - Homo sapiens (Human), 407 aa.	3.00E-227	2 (2cen)
174	cg43929685	256	CATAGAAGGCCAG GAGTCAGGAGACI C/TJGGGTTCTGTC CTGGATTATACAC C	C	T	Gln	Gln	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P29080 (2'- 5')OLIGOADENYLATE SYNTHETASE 1B (EC 2.7.7.-) ((2-5')OLIGO(A) SYNTHETASE 1B) (2-5A SYNTHETASE 1B) - Mus musculus (Mouse), 414 aa.	2.40E-225	12
175	cg43929685	268	GGAGTCAGGAGAC CTGGGTTCTGTC[C/ TTGGATTATACAC CAGCTCACTGAGG	C	T	Gln	Gln	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P29080 (2'- 5')OLIGOADENYLATE SYNTHETASE 1B (EC 2.7.7.-) ((2-5')OLIGO(A) SYNTHETASE 1B) (2-5A SYNTHETASE 1B) - Mus musculus (Mouse), 414 aa.	2.40E-225	12
176	cg43918561	53	CCATGCCACCC CGACGCCACACAC[G /CJCCACAGGCCAA GGGCTTCCGCAGG G	G	C	Thr	Thr	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P04177 TYROSINE 3- MONOOXYGENASE (EC 1.14.16.2) (TYROSINE 3- HYDROXYLASE) (TH) - Rattus norvegicus (Rat), 498 aa.	2.10E-224	11 (1p15.5)

177	cg42343176	1885	ATTTAATGAATTTC CTGAAGACTGT(A/ GJAGAAGTACAACT GAGAAATCCCTTT	A	G	Val	Val	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P14902 INDOLEAMINE 2,3-DIOXYGENASE (EC 1.13.11.42) (IDO) (INDOLEAMINE-PYRROLE 2,3- DIOXYGENASE) - Homo sapiens (Human), 403 aa.	3.90E-218	8 (8p12)
178	cg43956382	1146	AAAACAATGATAT CGATGAAGTTATC /TATCCACAGCT CCCTTATACAAAC	C	T	Ile	Ile	SILENT- CODING	UNCLASSIFIED	Human Gene SPTREMBL- ACC:Q99816 TUMOR SUSCEPTIBILITY PROTEIN - HOMO SAPIENS (HUMAN), 390 aa.	4.90E-211	11
179	cg43984681	979	CACCATGAAGCAG TTGCTGCGGGCC(C/ TTGGAGGAGGGCC GCGTGCGGGAAGT	C	T	Leu	Leu	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:O15382 BRANCHED- CHAIN AMINO ACID AMINOTRANSFERASE, MITOCHONDRIAL PRECURSOR (EC 2.6.1.42) (BCAT(M)) - Homo sapiens (Human), 392 aa.	1.30E-210	19 (19q13)
180	cg43984681	1074	TCCTGTACAAAGA CAGGAACCTCCAIC /TATCCCAACCATG GAAAATGGGCCTG	C	T	His	His	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:O15382 BRANCHED- CHAIN AMINO ACID AMINOTRANSFERASE, MITOCHONDRIAL PRECURSOR (EC 2.6.1.42) (BCAT(M)) - Homo sapiens (Human), 392 aa.	1.30E-210	19 (19q13)

181	cg43950996	1762	CTGCGGTGGAGAC GTCAGAGCTGCC[A /GJGGGAGGGGC TCCTGCGCCACAG C	A	G	Pro	SILENT- CODING	UNCLASSIFIE D	Human Gene SPTREMBL- ACC:P78545 ESE-1B - HOMO SAPIENS (HUMAN), 371 aa.	6.20E-204	1
182	cg44024506	988	ACCAGCTGCTCGT AGTACACAGGCAI G/AJGCACTTCTCT TGCTACCTCCATG	G	A	Leu	SILENT- CODING	UNCLASSIFIE D	Human Gene SPTREMBL- ACC:O60704 TYROSYLPROTEIN SULFOTRANSFERASE-2 - HOMO SAPIENS (HUMAN), 377 aa.	1.90E-200	22
183	cg43980381	1114	CTACCGCCTAACTA TGACTTTGTCT[C/ GJAAGAAGCGGAC CTTCACCAAGGGA G	C	G	Leu	SILENT- CODING	UNCLASSIFIE D	Human Gene SWISSNEW- ACC:Q03385 GUANTINE NUCLEOTIDE DISSOCIATION STIMULATOR RALGDS FORM A (RALGEF) - Mus musculus (Mouse), 852 aa.	5.60E-191	9

184	cg42650960	501	TCCCTGGCAGAA CTACCACTGAAIC /TJGACTGGATGGA GGAGGAATACCCG C	C	T	Asn	Asn	SILENT- CODING	UNCLASSIFIED D	Human Gene SWISSPROT- ACC:Q10981 GALACTOSIDE 2- L-FUCOSYLTRANSFERASE 2 (EC 2.4.1.69) (GDP-L- FUPOSE:BETA- D- GALACTOSIDE 2-ALPHA-L- FUCOSYLTRANSFERASE 2) (ALPHA(1,2)FT 2) (FUCOSYLTRANSFERASE 2) (SECRETOR BLOOD GROUP ALPHA-2- FUCOSYLTRANSFERASE) (SECRETOR FACTOR) (SE) (SE2) - Homo sapiens (Human), 343 aa.	2.00E-189	
185	cg43249389	1497	ACATCCAGGTGGT GTTCCGACGCCGTIC/ TJACCGACATCATC ATTGCCAACAAACC	C	T	Val	Val	SILENT- CODING	UNCLASSIFIED D	Human Gene SWISSPROT- ACC:P09471 GUANTINE NUCLEOTIDE-BINDING PROTEIN Q(O), ALPHA SUBUNIT 1 - Homo sapiens (Human), 353 aa.	1.40E-188	15
186	cg43946951	615	CAGTGACGGCAGG GTCAAAGTCCTTIG/ AJCGTAGCCCTCG TTAAGGCTGTAGA	G	A	Ala	Ala	SILENT- CODING	UNCLASSIFIED D	Human Gene SWISSPROT- ACC:P09467 FRUCTOSE-1,6- BISPHOSPHATASE (EC 3.1.3.11) (D-FRUCTOSE-1,6- BISPHOSPHATE 1- PHOSPHOHYDROLASE) (FBPASE) - Homo sapiens (Human), 337 aa.	3.50E-178	9 (9q22.2)

187	cg43248117	1054	AACCAGCCACTG TGAGAAAGACACI G/CJGTGTTCAAAGTC TTTGGGAATGGCA G	G	C	Thr	Thr	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:Q14894 MU-CRYSTALLIN HOMOLOG (NADP- REGULATED THYROID- HORMONE BINDING PROTEIN) - Homo sapiens (Human), 314 aa.	1.20E-161	16 (16p13.1 1)
188	cg44027049	482	CCACAATGTTAGG AGGGTATTTTATC/ TJATCCCTCCAGTT AACAAATACAGCA	C	T	Tyr	Tyr	SILENT- CODING	UNCLASSIFIED	Human Gene SWISSNEW- ACC:P11245 ARYLAMINE N- ACETYLTRANSFERASE, POLYMORPHIC (BC 2.3.1.5) (PNAT) (NAT-2) (ARYLAMINE ACETYLASE) - Homo sapiens (Human), 290 aa.	5.40E-157	8 (8p23.1)
189	cg43982075	499	CTGCCATCTTTCAG CCCTCTGAAACI/T JGTGTCCAGCACAG AATCTTCCCTGG	C	T	Thr	Thr	SILENT- CODING	UNCLASSIFIED	Human Gene SPTREMBL- ACC:Q15729 THYROTROPH EMBRYONIC FACTOR - HOMO SAPIENS (HUMAN), 303 aa.	1.20E-154	22
190	cg43942977	350	GCGCTTCCAGGT COGACAAATTCGIG /TJCAGACTATTGTC AAACTGGGGAATA	G	T	Arg	Arg	SILENT- CODING	UNCLASSIFIED	Human Gene Homologous to SWISSNEW-ACC:Q12846 SYNTAXIN 4 - Homo sapiens (Human), 297 aa.	9.60E-148	
191	cg43942977	701	GGCAGCTGAAGAT CACCAATGCTGGIG /CIATGGTGTCTGAT GAGGAGTTGGAGC	G	C	Gly	Gly	SILENT- CODING	UNCLASSIFIED	Human Gene Homologous to SWISSNEW-ACC:Q12846 SYNTAXIN 4 - Homo sapiens (Human), 297 aa.	9.60E-148	

192	cg43942977	773	GCGAGGTGTTTGT GTCCAATATCCTG/ TAAAGGACACGCA GGTGACTCGACAG G	G	T	Leu	Leu	SILENT- CODING	UNCLASSIFIED	Human Gene Homologous to SWISSNEW-ACC:Q12846 SYNTAXIN 4 - Homo sapiens (Human), 297 aa.	9.60E-148	
193	cg43985220	753	TGTACACTGCCAG AAAAAGGAAAAAGG/ T/GGCCCTTTGTAA TGGTCAAAAACTA C	T	G	Gly	Gly	SILENT- CODING	UNCLASSIFIED	Human Gene Homologous to SWISSNEW-ACC:P29218 MYO- INOSITOL-1(OR 4)- MONOPHOSPHATASE (EC 3.1.3.25) (IMP) (INOSITOL MONOPHOSPHATASE) (LITHIUM-SENSITIVE MYO- INOSITOL MONOPHOSPHATASE A1) - Homo sapiens (Human), 277 aa.	5.10E-145	8
194	cg43985220	837	TCTTGGTGACTGA GTTGGGCTCTTC[C/ TJAGAACACACAGA GACTGTGAGAATG G	C	T	Ser	Ser	SILENT- CODING	UNCLASSIFIED	Human Gene Homologous to SWISSNEW-ACC:P29218 MYO- INOSITOL-1(OR 4)- MONOPHOSPHATASE (EC 3.1.3.25) (IMP) (INOSITOL MONOPHOSPHATASE) (LITHIUM-SENSITIVE MYO- INOSITOL MONOPHOSPHATASE A1) - Homo sapiens (Human), 277 aa.	5.10E-145	8
195	cg43946394	321	TAGAGCGCACACA GGCCTCCAGCTG/A /GJCCCATGTCCGTC TCATCATCCCAAG	A	G	Ala	Ala	SILENT- CODING	UNCLASSIFIED	Human Gene Homologous to SWISSPROT-ACC:P29692 ELONGATION FACTOR 1- DELTA (EF-1-DELTA) - Homo sapiens (Human), 281 aa.	2.80E-144	19

196	cg43119818	1329	CTGACAGCTACAG GCTCTTTCAGTTTC/ TJCATTTTCAGTGG GGCAGTACAAATG	C	T	Phe	Phe	SILENT- CODING	UNCLASSIFIED	Human Gene Homologous to SWISSPROT-ACC:P00915 CARBONIC ANHYDRASE I (EC 4.2.1.1) (CARBONATE DEHYDRATASE I) - Homo sapiens (Human), 260 aa.	6.90E-141	8 (8q22)
197	cg43118279	735	AGAAAGTTGAAGGG GCTGGTGCCACTTT/ GJGGACCCGAATCA AGTCGACACACTA	T	G	Leu	Leu	SILENT- CODING	UNCLASSIFIED	Human Gene Homologous to SWISSPROT-ACC:Q05195 MAD PROTEIN (MAX DIMERIZER) - Homo sapiens (Human), 221 aa.	1.20E-111	2 (2p13)
198	cg43325007	866	TGGGTTTCAGGGAT GTAGCCCTTCTCTT/ CJACAGCCAGGCG GCTCAGGGCAAAC A	T	C	Val	Val	SILENT- CODING	UNCLASSIFIED	Human Gene Homologous to TREMBLNEW-ACC:AAD43195 PEROXISOMAL MEMBRANE PROTEIN PMP 24 - HOMO SAPIENS (HUMAN), 212 aa.	4.80E-110	20
199	cg39524111	402	GCCAAATATAGGAT AGGGCACTACAGI A/GJTTCCGGTACA GTGACACCCCTGGA GC	A	G	Arg	Arg	SILENT- CODING	UNCLASSIFIED	Human Gene Similar to TREMBLNEW-ACC:BAA13472 CD89 U08 - HOMO SAPIENS (HUMAN), 191 aa.	2.10E-100	19 (19q13.4)
200	cg43280516	629	ACGGGAGGAGCT GCAGATGGAACCTC /TJGTGTGAGGTGTC TTCTGGGACCTGC	C	T	Pro	Pro	SILENT- CODING	UNCLASSIFIED	Human Gene Similar to TREMBLNEW-ACC:CAB43107 PRENYLATED RAB ACCEPTOR 1 (PRA1) - HOMO SAPIENS (HUMAN), 185 aa.	6.80E-95	19

201	cg43963913	871	AGAGGTTGGGGG CGCCGAGCGCAI G/AJCGGCCCGAA AGGGGCTGGGCTC CT	G	A	Arg	SILENT- CODING	UNCLASSIFIED	Human Gene Similar to SPTREMBL-ACC:O14803 BCL- X/BCL-2 BINDING PROTEIN - HOMO SAPIENS (HUMAN), 168 aa (fragment).	5.10E-90	11
202	cg40262905	682	TAGTGAAGGCCT GAAATATATGCTG /CJGAGGTGGAAT TGGCAGAACTACC T	G	C	Leu	SILENT- CODING	UNCLASSIFIED	Human Gene Similar to TREMBLNEW-ACC:BAA34941 HUMAN CMAP - HOMO SAPIENS (HUMAN), 167 aa.	1.30E-89	
203	cg43918168	915	CTCCATCAACAGC ATCCGGACTGCAIT /CJGGCGGCTCGCC GTGCGGCTGGGGC C	T	C	Pro	SILENT- CODING	UNCLASSIFIED	Human Gene Similar to SWISSPROT-ACC:P09496 CLATHRIN LIGHT CHAIN A (BRAIN AND LYMPHOCYTE LCA) - Homo sapiens (Human), 248 aa.	3.80E-85	9 (12q23)
204	cg43259701	136	CGACGAGGTGCTA CGCGAGGGCGAGI C/TJTGAGAGAGCG CAGCGACAGCCTC TT	C	T	Leu	SILENT- CODING	UNCLASSIFIED	Human Gene Similar to SPTREMBL-ACC:O00496 IPL (IPL) - HOMO SAPIENS (HUMAN), 152 aa.	1.30E-77	11
205	cg1527767	162	TTTTTCCAGCTTA CAATGGTACAGIA/ GICAGGAGCCTGG GGAAGGTCCTGTC C	A	G	Arg	SILENT- CODING	UNCLASSIFIED	Human Gene Similar to REMTREMBL-ACC:G36907 T- CELL RECEPTOR ALPHA- CHAIN HAP58 V(A)10.1-J(A)T - HOMO SAPIENS (HUMAN), 135 aa (fragment).	5.60E-68	

206	cg40968986	316	AGAAGAGAGCCTG TGACACTGCCACIC /TTGTGTGACTCAT CGGCTGGCAGGCT	C	T	Thr	Thr	SILENT- CODING	UNCLASSIFIED	Human Gene Similar to SWISSNEW-ACC:P06881 CALCITONIN GENE-RELATED PEPTIDE 1 PRECURSOR (CGRP- D) (ALPHA-TYPE CGRP) - Homo sapiens (Human), 128 aa.	5.10E-58	11 (11p15.2)
207	cg42550133	300	TCATCTGAGTTCT AAGAAGCTCCTT/ CCTCAGTGACTCT GGCTTCTATCTCT	T	C	Leu	Leu	SILENT- CODING	UNCLASSIFIED	Human Gene Similar to REMTEMBL-ACC:D1002898 T- CELL RECEPTOR BETA-CHAIN V REGION - HOMO SAPIENS (HUMAN), 112 aa (fragment).	8.50E-56	7 (7q35)
208	cg25226759	317	CTCTGGTTGTCCAC GAGGGAGACACT/ CJGTAACCTCTCAAT TGCAGTTATGAAG	T	C	Thr	Thr	SILENT- CODING	UNCLASSIFIED	Human Gene Similar to REMTEMBL-ACC:G33509 T CELL RECEPTOR - HOMO SAPIENS (HUMAN), 118 aa (fragment).	1.60E-54	
209	cg41664708	249	AAGTCTGTGCTGA TCCACAAGCCACIA /GJTGGGTGAGAGA CGTGGTCAGGAGC A	A	G	Thr	Thr	SILENT- CODING	UNCLASSIFIED	Human Gene Similar to SWISSNEW-ACC:P47992 LYMPHOTACTIN PRECURSOR (CYTOKINE SCM-1) (ATAC) (LYMPHOTAXIN) (SCM-1- ALPHA) - Homo sapiens (Human), 114 aa.	2.00E-54	1
210	cg43300673	1571	AGGGAGGCGGGGA GGGTAGCATGGG/ G/gap]CACACGGCC CTCACAGGGACTC ACT	G	gap			SILENT- NONCODING	ATPase associated	Human Gene SPTREMBL- ID:Q93050 VACUOLAR-TYPE H(+)-ATPASE 115 KDA SUBUNIT - HOMO SAPIENS (HUMAN), 831 aa.	0	17

211	cg43284434	2570	AGTTGAAATCAGA GAGGAATAAAAAA]g ap/A]GACATTTTAT ATTTTATTCGCTC C	gap	A				SILENT- NONCOD ING	ATPase_associat ed	Human Gene Homologous to SPTREMBL-ID:Q18788 C52E4.5 - CAENORHABDITIS ELEGANS, 590 aa.	4.00E-121	6
212	cg43132502	196	TAAGCATGAGGTG GCACGAGGACGGI A/C]GTTGGCGATG CCACCTGGGGGTC AC	A	C				SILENT- NONCOD ING	ATPase_associat ed	Human Gene Similar to SPTREMBL-ID:Q15332 GAMMA SUBUNIT OF SODIUM POTASSIUM ATPASE LIKE - HOMO SAPIENS (HUMAN), 126 aa.	9.40E-58	11
213	cg43931765	606	GGTCCCTTGCTTT ATCCCAAGCTC[G/T]GAGGGACGCAGC CTGGCATGGCTCT	G	T				SILENT- NONCOD ING	cadherin	Human Gene SWISSPROT- ID:P18084 INTEGRIN BETA-5 SUBUNIT PRECURSOR - HOMO SAPIENS (HUMAN), 799 aa.	0	3
214	cg43931765	607	GTCCCTTGCTTTA TCCCAAGCTC[G/T]AGGGACGCAGCCT GGCATGGCTCTG	G	T				SILENT- NONCOD ING	cadherin	Human Gene SWISSPROT- ID:P18084 INTEGRIN BETA-5 SUBUNIT PRECURSOR - HOMO SAPIENS (HUMAN), 799 aa.	0	3
215	cg43931765	615	CTTTATCCCAAGCT CGGAGGGACGC]ga p/G]AGCCTGGCATG GCTCTGGCCTAGC A	gap	G				SILENT- NONCOD ING	cadherin	Human Gene SWISSPROT- ID:P18084 INTEGRIN BETA-5 SUBUNIT PRECURSOR - HOMO SAPIENS (HUMAN), 799 aa.	0	3

216	cg43931765	660	TAGCAGCCAGGTG ACATGGCCAGGCig ap/TJACCTTCCTGT ACAGGCACTGTGG GC	gap	T				SILENT- NONCOD ING	cadherin	Human Gene SWISSPROT- ID:P18084 INTEGRIN BETA-5 SUBUNIT PRECURSOR - HOMO SAPIENS (HUMAN), 799 aa.	0		3
217	cg43931765	665	GCCAGGTGACATG GCCAGGCACCTTig ap/TJCCTGTACAGG CACTGTGGGCTCCT G	gap	T				SILENT- NONCOD ING	cadherin	Human Gene SWISSPROT- ID:P18084 INTEGRIN BETA-5 SUBUNIT PRECURSOR - HOMO SAPIENS (HUMAN), 799 aa.	0		3
218	cg43931765	668	AGGTGACATGGCC AGGCACCTTCCTIga p/TJGTACAGGCACT GTGGGCTCCTGGC C	gap	T				SILENT- NONCOD ING	cadherin	Human Gene SWISSPROT- ID:P18084 INTEGRIN BETA-5 SUBUNIT PRECURSOR - HOMO SAPIENS (HUMAN), 799 aa.	0		3
219	cg43931765	668	AGGTGACATGGCC AGGCACCTTCCTIga p/TJGTACAGGCACT GTGGGCTCCTGGC C	gap	T				SILENT- NONCOD ING	cadherin	Human Gene SWISSPROT- ID:P18084 INTEGRIN BETA-5 SUBUNIT PRECURSOR - HOMO SAPIENS (HUMAN), 799 aa.	0		3
220	cg43931765	668	AGGTGACATGGCC AGGCACCTTCCTIga p/TJGTACAGGCACT GTGGGCTCCTGGC C	gap	T				SILENT- NONCOD ING	cadherin	Human Gene SWISSPROT- ID:P18084 INTEGRIN BETA-5 SUBUNIT PRECURSOR - HOMO SAPIENS (HUMAN), 799 aa.	0		3

221	cg43932088	4769	AATCCACAATCGG CATCAGGAAGCCIA /CJAAGTCCAGTG GCCATTAGGTCC T	A	C				SILENT- NONCOD ING	cadherin	Human Gene SPTRMBL- ID:Q15065 OB-CADHERIN-1 - HOMO SAPIENS (HUMAN), 796 aa.	0	16
222	cg44010957	1406	TCCCTATGAGCCTG CAAAGGAGACAIG/ TJTCAGGAATGAGT TCCATGTTTCGAGA	G	T				SILENT- NONCOD ING	cadherin	Human Gene SWISSPROT- ID:P20701 LEUKOCYTE ADHESION GLYCOPROTEIN LFA-1 ALPHA CHAIN PRECURSOR (LEUKOCYTE FUNCTION ASSOCIATED MOLECULE 1, ALPHA CHAIN) (CD11A) (INTEGRIN ALPHA-L) - HOMO SAPIENS (HUMAN), 1170 aa.	0	16 (16p11.2)
223	cg43956560	1463	CAGTGCATCTGGG AAGATTCTACCTI/ CTGACCAACAGTTC CTTCAGCTTCCAT	T	C				SILENT- NONCOD ING	cadherin	Human Gene SWISSPROT- ID:P14151 L-SELECTIN PRECURSOR (LYMPH NODE HOMING RECEPTOR) (LEUKOCYTE ADHESION MOLECULE-1) (LAM-1) (LEUKOCYTE SURFACE ANTIGEN LEU-8) (TQ1) (GP90- MEL) (LEUKOCYTE- ENDOTHELIAL CELL ADHESION MOLECULE 1) (LECAM1) (CD62L) - HOMO SAPIENS (HUMAN), 372 aa.	1,00E-218	1 (1q23)

224	cg43956560	1492	CAACAGTTCCTTCA GCTTCCATTTC[G/A] JCCCTCATTTATC CCTCAACCCCA	G	A			SILENT- NONCOD ING	cadherin	Human Gene SWISSPROT- ID:P14151 L-SELECTIN PRECURSOR (LYMPH NODE HOMING RECEPTOR) (LEUKOCYTE ADHESION MOLECULE-1) (LAM-1) (LEUKOCYTE SURFACE ANTIGEN LEU-8) (TQ1) (GP90- MEL) (LEUKOCYTE- ENDOTHELIAL CELL ADHESION MOLECULE 1) (LECAM1) (CD62L) - HOMO SAPIENS (HUMAN), 372 aa.	1.00E-218	1 (1q23)
225	cg43956560	2242	TGCTCTCCTTTCCC CTGCCCCAGAC[C/ A]CTTTATCCACT TACCTAGATTCTA	C	A			SILENT- NONCOD ING	cadherin	Human Gene SWISSPROT- ID:P14151 L-SELECTIN PRECURSOR (LYMPH NODE HOMING RECEPTOR) (LEUKOCYTE ADHESION MOLECULE-1) (LAM-1) (LEUKOCYTE SURFACE ANTIGEN LEU-8) (TQ1) (GP90- MEL) (LEUKOCYTE- ENDOTHELIAL CELL ADHESION MOLECULE 1) (LECAM1) (CD62L) - HOMO SAPIENS (HUMAN), 372 aa.	1.00E-218	1 (1q23)

226	cg43264626	428	TGGCCACAGTGAA AAAGGTCATGGGTT /AJGGAGAGAAGCA AAGTAGGAAGGAT C	T	A			SILENT- NONCOD ING	cathepsin	Human Gene SWISSPROT- ID:P43235 CATHEPSIN K PRECURSOR (EC 3.4.22.38) (CATHEPSIN O) (CATHEPSIN X) (CATHEPSIN O2) - HOMO SAPIENS (HUMAN), 329 aa.	4.10E-183	1
227	cg43011543	1972	ACCGCACCTTTCC ACCGGTGGGGC/ GJCCAGTGAAGTT TAACAAACTGCTG	C	G			SILENT- NONCOD ING	collagen	Human Gene SWISSPROT- ID:P27658 COLLAGEN ALPHA 1(VIII) CHAIN PRECURSOR (ENDOTHELIAL COLLAGEN) - HOMO SAPIENS (HUMAN), 744 aa.	0	
228	cg43011543	2096	CATACCACGTTC CTGCAAGGGGGG/ C/GJACGTGTGGG TTGCTCTATTCAAG A	C	G			SILENT- NONCOD ING	collagen	Human Gene SWISSPROT- ID:P27658 COLLAGEN ALPHA 1(VIII) CHAIN PRECURSOR (ENDOTHELIAL COLLAGEN) - HOMO SAPIENS (HUMAN), 744 aa.	0	
229	cg43933757	2546	GAACCCAGTAGG CTCTGGAGGCCIA /CTGGTCAGCTTGC TTGGAAATCCAGCA	A	C			SILENT- NONCOD ING	complement	Human Gene SWISSPROT- ID:P10643 COMPLEMENT COMPONENT C7 PRECURSOR - HOMO SAPIENS (HUMAN), 843 aa.	0	5 (5p13)
230	cg41553795	64	TGGTGGTGCTACC CTTGGCTCCCAIC/ GJAGTCTGCCACC CTGCTGCCGCCAC	C	G			SILENT- NONCOD ING	complement	Human Gene Homologous to SWISSPROT-ID:P07360 COMPLEMENT C8 GAMMA CHAIN PRECURSOR - HOMO SAPIENS (HUMAN), 202 aa.	1.40E-104	9 (9q34.3)

231	cg42342496	168	AGCCCTTCTCCACC CGGATAGATTCTC/CT JTCACCCCTTGCCCC GCCTTTGCCCCA	C	T			SILENT- NONCOD ING	csf	Human Gene SWISSPROT- ID:P40225 THROMBOPOIETIN PRECURSOR (MEGAKARYOCYTE COLONY STIMULATING FACTOR) (C- MPL LIGAND) (ML) (MEGAKARYOCYTE GROWTH AND DEVELOPMENT FACTOR) (MGDF) - HOMO SAPIENS (HUMAN), 353 aa.	1.20E-189	3 (3q26.3)
232	cg42342496	179	ACCCGGATAGATT CCTCACCCCTTGG/C/ TJCCGCTTTGCC CACCCCTACTCTGC	C	T			SILENT- NONCOD ING	csf	Human Gene SWISSPROT- ID:P40225 THROMBOPOIETIN PRECURSOR (MEGAKARYOCYTE COLONY STIMULATING FACTOR) (C- MPL LIGAND) (ML) (MEGAKARYOCYTE GROWTH AND DEVELOPMENT FACTOR) (MGDF) - HOMO SAPIENS (HUMAN), 353 aa.	1.20E-189	3 (3q26.3)
233	cg41533258	1356	GTGCCTGGACATTT GCCTTGCTGGA/C/CT JGGGGACTGGGA TGTGGGAGGGAGC	C	T			SILENT- NONCOD ING	csf	Human Gene Homologous to SWISSPROT-ID:P09919 GRANULOCYTE COLONY- STIMULATING FACTOR PRECURSOR (G-CSF) (PLURIPROETIN) - HOMO SAPIENS (HUMAN), 207 aa.	1.50E-107	17 (17q11.2)

234	cg2753430	657	ACGACTTTGAGCC TCGCGATCTTTT[ga p/GJAGTCCACGTC CAGCTCGTTCTCTG	gap	G			SILENT- NONCOD ING	csf	Human Gene Similar to SWISSNEW-ID:P08700 INTERLEUKIN-3 PRECURSOR (IL-3) (MULTIPOTENTIAL COLONY-STIMULATING FACTOR) (HEMATOPOIETIC GROWTH FACTOR) (P-CELL STIMULATING FACTOR) (MAST-CELL GROWTH FACTOR) (MCGF) - HOMO SAPIENS (HUMAN), 152 aa.lpcIs:SWISSPROT-ID:P08700 INTERLEUKIN-3 PRECURSOR (IL-3) (MULTIPOTENTIAL COLONY-STIMULATING FACTOR) (HEMATOPOIETIC GROWTH FACTOR) (P-CELL STIMULATING FACTOR) (MAST-CELL GROWTH FACTOR) (MCGF) - HOMO SAPIENS (HUMAN), 152 aa.	1.10E-77	5
235	cg44036323	225	TGGGGCTTAAAAG GGCAACCCCGCG[G /C]GGACCCCTTCCTC CCTAGTCGCGGGG	G	C			SILENT- NONCOD ING	dehydrogenase	Human Gene SWISSPROT- ID:P0367 GLUTAMATE DEHYDROGENASE 1 PRECURSOR (EC 1.4.1.3) (GDH) - HOMO SAPIENS (HUMAN), 558 aa.	5.80E-303	10

236	cg43918671	766	GAGAGACCATTTA CTTACATCAGTTTC/ TJGGTTATAGACA TTTGAATCATATC	C	T			SILENT- NONCOD ING	dehydrogenase	Human Gene SPTREMBL- ID:Q14131 DIHYDROLIPOAMIDE DEHYDROGENASE - HOMO SAPIENS (HUMAN), 511 aa.	5.10E-272	7 (7q31)
237	cg43057018	1995	AGTTTCATTATACT TTTCTCTCCAC[GJTTCGCTATGTT GAAAATTTTCTG	gap	G			SILENT- NONCOD ING	dehydrogenase	Human Gene SWISSNEW- ID:P08319 ALCOHOL DEHYDROGENASE CLASS II PI CHAIN (EC 1.1.1.1) - HOMO SAPIENS (HUMAN), 391 aa.[pcis:SWISSPROT-ID:P08319 ALCOHOL DEHYDROGENASE CLASS II PI CHAIN (EC 1.1.1.1) - HOMO SAPIENS (HUMAN), 391 aa.	1.30E-209	4 (4q22)
238	cg44005808	3691	ACAAGACAGAAAGC TGAAAGTGCATCCIG ap/CJAAAAGGTGCTC AGAGAGCCGGCCC GC	gap	C			SILENT- NONCOD ING	dna_rna_bind	Human Gene SWISSNEW- ID:P19838 NUCLEAR FACTOR NF-KAPPA-B P105 SUBUNIT (DNA-BINDING FACTOR KBFI) (EBP- 1) [CONTAINS: NUCLEAR FACTOR NF-KAPPA- B P50 SUBUNIT] - HOMO SAPIENS (HUMAN), 969 aa.[pcis:SWISSPROT-ID:P19838 NUCLEAR FACTOR NF-KAPPA- B P105 SUBUNIT (CONTAINS: NUCLEAR FACTOR NF- KAPPA-B P50 SUBUNIT) (DNA- BINDING FACTOR KBFI) (EBP- 1) - HOMO SAPIENS (HUMAN), 969 aa.	0	

239	cg44005808	630	TCTTCCTTCTCCAG CCGGCAGGCCCG[/GJCGCGCTTAGG AGGGAGAGCCCCAC C	gap	G				SILENT- NONCOD ING	dna_rna_bind	Human Gene SWISSNEW- ID:P19838 NUCLEAR FACTOR NF-KAPPA-B P105 SUBUNIT (DNA-BINDING FACTOR KBFI) (EBP- 1) [CONTAINS: NUCLEAR FACTOR NF-KAPPA- B P50 SUBUNIT] - HOMO SAPIENS (HUMAN), 969 aa. pcis:SWISSPROT-ID:P19838 NUCLEAR FACTOR NF-KAPPA- B P105 SUBUNIT (CONTAINS: NUCLEAR FACTOR NF- KAPPA-B P50 SUBUNIT) (DNA- BINDING FACTOR KBFI) (EBP- 1) - HOMO SAPIENS (HUMAN), 969 aa.	0		
240	cg43956159	1244	TGGCGAGTCCAGG GTCACCCACATA[ap/A]CCATGCACCA CGGGTGCTATGCC GC	gap	A				SILENT- NONCOD ING	dna_rna_bind	Human Gene SPTREMBL- ID:Q99612 DNA-BINDING PROTEIN CPBP - HOMO SAPIENS (HUMAN), 290 aa (fragment).	1.40E-159	10	
241	cg43956159	1248	GAGTCCAGGGTCA CCCACATACCAT[p/TJGCACCCACGGGT GCTATGCCGCTTCT	gap	T				SILENT- NONCOD ING	dna_rna_bind	Human Gene SPTREMBL- ID:Q99612 DNA-BINDING PROTEIN CPBP - HOMO SAPIENS (HUMAN), 290 aa (fragment).	1.40E-159	10	

242	cg43956159	1268	TACCATGCACCAC GGGTGCTATGCCG /AJCTTCTTACAGGA CCTTTTAGCCCT	G	A			SILENT- NONCOD ING	dna_rna_bind	Human Gene SPTREMBL- ID:Q99612 DNA-BINDING PROTEIN CPBP - HOMO SAPIENS (HUMAN), 290 aa (fragment).	1.40E-159	10
243	cg43956159	1342	CCTGGAGGCAACT GGGTAGGGTGCAI G/CJAACGGCATGC TTTGGCTGGAACA CG	G	C			SILENT- NONCOD ING	dna_rna_bind	Human Gene SPTREMBL- ID:Q99612 DNA-BINDING PROTEIN CPBP - HOMO SAPIENS (HUMAN), 290 aa (fragment).	1.40E-159	10
244	cg43956159	1364	CAGAACGGCATGC TTTGGCTGGAAC[ga p/CJACGCATCCCTC CTTCCACGGCCGG C	gap	C			SILENT- NONCOD ING	dna_rna_bind	Human Gene SPTREMBL- ID:Q99612 DNA-BINDING PROTEIN CPBP - HOMO SAPIENS (HUMAN), 290 aa (fragment).	1.40E-159	10
245	cg43971258	471	CAGAGCTAGCTCT GGCTCTTCAGGC/C/ TJACAAAGTTCACAG TCCTTCGGCTCCTG	C	T			SILENT- NONCOD ING	dna_rna_bind_in hib	Human Gene Similar to SWISSNEW-ID:Q02535 DNA- BINDING PROTEIN INHIBITOR ID-3 (ID-LIKE PROTEIN INHIBITOR HLH 1R21) (HELIX- LOOP-HELIX PROTEIN HEIR-1) - HOMO SAPIENS (HUMAN), 119 aa. lpcis:SWISSPROT- ID:Q02535 DNA-BINDING PROTEIN INHIBITOR ID-3 (ID- LIKE PROTEIN INHIBITOR HLH 1R21) (HELIX-LOOP- HELIX PROTEIN HEIR-1) - HOMO SAPIENS (HUMAN), 119 aa.	1.30E-60	1 (1p36.13)

246	cg43971258	508	GTCCTTCGCTCCTG AGCACCAGGTTT/ CJAGTCTCCAGGAA GGGATTGGTGAA	T	C				SILENT- NONCOD ING	dna_rna_bind_in hib	Human Gene Similar to SWISSNEW-ID:Q02535 DNA- BINDING PROTEIN INHIBITOR ID-3 (ID-LIKE PROTEIN INHIBITOR HLH 1R21) (HELIX- LOOP-HELIX PROTEIN HEIR-1) - HOMO SAPIENS (HUMAN), 119 aa.lpcis:SWISSPROT- ID:Q02535 DNA-BINDING PROTEIN INHIBITOR ID-3 (ID- LIKE PROTEIN INHIBITOR HLH 1R21) (HELIX-LOOP- HELIX PROTEIN HEIR-1) - HOMO SAPIENS (HUMAN), 119 aa.	1.30E-60	1 (1p36.13)
247	cg43982507	3373	GATACCTTTGGGTG GATCAAGCTTG[ga /CJGTACTTGACCG TTTTATATTACT	gap	C				SILENT- NONCOD ING	eph	Human Gene SWISSPROT- ID:P98155 VERY LOW- DENSITY LIPOPROTEIN RECEPTOR PRECURSOR (VLDL RECEPTOR) - HOMO SAPIENS (HUMAN), 873 aa.	0	9 (9p24)
248	cg43982507	3739	CAAAAAAATTTAT AAACTAATTTG[ga p/GJTACGTATGAAT GATATCTTTGACCT	gap	G				SILENT- NONCOD ING	eph	Human Gene SWISSPROT- ID:P98155 VERY LOW- DENSITY LIPOPROTEIN RECEPTOR PRECURSOR (VLDL RECEPTOR) - HOMO SAPIENS (HUMAN), 873 aa.	0	9 (9p24)
249	cg43982507	514	CCTCCTTCTCCCC TTTCCCTCCCA/C JGCCCCACCTTCT TCCTCCTTTCCG	A	C				SILENT- NONCOD ING	eph	Human Gene SWISSPROT- ID:P98155 VERY LOW- DENSITY LIPOPROTEIN RECEPTOR PRECURSOR (VLDL RECEPTOR) - HOMO SAPIENS (HUMAN), 873 aa.	0	9 (9p24)

250	cg41554010	1371	CTGCCCTGCCACCT GTCTGTCTGTCT[<u>gap</u>]/ TJCCTAAAGAGTTCTC TGGTATGAACTTG	gap	T			SILENT- NONCOD ING	eph	Human Gene SWISSNEW- ID:P06727 APOLOPROTEIN A- IV PRECURSOR (APO-AIV) - HOMO SAPIENS (HUMAN), 396 aa.lpcis:SWISSPROT-ID:P06727 APOLOPROTEIN A-IV PRECURSOR (APO-AIV) - HOMO SAPIENS (HUMAN), 396 aa.	1.80E-203	11 (11q23)
251	cg41554010	1371	CTGCCCTGCCACCT GTCTGTCTGTCT[<u>gap</u>]/ TJCCTAAAGAGTTCTC TGGTATGAACTTG	gap	T			SILENT- NONCOD ING	eph	Human Gene SWISSNEW- ID:P06727 APOLOPROTEIN A- IV PRECURSOR (APO-AIV) - HOMO SAPIENS (HUMAN), 396 aa.lpcis:SWISSPROT-ID:P06727 APOLOPROTEIN A-IV PRECURSOR (APO-AIV) - HOMO SAPIENS (HUMAN), 396 aa.	1.80E-203	11 (11q23)
252	cg43984905	2376	TCCCTCCAGGACT AGGCTGGAGGAAI G/CJCCAGTGGGT CCCCCTGAGTGG GC	G	C			SILENT- NONCOD ING	esterase	Human Gene SWISSPROT- ID:P51178 1- PHOSPHATIDYLINOSITOL-4,5- BISPHOSPHATE PHOSPHODIESTERASE DELTA 1 (EC 3.1.4.11) (PLC-DELTA-1) (PHOSPHOLIPASE C-DELTA-1) (PLC-III) - HOMO SAPIENS (HUMAN), 756 aa.	0	3

253	cg43984905	2440	CACATGTGGGGAC AGGGCTGGTGTG/G /CJCTGCTCCAGCC TCTTGCTCAGAGC	G	C				SILENT- NONCOD ING	esterase	Human Gene SWISSPROT- ID:P51178 1- PHOSPHATIDYLINOSITOL-4,5- BISPHOSPHATE PHOSPHODIESTERASE DELTA 1 (EC 3.1.4.11) (PLC-DELTA-1) (PHOSPHOLIPASE C-DELTA-1) (PLC-III) - HOMO SAPIENS (HUMAN), 756 aa.	0	3
254	cg43992911	382	CTAAAGTCGGAGT ATCTTCTTCCAA/G/ AJATTTCACGTCTT GGCGGCCGTTCCA	G	A				SILENT- NONCOD ING	glycoprotein	Human Gene SWISSPROT- ID:P08183 MULTIDRUG RESISTANCE PROTEIN 1 (P- GLYCOPROTEIN 1) - HOMO SAPIENS (HUMAN), 1280 aa.	0	7
255	cg43932434	267	TTTCTAGAGGGGG TCTGTGAAGAT/G/ AJTGTAAGTAGTAC ACCCCAAGCCCCCA	G	A				SILENT- NONCOD ING	glycoprotein	Human Gene SWISSPROT- ID:P16070 CD44 ANTIGEN PRECURSOR (PHAGOCYTIC GLYCOPROTEIN 1) (PGP-1) (HUTCH-1) (EXTRACELLULAR MATRIX RECEPTOR-III) (ECMR-III) (GP90 LYMPHOCYTE HOMING/ADHESION RECEPTOR) (HERMES ANTIGEN) (HYALURONATE RECEPTOR) (HEPARAN SULFATE PROTEOGLYCAN) (EPICAN) (CDW44) - HOMO SAPIENS (HUMAN), 742 aa.	1.80E-195	11 (11pter)

256	cg43932434	306	CCCCAACCCCAA CCTCAGTGGAAAT /GJCAATGCCAGG GATTAGGCTATGG A	A	G			SILENT- NONCOD ING	glycoprotein	Human Gene SWISSPROT- ID:P16070 CD44 ANTIGEN PRECURSOR (PHAGOCYTIC GLYCOPROTEIN I) (PGP-I) (HUTCH-I) (EXTRACELLULAR MATRIX RECEPTOR-III) (ECMR-III) (GP90 LYMPHOCYTE HOMING/ADHESION RECEPTOR) (HERMES ANTIGEN) (HYALURONATE RECEPTOR) (HEPARAN SULFATE PROTEOGLYCAN) (EPICAN) (CDW44) - HOMO SAPIENS (HUMAN), 742 aa.	1.80E-195	11 (11pter)
257	cg43318219	366	GCGCAGGTCAGAG GGCGGCCGACGI A/GJGGCCTCCGG AGGTCCCCACGCC GG	A	G			SILENT- NONCOD ING	glycoprotein	Human Gene SWISSNEW- ID:P15813 T-CELL SURFACE GLYCOPROTEIN CD1D PRECURSOR (CD1D ANTIGEN) (R3G1) - HOMO SAPIENS (HUMAN), 335 aa.lpcis:SWISSPROT-ID:P15813 T-CELL SURFACE GLYCOPROTEIN CD1D PRECURSOR (CD1D ANTIGEN) (R3G1) - HOMO SAPIENS (HUMAN), 335 aa.	3.10E-185	1 (1q21)

238	cg43967861	1934	CTCTATACTGTACAA CTCACCCATAAT/g ap]TCAAAACAATT CACCATGGTATAA A	T	gap			SILENT- NONCOD ING	glycoprotein	Human Gene Similar to SWISSPROT-ID:Q08878 FIBULIN-1, ISOFORM C PRECURSOR (BASEMENT- MEMBRANE PROTEIN 90) (BM- 90) - MUS MUSCULUS (MOUSE), 685 aa.	8.20E-67	2
239	cg43967861	1935	TCTATACTGTACAC TCACCCATAAT/g ap]CAAAACAATTAC ACCATGGTATAAA G	T	gap			SILENT- NONCOD ING	glycoprotein	Human Gene Similar to SWISSPROT-ID:Q08878 FIBULIN-1, ISOFORM C PRECURSOR (BASEMENT- MEMBRANE PROTEIN 90) (BM- 90) - MUS MUSCULUS (MOUSE), 685 aa.	8.20E-67	2
260	cg43965366	1411	GCCGAATAGCCTG GGTTTGGAAAAGIC /TATGTTTTTGAAA TATGTGGGATCTC	C	T			SILENT- NONCOD ING	glycoprotein	Human Gene Similar to SWISSPROT-ID:P49222 ERYTHROCYTE MEMBRANE PROTEIN BAND 4.2 (P4.2) (PALLIDIN) - MUS MUSCULUS (MOUSE), 690 aa.	8.90E-61	6 (6p25)
261	cg43965366	385	TACTGACCTAAAT CACACCCCTAGACIA /TTATCAGAGGGA AATTCTGACCATA A	A	T			SILENT- NONCOD ING	glycoprotein	Human Gene Similar to SWISSPROT-ID:P49222 ERYTHROCYTE MEMBRANE PROTEIN BAND 4.2 (P4.2) (PALLIDIN) - MUS MUSCULUS (MOUSE), 690 aa.	8.90E-61	6 (6p25)

262	cg43322513	1255	TGTCCTTGAAGAA CATGCACTTGGCJA /GJCGGATGGCACA AGCAAAATGGTAG A	A	G				SILENT- NONCOD ING	glycoprotein	Human Gene Similar to SWISSPROT-ID:P13983 EXTENSIN PRECURSOR (CELL WALL HYDROXYPROLINE- RICH GLYCOPROTEIN) - NICOTIANA TABACUM (COMMON TOBACCO), 620 aa.	3.30E-54	12
263	cg41637704	1397	CCCGCGCCCCAGT AGGAGCCCCGCGlg ap/GJCCAGCAGGT GCGCGCGCACGG AG	gap	G				SILENT- NONCOD ING	homeobox	Human Gene SWISSPROT- ID:P50219 HOMEBOX PROTEIN HB9 - HOMO SAPIENS (HUMAN), 401 aa.	1.20E-224	7
264	cg41637704	1423	CCAGCAGGTGCGG CGCGCACGGAGCig ap/GJCGCGGCGCGG CGGCTTCTCCCGG AG	gap	G				SILENT- NONCOD ING	homeobox	Human Gene SWISSPROT- ID:P50219 HOMEBOX PROTEIN HB9 - HOMO SAPIENS (HUMAN), 401 aa.	1.20E-224	7
265	cg41637704	1817	TGAAACTTGAAAC CGCCTCTGGAGC[C /TJGCCAATCTGCAG AGTATTGGAAAA	C	T				SILENT- NONCOD ING	homeobox	Human Gene SWISSPROT- ID:P50219 HOMEBOX PROTEIN HB9 - HOMO SAPIENS (HUMAN), 401 aa.	1.20E-224	7
266	cg43980506	939	TCCAAGAAAGGGT CATGGAAGCTTAT /CJTGGGAATAATC CTCTCAATTAGAA A	T	C				SILENT- NONCOD ING	homeobox	Human Gene TREMBLNEW- ID:G2896172 LIM HOMEBOX PROTEIN COFACTOR - HOMO SAPIENS (HUMAN), 373 aa.	1.60E-206	

267	cg43961305	100	GGGGGGTTTTT TTTTTCTCTG[G/T] TTTTTTTTTTTT TTTTTTTTT	G	T		SILENT- NONCOD ING	hydrolase	Human Gene SWISSPROT- ID:P37980 INORGANIC PYROPHOSPHATASE (EC 3.6.1.1) (PYROPHOSPHATE PHOSPHO- HYDROLASE) (PPASE) - BOS TAURUS (BOVINE), 289 aa.	1.30E-156	10
268	cg43998672	503	CTGGGGGTTTCG GGGAGGAACCAAG G/gap]GGCTCACGG AGCCTCCTGTGCTG CA	G	gap		SILENT- NONCOD ING	hydroxysteroid	Human Gene SPTREMBL- ID:Q13194 11-BETA- HYDROXYSTEROID DEHYDROGENASE TYPE 2 - HOMO SAPIENS (HUMAN), 405 aa.	2.00E-220	16 (16q22)
269	cg43998672	505	GGGGGTTTTCGGG GAGGAACCAAGG G/gap]CTCACGGAG CCTCCTGTGCTGCA GT	G	gap		SILENT- NONCOD ING	hydroxysteroid	Human Gene SPTREMBL- ID:Q13194 11-BETA- HYDROXYSTEROID DEHYDROGENASE TYPE 2 - HOMO SAPIENS (HUMAN), 405 aa.	2.00E-220	16 (16q22)
270	cg42908571	1031	GAGTTAATTATGT AAGTCATATTT[TATATTTTAAAG AGTACCACTTGAA	gap	T		SILENT- NONCOD ING	interleukin	Human Gene Homologous to SWISSPROT-ID:P05231 INTERLEUKIN-6 PRECURSOR (IL-6) (B-CELL STIMULATORY FACTOR 2) (BSF-2) (INTERFERON BETA-2) (HYBRIDOMA GROWTH FACTOR) - HOMO SAPIENS (HUMAN), 212 aa.	3.40E-108	7 (7p21)

271	cg42908571	1178	CTTACCTCAAATA AATGGCTAACTTga pTJATACATATTT TAAAGAAATATTT A	gap	T				SILENT- NONCOD ING	interleukin	Human Gene Homologous to SWISSPROT-ID:P05231 INTERLEUKIN-6 PRECURSOR (IL-6) (B-CELL STIMULATORY FACTOR 2) (BSF-2) (INTERFERON BETA-2) (HYBRIDOMA GROWTH FACTOR) - HOMO SAPIENS (HUMAN), 212 aa.	3.40E-108	7 (7p21)
272	cg42164914	1617	CAGCCCCATTGT GGTCACAGGAAGT /CJAGAGGAGGCCA CGTTCCTACTAGTT	T	C				SILENT- NONCOD ING	interleukinrecept	Human Gene SWISSPROT- ID:P25025 HIGH AFFINITY INTERLEUKIN-8 RECEPTOR B (IL-8R B) (CXCR-2) (GRO/MGSA RECEPTOR) (IL-8 RECEPTOR TYPE 2) - HOMO SAPIENS (HUMAN), 360 aa.	9.60E-191	2 (2q35)
273	cg43958501	1133	CCCAACCTGGGTTT GGCAGACATCAIA/ GJAATGATGGAGTA CATTITGCAGATA	A	G				SILENT- NONCOD ING	isomerase	Human Gene SWISSPROT- ID:P46926 PUTATIVE GLUCOSAMINE-6-PHOSPHATE ISOMERASE (EC 5.3.1.10) (GLUCOSAMINE- 6- PHOSPHATE DEAMINASE) (OSCILLIN) (K1AA0060) - HOMO SAPIENS (HUMAN), 289 aa.	1.60E-156	5

274	cg43958501	805	CACCCCCAGGTTCT CCTAGTTCAGA/G/ AJAAAAGCTGTGA AAGTGAAGAAGG A	G	A			SILENT- NONCOD ING	isomerase	Human Gene SWISSPROT- ID:P46926 PUTATIVE GLUCOSAMINE-6-PHOSPHATE ISOMERASE (EC 5.3.1.10) (GLUCOSAMINE- 6- PHOSPHATE DEAMINASE) (OSCILLIN) (KIAA0060) - HOMO SAPIENS (HUMAN), 289 aa.	1.60E-136	5
275	cg43090990	2710	TTTATTCTATTGCT ATCTGTGGATG/T/G JGTAAATGGCTGGG GGGCCAGCCCTG	T	G			SILENT- NONCOD ING	kinase	Human Gene SWISSPROT- ID:Q04759 PROTEIN KINASE C, THETA TYPE (EC 2.7.1.1-) (NPKC-THETA) - HOMO SAPIENS (HUMAN), 706 aa.	0	10
276	cg42879455	2259	AGCCTTTGTGCTCC CACTCAATACAJA/ CJAAAGGCCCTCT CTACATCTGGGAA	A	C			SILENT- NONCOD ING	kinase	Human Gene SWISSPROT- ID:Q06187 TYROSINE-PROTEIN KINASE BTK (EC 2.7.1.112) (BRUTON'S TYROSINE KINASE) (AGAMMAGLOBULINAEMIA TYROSINE KINASE) (ATK) (B CELL PROGENITOR KINASE) (BPK) - HOMO SAPIENS (HUMAN), 659 aa.	0	X (Xq21.3)

277	cg42879455	2283	AAAAAGGCCCTC TCTACATCTGGG[A] GJATGCACCTCTTC TTTGATTCCTGG	A	G				SILENT- NONCOD ING	kinase	Human Gene SWISSPROT- ID:Q06187 TYROSINE-PROTEIN KINASE BTK (EC 2.7.1.112) (BRUTON'S TYROSINE KINASE) (AGAMMAGLOBULINAEMIA TYROSINE KINASE) (ATK) (B CELL PROGENITOR KINASE) (BPK) - HOMO SAPIENS (HUMAN), 659 aa.	0		X (Xq21.3)
278	cg43971741	2151	AGCAACTTGGCTG AGCCCCACTACA[C] /TACAGAGAAATC ATCAACCTGACTT A	C	T				SILENT- NONCOD ING	kinase	Human Gene SPTREMBL- ID:Q92749 TYPE I PHOSPHATIDYLINOSITOL-4- PHOSPHATE 5-KINASE BETA (EC 2.7.1.68) (STM-7 PROTEIN) - HOMO SAPIENS (HUMAN), 540 aa.	1.40E-290	9	
279	cg43971741	2200	TAAGAGTTTCAA GATGTCAAACCTT[C] AJAGGCTGATCAGC AGATGGGATGTGA	C	A				SILENT- NONCOD ING	kinase	Human Gene SPTREMBL- ID:Q92749 TYPE I PHOSPHATIDYLINOSITOL-4- PHOSPHATE 5-KINASE BETA (EC 2.7.1.68) (STM-7 PROTEIN) - HOMO SAPIENS (HUMAN), 540 aa.	1.40E-290	9	
280	cg43971741	2451	TTTTAAAAATCCA TCCACACACAT[/TGGTAAATTAAG TATAAATTCCTTTG	gap	T				SILENT- NONCOD ING	kinase	Human Gene SPTREMBL- ID:Q92749 TYPE I PHOSPHATIDYLINOSITOL-4- PHOSPHATE 5-KINASE BETA (EC 2.7.1.68) (STM-7 PROTEIN) - HOMO SAPIENS (HUMAN), 540 aa.	1.40E-290	9	

281	cg43947749	1996	AACGTGGATTGCG ACCGTCCAACCTG /gapJGCCCCGCC TCCTACAGCTGTA AC	G	gap			SILENT- NONCOD ING	kinase	Human Gene SWISSPROT- ID:P49840 GLYCOGEN SYNTHASE KINASE-3 ALPHA (EC 2.7.1.37) (GSK-3 ALPHA) - HOMO SAPIENS (HUMAN), 483 aa.	5.60E-267	19
282	cg43947749	1997	ACGTGGATTGCA CCGTCCAACCTG /gapJGCCCCGCCCTC CTACAGCTGTAAC T	G	gap			SILENT- NONCOD ING	kinase	Human Gene SWISSPROT- ID:P49840 GLYCOGEN SYNTHASE KINASE-3 ALPHA (EC 2.7.1.37) (GSK-3 ALPHA) - HOMO SAPIENS (HUMAN), 483 aa.	5.60E-267	19
283	cg44131752	1535	CACTTAATACCAG AGACCCCCCCCCG ap/CJTCCCTCCCT CTTCCCTCCCT	gap	C			SILENT- NONCOD ING	kinase	Human Gene SPTREMBL- ID:Q15599 TYROSINE KINASE ACTIVATOR PROTEIN 1 (TKA- 1) - HOMO SAPIENS (HUMAN), 450 aa.	7.80E-173	16
284	cg43917718	306	AGACGTGTCTGCC ACAGGTCTCAGGIA /GTTAACAGATGCC CTGTCCACTGAGA G	A	G			SILENT- NONCOD ING	kinase	Human Gene Similar to SPTREMBL-ID:Q15599 TYROSINE KINASE ACTIVATOR PROTEIN 1 (TKA- 1) - HOMO SAPIENS (HUMAN), 450 aa.	1.40E-79	17
285	cg43928048	1876	TTTGATGGAAAGG TTGTCCACACTG/G/ A/GAAATTACACAC ACTTGATCAGGAA	G	A			SILENT- NONCOD ING	kinase	Human Gene Similar to SWISSPROT-ID:P20505 30 KD PROTEIN KINASE HOMOLOG (EC 2.7.1.-) (PROTEIN B1) - VACCINIA VIRUS (STRAIN COPENHAGEN), 300 aa.	5.30E-35	

286	cg42714731	208	CCCTCCGGATTGG GCGCGCGTGCGGJC /MJCCGCCCGGAGT GAGGGTTTTCGTG G	C	M			SILENT- NONCOD ING	kinaseinhibitor	Human Gene Similar to SWISSPROT-ID:P42771 CYCLIN-DEPENDENT KINASE 4 INHIBITOR A (CDK4) (P16- INK4) (P16-INK4A) (MULTIPLE TUMOR SUPPRESSOR 1) (MTS1) - HOMO SAPIENS (HUMAN), 156 aa.	2.60E-53	9 (9p21)
287	cg43322545	2943	TCCAAGCTAAGCA CTGCCACTGGGJA /GJAAACTCCACCTT CCCACTTTCCAC	A	G			SILENT- NONCOD ING	kinasereceptor	Human Gene SWISSNEW- ID:P30530 TYROSINE-PROTEIN KINASE RECEPTOR UFO PRECURSOR (EC 2.7.1.112) (AXL ONCOGENE) - HOMO SAPIENS (HUMAN), 887 aa. pcis:SWISSPROT-ID:P30530 TYROSINE-PROTEIN KINASE RECEPTOR UFO PRECURSOR (EC 2.7.1.112) (AXL ONCOGENE) - HOMO SAPIENS (HUMAN), 887 aa.	0	19 (19q13.1)

288	cg43322545	3037	CCAGCTCCATCCCA GACAGGTCCTTC/ G/CCTTCTCTGTG CAGTAGCATCAAC	C	G				SILENT- NONCOD ING	kinasereceptor	Human Gene SWISSNEW- ID:P30530 TYROSINE-PROTEIN KINASE RECEPTOR UFO PRECURSOR (EC 2.7.1.112) (AXL ONCOGENE) - HOMO SAPIENS (HUMAN), 887 aa.lpcsl:SWISSPROT-ID:P30530 TYROSINE-PROTEIN KINASE RECEPTOR UFO PRECURSOR (EC 2.7.1.112) (AXL ONCOGENE) - HOMO SAPIENS (HUMAN), 887 aa.	0	19 (19q13.1)
289	cg43322545	3038	CACCTCCATCCCA GACAGGTCCTTC/ G/CCTTCTCTGTG AGTAGCATCACCT	C	G				SILENT- NONCOD ING	kinasereceptor	Human Gene SWISSNEW- ID:P30530 TYROSINE-PROTEIN KINASE RECEPTOR UFO PRECURSOR (EC 2.7.1.112) (AXL ONCOGENE) - HOMO SAPIENS (HUMAN), 887 aa.lpcsl:SWISSPROT-ID:P30530 TYROSINE-PROTEIN KINASE RECEPTOR UFO PRECURSOR (EC 2.7.1.112) (AXL ONCOGENE) - HOMO SAPIENS (HUMAN), 887 aa.	0	19 (19q13.1)
290	cg43980494	1040	GTCTGATAGAAGA GGAGCAGGAGAA/ A/GJCAAAATCGTTA AAACCTAGCGAAT TC	A	G				SILENT- NONCOD ING	kinesin	Human Gene SPTREMBL- ID:Q14807 KID (KINESIN-LIKE DNA BINDING PROTEIN) - HOMO SAPIENS (HUMAN), 665 aa.	0	16

291	cg43925424	374	TCAGGAGCAAGGC GAATGTATGACAIA /CJCATGTCCACAAT GGTGTACATAAAG	A	C				SILENT- NONCOD ING	kinesin	Human Gene SWISSPROT- ID:Q07866 KINESIN LIGHT CHAIN (KLC) - HOMO SAPIENS (HUMAN), 569 aa.	1.90E-304	14
292	cg42479188	305	TTCGTAAGAGGCT GACGATTTACT/A GJTCTCATTTTTTC CTTCTCCAGAA	A	G				SILENT- NONCOD ING	MHC	Human Gene Homologous to SWISSPROT-ID:P13765 HLA CLASS II HISTOCOMPATIBILITY ANTIGEN, DO BETA CHAIN PRECURSOR - HOMO SAPIENS (HUMAN), 273 aa.	3.40E-147	6 (6p21.3)
293	cg42686658	1167	CTAGCTTCCCTCC CATTCAACACA/A CJACACACATCTT GCTCTACCCAAAG	A	C				SILENT- NONCOD ING	MHC	Human Gene Homologous to SWISSPROT-ID:P06340 HLA CLASS II HISTOCOMPATIBILITY ANTIGEN, DZ ALPHA CHAIN PRECURSOR (MHC DN-ALPHA) - HOMO SAPIENS (HUMAN), 250 aa.	3.70E-134	6 (6p21.3)
294	cg38337333	1122	TGTCTCAAAACCA GCTTGCCAGCTCT/ CJAATGTACCAGCA GCTGGAATCTGAA	T	C				SILENT- NONCOD ING	MHC	Human Gene Homologous to SPTREMBL-ID:Q95368 HLA CLASS I INHIBITORY NK RECEPTOR - HOMO SAPIENS (HUMAN), 455 aa.	1.80E-113	19
295	cg27803682	2506	GTGGCTGGGCTAT TCCATCCATCTGT/ GJAAGCACATTGA GCCTCCAGGCTTC	T	G				SILENT- NONCOD ING	misc_channel	Human Gene Similar to SPTREMBL-ID:P91197 SIMILAR TO LIGAND-GATED IONIC CHANNEL PROTEIN - CAENORHABDITIS ELEGANS, 461 aa.	3.50E-81	

296	cg21413267	1440	CGAGCGGCACCCA GAGCCTGCACCCIT /GJCCCTCACCGTCC TTCTGCGTCCCCC	T	G			SILENT- NONCOD ING	misc_channel	Human Gene Similar to SPTREMBL-ID:P91197 SIMILAR TO LIGAND-GATED IONIC CHANNEL PROTEIN - CAENORHABDITIS ELEGANS, 461 aa.	7.90E-79	
297	cg21413267	1860	AGGAGCCCTCTTC GGTGTCCTCCGAGIT /CJGCCACGGTCAA GACCGGCAGCACC A	T	C			SILENT- NONCOD ING	misc_channel	Human Gene Similar to SPTREMBL-ID:P91197 SIMILAR TO LIGAND-GATED IONIC CHANNEL PROTEIN - CAENORHABDITIS ELEGANS, 461 aa.	7.90E-79	
298	cg21413267	1890	CGGTCAAGACCCG CAGCACCAAGC A/GJCCGCCCGGC ACCTGCCCTGTGG C	A	G			SILENT- NONCOD ING	misc_channel	Human Gene Similar to SPTREMBL-ID:P91197 SIMILAR TO LIGAND-GATED IONIC CHANNEL PROTEIN - CAENORHABDITIS ELEGANS, 461 aa.	7.90E-79	
299	cg42481172	1541	GAGCCGTGTGGCT GTGGCCTCCGGGA /CJGGCGGTGGACG GCGTGGCTTCATC	A	C			SILENT- NONCOD ING	misc_channel	Human Gene Similar to SPTREMBL-ID:P91197 SIMILAR TO LIGAND-GATED IONIC CHANNEL PROTEIN - CAENORHABDITIS ELEGANS, 461 aa.	2.30E-71	1
300	cg39518465	89	GGGTGCACGGCCG GCCCTGGGCAGGIG ap/CJGTAGCCATGG AGCTGTGGCGCCA AT	gap	C			SILENT- NONCOD ING	oncogene	Human Gene SWISSPROT- ID:P15498 VAV PROTO- ONCOGENE - HOMO SAPIENS (HUMAN), 846 aa.	0	

301	cg41972699	627	ATGGGGCCGGTGT CTGCCAGGAGGIG ap/CJGCAGACCCGG CTCCAGGGCCAGC GC	gap	C				SILENT- NONCOD ING	oncogene	Human Gene Similar to SWISSPROT-ID:Q64010 PROTO- ONCOGENE C-CRK (P38) (ADAPTER MOLECULE CRK) - MUS MUSCULUS (MOUSE), 304 aa.	2.40E-84	22 (22q11)
302	cg40333812	235	AGCATTGAGGAA GCATAACTGACGIC /TGTGAAGGGGT GTGGGGTACTTGC C	C	T				SILENT- NONCOD ING	oncogene	Human Gene Similar to SWISSPROT-ID:P31695 NEUROGENIC LOCUS NOTCH HOMOLOG PROTEIN 4 PRECURSOR (TRANSFORMING PROTEIN INT-3) - MUS MUSCULUS (MOUSE), 1964 aa.	1.40E-62	
303	cg43280482	2295	AGCATCTGCAGAC GACCCCCGACGIC /CJTTCCTCGGAC CCCCCTCGAAGCC	A	C				SILENT- NONCOD ING	oncogene	Human Gene Similar to TREMBL NEW-ID:G2952331 ARG/ABL-INTERACTING PROTEIN ARBP2A - HOMO SAPIENS (HUMAN), 666 aa.	3.90E-62	8
304	cg44014837	22	CACTGCTGTGCAG GGCAGGGAJA/TGTC TCCAGGCAGACAG CCCAGCAAAG	A	T				SILENT- NONCOD ING	oxidase	Human Gene SWISSNEW- ID:P08684 CYTOCHROME P450 3A4 (EC 1.14.14.1) (CYP1A4) (NIFEDIPINE OXIDASE) (NF-25) (P450-PCN1) - HOMO SAPIENS (HUMAN), 502 aa. pcis:SWISSPROT-ID:P08684 CYTOCHROME P450 IIA4 (EC 1.14.14.1) (NIFEDIPINE OXIDASE) (NF-25) (P450-PCN1) - HOMO SAPIENS (HUMAN), 502 aa.	8.00E-257	

305	cg41626506	3178	CAGCACAGCGAGC GCTCTCAATTCTG[A/ gap]CCTTTTTCCTC TTCTCAGCCAACT	A	gap				SILENT- NONCOD ING	peroxidase	Human Gene SWISSPROT- ID:P07202 THYROID PEROXIDASE PRECURSOR (EC 1.11.1.8) (TPO) - HOMO SAPIENS (HUMAN), 933 aa.	0	3 (3q26.3)
306	cg43918944	2938	TCTGTAGAGCTCTG AAAAGGTTGACTT/ GJATATAGAGGTCT TGTATGTTTTAC	T	G				SILENT- NONCOD ING	phosphatase	Human Gene SPTREMBL- ID:Q15172 PROTEIN PHOSPHATASE 2A B56-ALPHA - HOMO SAPIENS (HUMAN), 486 aa.	4.60E-246	1
307	cg43988365	1537	GACAGACGAGACA GTGAGGTATGTG[A /GJGGCTGCTCCGG AATGGTCCGGAGG C	A	G				SILENT- NONCOD ING	phosphatase	Human Gene SWISSPROT- ID:Q14642 TYPE I INOSITOL- 1,4,5-TRISPHOSPHATE 5- PHOSPHATASE (EC 3.1.3.56) (SPTASE) - HOMO SAPIENS (HUMAN), 412 aa. pcds:SPTREMBL-ID:Q14642 INOSITOL 1,4,5-TRIPHOSPHATE 5-PHOSPHATASE - HOMO SAPIENS (HUMAN), 412 aa.	2.60E-227	10
308	cg43969460	581	TAACTATGCAAGA CAAGACTTGGTC[C /GJTCACGTTCCGCT CTCTAGTTGATT	C	G				SILENT- NONCOD ING	phosphatase	Human Gene SWISSPROT- ID:P36876 PROTEIN PHOSPHATASE PP2A, 55 KD REGULATORY SUBUNIT, ALPHA ISOFORM (PROTEIN PHOSPHATASE PP2A B SUBUNIT ALPHA ISOFORM) (ALPHA-PR55) - RATTUS NORVEGICUS (RAT), 447 aa.	1.90E-202	

309	cg43933809	362	AATTAAACTCTA GGTGATACTTAT/ CJATGGAAGTAGT TATTTCCTATTA	T	C				SILENT- NONCOD ING	phosphatase	Human Gene SWISSPROT- ID:P37140 SERINE/THREONINE PROTEIN PHOSPHATASE PP1- BETA CATALYTIC SUBUNIT (EC 3.1.3.16) (PP-1B) - HOMO SAPIENS (HUMAN), RATTUS NORVEGICUS (RAT), MUS MUSCULUS (MOUSE), 327 aa.	1.60E-181	2 (2p23)
310	cg43931444	215	TGCTCGCGCCGTG CCACTAAGGTCAJC /TJTCCCGCCTCCGA GAGCCAGAGCCG	C	T				SILENT- NONCOD ING	phosphatase b	Human Gene Similar to SWISSPROT-ID:P39687 POTENT HEAT-STABLE PROTEIN PHOSPHATASE 2A INHIBITOR HPP2A (HLA-DR ASSOCIATED PROTEIN 1) (PHAP1) (ACIDIC NUCLEAR PHOSPHOPROTEIN PP32) (CEREBELLAR LEUCINE RICH ACIDIC NUCLEAR PROTEIN) - HOMO SAPIENS (HUMAN), 249 aa.	1.20E-89	9
311	cg42937321	1977	CTTTTCCCTCTTAC CCTCTCTCTCTG/TJ AACATCGTAAACA ACAGACTTACGT	G	T				SILENT- NONCOD ING	potassium_chan nel	Human Gene SWISSPROT- ID:P22001 VOLTAGE-GATED POTASSIUM CHANNEL PROTEIN KV1.3 (HPCN3) (HGK5) (HUKIII) (HLK3) - HOMO SAPIENS (HUMAN), 523 aa.	5.40E-284	1 (1p21)

312	cg42937321	1983	CCTCTTACCCTCTC TCTCTGAACAT[CT]GTAACAACACAGA CTTACGTTAAACT	C	T			SILENT- NONCOD ING	potassium_chan nel	Human Gene SWISSPROT- ID:P22001 VOLTAGE-GATED POTASSIUM CHANNEL PROTEIN KV1.3 (HPCN3) (HGK5) (HUK1) (HLK3) - HOMO SAPIENS (HUMAN), 523 aa.	5.40E-284	1 (1p21)
313	cg40991963	1357	CAAAATGTAACAG TGGCTTTTCAAC[A/ G]GGAGTAAAGCA AAGTCTCTAAAGC T	A	G			SILENT- NONCOD ING	potassium_chan nel	Human Gene SWISSPROT- ID:P48048 ATP-SENSITIVE INWARD RECTIFIER POTASSIUM CHANNEL 1 (POTASSIUM CHANNEL, INWARDLY RECTIFYING, SUBFAMILY 1, MEMBER 1) (ATP-REGULATED POTASSIUM CHANNEL ROM- K) (KIR1.1) - HOMO SAPIENS (HUMAN), 391 aa.	1.80E-205	11 (11q24)

314	cg43951366	2332	AAAGATGTTTGA TACTTAACACTG/ AJTCACAAGATGG AAATGCTGAAAG	G	A				SILENT- NONCOD ING	prostaglandin	Human Gene SWISSNEW- ID:P33354 PROTAGLANDIN G/H SYNTHASE 2 PRECURSOR (EC 1.14.99.1) (CYCLOOXYGENASE -2) (COX- 2) (PROTAGLANDIN- ENDOPEROXIDE SYNTHASE 2) (PROTAGLANDIN H2 SYNTHASE 2) (PGH SYNTHASE 2) (PGHS-2) (PHS II) - HOMO SAPIENS (HUMAN), 604 aa.lpcis:SPTRMBL-ID:Q16876 PROTAGLANDIN ENDOPEROXIDE SYNTHASE-2 PRECURSOR (EC 1.14.99.1) (PROTAGLANDIN- ENDOPEROXIDE SYNTHASE) (PROTAGLANDIN SYNTHASE) (PROTAGLANDIN G/H SYNTHASE) - HOMO SAPIENS (HUMAN), 604 aa	0	1 (1q25.2)
-----	------------	------	--	---	---	--	--	--	--------------------------	---------------	---	---	---------------

315	cg43951366	2829	TGGTGGAGCCACT GCAGTGTATCTTT/ CJAAAATAAGAAT ATTGTGTGAGATA	T	C			SILENT- NONCOD ING	prostaglandin	Human Gene SWISSNEW- ID:P33354 PROSTAGLANDIN G/H SYNTHASE 2 PRECURSOR (EC 1.14.99.1) (CYCLOOXYGENASE -2) (COX- 2) (PROSTAGLANDIN- ENDOPEROXIDE SYNTHASE 2) (PROSTAGLANDIN H2 SYNTHASE 2) (PGH SYNTHASE 2) (PGHS-2) (PHS II) - HOMO SAPIENS (HUMAN), 604 aa. pcis:SPTREMBL-ID:Q16876 PROSTAGLANDIN ENDOPEROXIDE SYNTHASE-2 PRECURSOR (EC 1.14.99.1) (PROSTAGLANDIN- ENDOPEROXIDE SYNTHASE) (PROSTAGLANDIN SYNTHASE) (PROSTAGLANDIN G/H SYNTHASE) - HOMO SAPIENS (HUMAN), 604 aa.	0	1 (1q25.2)
316	cg43306254	1431	CACTTAACTTGCAAT GTGCACAGCTTT/C JTGCTAACAAATAT CGCTAAACCTTA	T	C			SILENT- NONCOD ING	prostaglandin	Human Gene SPTREMBL- ID:O00325 PROSTAGLANDIN EP3 RECEPTOR SUBTYPE ISOFORM - HOMO SAPIENS (HUMAN), 402 aa.	1.40E-211	1 (1p31.2)

317	cg43306254	1666	ATGTGATTAAATTAT GTGATGAAAACAC/A/ TJTITTTTATAAAT GATCTTGGTCTAT	A	T			SILENT- NONCOD ING	prostaglandin	Human Gene SPTREMBL- ID:000325 PROSTAGLANDIN EP3 RECEPTOR SUBTYPE ISOFORM - HOMO SAPIENS (HUMAN), 402 aa.	1.40E-211	1 (1p31.2)
318	cg42918089	1064	CAATCAGAATTGA TAAGCACTGTTC/C/ TJTCCACTCCATT AGCAATTATGTC	C	T			SILENT- NONCOD ING	protease	Human Gene Homologous to SWISSNEW-ID:P09237 MATRILYSIN PRECURSOR (EC 3.4.24.23) (PUMP-1 PROTEASE) (UTERINE METALLOPROTEINASE) (MATRIX METALLOPROTEINASE-7) (MMP-7) (MATRIN) - HOMO SAPIENS (HUMAN), 267 aa. pcls: SWISSPROT-ID:P09237 MATRILYSIN PRECURSOR (EC 3.4.24.23) (PUMP-1 PROTEASE) (UTERINE METALLOPROTEINASE) (MATRIX METALLOPROTEINASE-7) (MMP-7) (MATRIN) - HOMO SAPIENS (HUMAN), 267 aa.	2.40E-146	11 (11q21)
319	cg44032168	1703	TCCATCCCTCTTT GGGCTCTCTG/C TAGGGAAGTAACA TTTACTGAGCACC	G	C			SILENT- NONCOD ING	protease	Human Gene Similar to SWISSPROT-ID:P25155 COAGULATION FACTOR X PRECURSOR (EC 3.4.21.6) (STUART FACTOR) (VIRUS ACTIVATING PROTEASE) (VAP) - GALLUS GALLUS (CHICKEN), 475 aa.	2.40E-82	2 (2q13)

320	cg43154190	1250	TACCGGAAGTTG AGCTCAATTTCAT/ C/TTCGTGTTTCTGG CCACAACACTGCCA	T	C				SILENT- NONCOD ING	protease	Human Gene Similar to SWISSPROT-ID:P50280 MATRILYSIN PRECURSOR (EC 3.4.24.23) (PUMP-1 PROTEASE) (UTERINE METALLOPROTEINASE) (MATRIX METALLOPROTEINASE-7) (MMP-7) (MATRIN) - RATTUS NORVEGICUS (RAJ), 267 aa.	2.40E-39	11 (11q22)
321	cg43927549	175	CCAGTGCTGCGG CTCTACTGGGGA /CJGTGCGCTGGTC GGAAGATTGCTGG A	A	C				SILENT- NONCOD ING	reductase	Human Gene Homologous to SWISSPROT-ID:P16083 NAD(P)H DEHYDROGENASE (QUINONE) 2 (EC 1.6.99.2) (QUINONE REDUCTASE) (DT- DIAPHORASE) (AZOREDUCTASE) (PHYLLOQUINONE REDUCTASE) (MENADIONE REDUCTASE) - HOMO SAPIENS (HUMAN), 231 aa.	1.60E-124	6 (6pter)
322	cg43927549	191	TACTGGGAGTGC GCTGGTCGGAAGlg ap/G/ATTGCTGGAC TCGCTGAAGAGAG AC	gap	G				SILENT- NONCOD ING	reductase	Human Gene Homologous to SWISSPROT-ID:P16083 NAD(P)H DEHYDROGENASE (QUINONE) 2 (EC 1.6.99.2) (QUINONE REDUCTASE) (DT- DIAPHORASE) (AZOREDUCTASE) (PHYLLOQUINONE REDUCTASE) (MENADIONE REDUCTASE) - HOMO SAPIENS (HUMAN), 231 aa.	1.60E-124	6 (6pter)

323	cg43927549	52	CGGTCCGTGGTCC CCGGGGGCGCAGGg ap/GJTGGCAGCGCT CCCGCCCTCCAGG CG	gap	G				SILENT- NONCOD ING	reductase	Human Gene Homologous to SWISSPROT-ID:P16083 NAD(P)H DEHYDROGENASE (QUINONE) 2 (EC 1.6.99.2) (QUINONE REDUCTASE) (DT- DIAPHORASE) (AZOREDUCTASE) (PHYLLOQUINONE REDUCTASE) (MENADIONE REDUCTASE) - HOMO SAPIENS (HUMAN), 231 aa.	1.60E-124	6 (6pter)
324	cg43947066	780	TTCTCAAAAGGCT GGGGGTATTATJA /GJTAAGAACTTATT CCAAAGTGACTCT	A	G				SILENT- NONCOD ING	struct	Human Gene SWISSPROT- ID:O15142 ACTIN-LIKE PROTEIN 2 - HOMO SAPIENS (HUMAN), 394 aa.	3.30E-207	2
325	cg43923264	113	AGGAAAGCCGGAG AATTGGGACACG[C /gap]AAGAGGGGGG GCTTTGATGACCC GC	C	gap				SILENT- NONCOD ING	struct	Human Gene SWISSPROT- ID:Q14012 CALCIUM/CALMODULIN- DEPENDENT PROTEIN KINASE TYPE I (EC 2.7.1.123) (CAM KINASE I) - HOMO SAPIENS (HUMAN), 370 aa.	1.70E-200	3
326	cg43942332	1926	AGATTCATCAGAA TAGGATTTTGC[A/ CJAAATCCACCCCA TATGCTGTTGAGC	A	C				SILENT- NONCOD ING	struct	Human Gene Homologous to SPTREMBL-ID:000379 DELTA- CATENIN - HOMO SAPIENS (HUMAN), 792 aa.	2.10E-124	11

327	cg43274705	580	CCGCTGTCTCTGTGTC TTCGCTTTTAA[G/T] TCAAGAAGAAATAA TGCGACGAAAAT	G	T			SILENT- NONCOD ING	struct	Human Gene Homologous to SPTRMBL-ID:Q28910 MUCIN - BOS TAURUS (BOVINE), 600 aa (fragment).	4.80E-110	12
328	cg42207316	146	CCACTTCTCTGGGA CACATTCGCTT[CT JTGTTTCTCCAGC ATGCGCTTGCTC	C	T			SILENT- NONCOD ING	struct	Human Gene Similar to SWISSNEW-ID:P12273 PROLACTIN-INDUCIBLE PROTEIN PRECURSOR (SECRETORY ACTIN-BINDING PROTEIN) (SABP) (GROSS CYSTIC DISEASE FLUID PROTEIN 15) (GCDFF-15) (GPI7) - HOMO SAPIENS (HUMAN), 146 aa.lpcis:SWISSPROT-ID:P12273 PROLACTIN-INDUCIBLE PROTEIN PRECURSOR (SECRETORY ACTIN-BINDING PROTEIN) (SABP) (GROSS CYSTIC DISEASE FLUID PROTEIN 15) (GCDFF-15) - HOMO SAPIENS (HUMAN), 146 aa.	3.50E-74	7 (7q32)
329	cg43927885	546	CATCATCATCATA GTTTACTTCAGC[A/ T]CTTAAATCCCCG AGGAGTCTGCCCT	A	T			SILENT- NONCOD ING	struct	Human Gene Similar to SWISSPROT-ID:P19065 SYNAPTOBREVIN 2 (VESICLE ASSOCIATED MEMBRANE PROTEIN 2) (VAMP-2) - HOMO SAPIENS (HUMAN), AND BOS TAURUS (BOVINE), 115 aa.	1.20E-35	17

330	cg40388639	5029	CTCTTGCCAGCCG GCTGCAAGTTTgap /TGTAAAGCGCGG ACAGACACTGCTG A	gap	T				SILENT- NONCOD ING	synthase	Human Gene SWISSPROT- ID:P29473 NITRIC-OXIDE SYNTHASE, BRAIN (EC 1.14.13.39) (NOS, TYPE I) (NEURONAL NOS) (NNOS) - HOMO SAPIENS (HUMAN), 1434 aa.	0	12 (12q24.2)
331	cg43949316	555	AGGTTACCAACA GGAATACACAC[C /TTCCTCCCTTT CTGCTCTAGAAGG	C	T				SILENT- NONCOD ING	synthase	Human Gene SWISSPROT- ID:P48651 PHOSPHATIDYL SERINE SYNTHASE I (SERINE- EXCHANGE ENZYME I) (EC 2.7.8.-) (KJAA0024) - HOMO SAPIENS (HUMAN), 473 aa.	9,80E-269	8
332	cg43958714	1565	TGGGTGATGATCA CTGTGCTGCTTGTT CJGGCTCATGGCAG AGCATTCAAGTGCC	T	C				SILENT- NONCOD ING	synthase	Human Gene Similar to SPTREMBL-ID:Q42761 SQUALENE SYNTHASE (EC 2.5.1.21) (FARNESYL- DIPHOSPHATE FARNESYLTRANSFERASE) (FARNESYLTRANSFERASE) (PRESQUALENE-DI- DIPHOSPHATE SYNTHASE) - GLYCRRHIZA GLABRA, 412 aa.	9,20E-83	8

333	cg43275028	2508	ACAGACTGGCTGC AGCATTAGGAATC /TJAGGTCATTCCGA AACTCATCATTGA	C	T			SILENT- NONCOD ING	synthase	Human Gene Similar to SWISSPROT-ID:P70490 MILK FAT GLOBULE-EGF FACTOR 8 PRECURSOR (MFG-E8) (O- ACETYL GD3 GANGLIOSIDE SYNTHASE) (AGS) (MFGM) - RATTUS NORVEGICUS (RAT), 427 aa.lpcis:SPTREMBL- ID:P70490 O-ACETYL GD3 GANGLIOSIDE SYNTHASE - RATTUS NORVEGICUS (RAT), 427 aa.	3.20E-65	1 (1q23)
334	cg43275028	2535	GGTCATTCCGAAA CTCATCATTGAAAT/ CJCAGGAAGAAGA AGAGITCAATCTT A	T	C			SILENT- NONCOD ING	synthase	Human Gene Similar to SWISSPROT-ID:P70490 MILK FAT GLOBULE-EGF FACTOR 8 PRECURSOR (MFG-E8) (O- ACETYL GD3 GANGLIOSIDE SYNTHASE) (AGS) (MFGM) - RATTUS NORVEGICUS (RAT), 427 aa.lpcis:SPTREMBL- ID:P70490 O-ACETYL GD3 GANGLIOSIDE SYNTHASE - RATTUS NORVEGICUS (RAT), 427 aa.	3.20E-65	1 (1q23)

335	cg43275028	2601	AGAAATGGCACTGA ATTCGTTTCTCTCA/ GJAACACAGATATA ATTGTTGGTTCAA	A	G				SILENT- NONCOD ING	synthase	Human Gene Similar to SWISSPROT-ID:P70490 MILK FAT GLOBULE-EGF FACTOR 8 PRECURSOR (MFG-E8) (O- ACETYL GD3 GANGLIOSIDE SYNTHASE) (AGS) (MFGM) - RATTUS NORVEGICUS (RAT), 427 aa, pcis:SPTREMBL- ID:P70490 O-ACETYL GD3 GANGLIOSIDE SYNTHASE - RATTUS NORVEGICUS (RAT), 427 aa.	3.20E-65	1 (1q23)
336	cg43275028	2873	CTTTCACITGGTGC TGGAGAATTCA/A/ GJAAGTCAAGAAC ATGCTAAGCATAA G	A	G				SILENT- NONCOD ING	synthase	Human Gene Similar to SWISSPROT-ID:P70490 MILK FAT GLOBULE-EGF FACTOR 8 PRECURSOR (MFG-E8) (O- ACETYL GD3 GANGLIOSIDE SYNTHASE) (AGS) (MFGM) - RATTUS NORVEGICUS (RAT), 427 aa, pcis:SPTREMBL- ID:P70490 O-ACETYL GD3 GANGLIOSIDE SYNTHASE - RATTUS NORVEGICUS (RAT), 427 aa.	3.20E-65	1 (1q23)

337	cg43275028	2894	TTCAAAAAGTCAAG AACATGCTAAGC/A /GJTAAGGGACCCA AGGTAGAAAAGAGA T	A	G				SILENT- NONCOD ING	synthase	Human Gene Similar to SWISSPROT-ID:P70490 MILK FAT GLOBULE-EGF FACTOR 8 PRECURSOR (MFG-E8) (O- ACETYL GD3 GANGLIOSIDE SYNTHASE) (AGS) (MFGM) - RATTUS NORVEGICUS (RAT), 427 aa. pcis:SPTREMBL- ID:P70490 O-ACETYL GD3 GANGLIOSIDE SYNTHASE - RATTUS NORVEGICUS (RAT), 427 aa.	3.20E-65	1 (1q23)
338	cg43275028	3073	TTCTCCTTCCAGAA TGAGGCCCTGGJA/ GJAGGACCCTCTTA GTGATCTGTACT	A	G				SILENT- NONCOD ING	synthase	Human Gene Similar to SWISSPROT-ID:P70490 MILK FAT GLOBULE-EGF FACTOR 8 PRECURSOR (MFG-E8) (O- ACETYL GD3 GANGLIOSIDE SYNTHASE) (AGS) (MFGM) - RATTUS NORVEGICUS (RAT), 427 aa. pcis:SPTREMBL- ID:P70490 O-ACETYL GD3 GANGLIOSIDE SYNTHASE - RATTUS NORVEGICUS (RAT), 427 aa.	3.20E-65	1 (1q23)

339	cg43275028	5590	ACTACATAAGGAC AGCAACATGCCTA /GJTGGACATGAGA GAATTGTCTTACT	A	G				SILENT- NONCOD ING	synthase	Human Gene Similar to SWISSPROT-ID:P70490 MILK FAT GLOBULE-EGF FACTOR 8 PRECURSOR (MFG-E8) (O- ACETYL GD3 GANGLIOSIDE SYNTHASE) (AGS) (MFGM) - RATTUS NORVEGICUS (RAT), 427 aa.lpcis:SPREMBL- ID:P70490 O-ACETYL GD3 GANGLIOSIDE SYNTHASE - RATTUS NORVEGICUS (RAT), 427 aa.	3.20E-65	1 (1q23)
340	cg43985000	1856	GAAAAAATCACA AGGCAACTGTGA[C /GJTCCGGGAATCT CTTCTCTGATCCTT	C	G				SILENT- NONCOD ING	tm7	Human Gene SWISSPROT- ID:P25101 ENDOTHELIN-1 RECEPTOR PRECURSOR (ET-A) - HOMO SAPIENS (HUMAN), 427 aa.	1.60E-236	4
341	cg39565524	1684	TCCGACCCACAC ACCTGAGGGAG[C /GJCCTACCCTAGCC TCAGCCGCTCCTG	C	G				SILENT- NONCOD ING	tm7	Human Gene SWISSPROT- ID:P51575 P2X PURINOCEPTOR 1 (ATP RECEPTOR) (P2X1) (PURINERGIC RECEPTOR) - HOMO SAPIENS (HUMAN), 399 aa.	2.00E-220	17
342	cg43306266	1603	ATAATCCATGCCTC TGAATATTAGAT/ GJTGTTTCTTGGG TGGGATTTTGAAT	T	G				SILENT- NONCOD ING	tm7	Human Gene SWISSPROT- ID:P43115 PROSTAGLANDIN E2 RECEPTOR, EP3 SUBTYPE (PROSTANOID EP3 RECEPTOR) (PGE RECEPTOR, EP3 SUBTYPE) - HOMO SAPIENS (HUMAN), 390 aa.	4.80E-212	1 (1p31.2)

343	cg43306266	1641	GGGATTTTGAATA TGCAATTTAAGAAIG ap/CJGTTGGGAAGA ATTTCACAGATGA TG	gap	C				SILENT- NONCOD ING	tm7	Human Gene SWISSPROT- ID:P43115 PROSTAGLANDIN E2 RECEPTOR, EP3 SUBTYPE (PROSTANOID EP3 RECEPTOR) (PGE RECEPTOR, EP3 SUBTYPE) - HOMO SAPIENS (HUMAN), 390 aa.	4.80E-212	1 (1p31.2)
344	cg43306266	1650	GAATATGCAATTA AGAAAGTTGGGAAG G/CJAATTTTCACAG ATGATGATTGGAG GA	G	C				SILENT- NONCOD ING	tm7	Human Gene SWISSPROT- ID:P43115 PROSTAGLANDIN E2 RECEPTOR, EP3 SUBTYPE (PROSTANOID EP3 RECEPTOR) (PGE RECEPTOR, EP3 SUBTYPE) - HOMO SAPIENS (HUMAN), 390 aa.	4.80E-212	1 (1p31.2)

345	cg43329467	683	TCGGCAAATCTTG AAAGCTGCAGG C/TJGCAGAGACAT GGATGTGACTTCC CA	C	T			SILENT- NONCOD ING	tm7	Human Gene SWISSNEW- ID:Q99527 CHEMOKINE RECEPTOR-LIKE 2 (IL8- RELATED RECEPTOR DRY12) (FLOW-INDUCED ENDOTHELIAL G PROTEIN- COUPLED RECEPTOR) (FEG-1) (G PROTEIN-COUPLED RECEPTOR GPR30) (GPCR-BR) - HOMO SAPIENS (HUMAN), 375 aa.lpcis:SWISSPROT- ID:Q99527 CHEMOKINE RECEPTOR-LIKE 2 (IL8- RELATED RECEPTOR DRY12) (FLOW-INDUCED ENDOTHELIAL G PROTEIN- COUPLED RECEPTOR) (FEG-1) (G PROTEIN-COUPLED RECEPTOR GPR30) - HOMO SAPIENS (HUMAN), 375 aa.lpcis:TREMBLNEW- ID:G2656121 G-PROTEIN COUPLED RECEPTOR - HOMO SAPIENS (HUMAN), 375 aa.	8.20E-201	7
346	cg2751286	439	AAGGCATAAGAAC TAGGAGCTGCTG ap/GJACATTTCAT ATGAAGGGCAACT CC	gap	G			SILENT- NONCOD ING	tm7	Human Gene SWISSPROT- ID:P50052 TYPE-2 ANGIOTENSIN II RECEPTOR (AT2) - HOMO SAPIENS (HUMAN), 363 aa.	2.00E-197	X

347	cg11731407	76	GAATGTGGGGATA AGGCATTGGGACIC /TCTATCAGGTATC CTGAGGAGAGACT	C	T			SILENT- NONCOD ING	tm7	Human Gene SWISSPROT- ID:P46089 PROBABLE G PROTEIN-COUPLED RECEPTOR GPR3 (ACCA ORPHAN RECEPTOR) - HOMO SAPIENS (HUMAN), 330 aa.	3.20E-176	1
348	cg43326635	135	CAGCCGGGAGCTC TGCCAGCTTTGGIC/ TGAAGGAGGGTG CTTGCCCTCGTGCCC	C	T			SILENT- NONCOD ING	tm7	Human Gene SWISSPROT- ID:P30542 ADENOSINE A1 RECEPTOR - HOMO SAPIENS (HUMAN), 326 aa.	1.10E-173	1
349	cg43326635	139	CGGGAGCTCTGCC AGCTTTGGCGAAIG /CJGAGGGTGCTTG CCTCGTGCCCCCTTG	G	C			SILENT- NONCOD ING	tm7	Human Gene SWISSPROT- ID:P30542 ADENOSINE A1 RECEPTOR - HOMO SAPIENS (HUMAN), 326 aa.	1.10E-173	1
350	cg43993798	1839	TGCTCTTGCTGCTG ATGGAGGAGGAJA/ GJGGGGTGGATCCC GTGGAGCCTCCAA	A	G			SILENT- NONCOD ING	tm7	Human Gene Homologous to SWISSPROT-ID:P31421 METABOTROPIC GLUTAMATE RECEPTOR 2 PRECURSOR - RATTUS NORVEGICUS (RAT), 872 aa.	6.90E-109	3 (3q21)

351	cg43040271	2130	ATGCTTCCCCAAC CCTAGGGAATC(A/ C)ACACTTAAGATA ATTGCCCACTTCT	A	C				SILENT- NONCOD ING	tm7	Human Gene Similar to SWISSPROT-ID:Q25322 TYRAMINE/OCTOPAMINE RECEPTOR 2 (TYR-LOC 2) - LOCUSTA MIGRATORIA (MIGRATORY LOCUST), 484 aa. pcls:SPTREMBL-ID:Q25322 GCR2 (G PROTEIN-COUPLED RECEPTOR) - LOCUSTA MIGRATORIA (MIGRATORY LOCUST), 484 aa.	2.90E-74	
352	cg43040271	2139	CCAACCCCTAGGGA ATCAACACTTAA/G /TJATAATTCGCCAC TTCTCCTCTTCT	G	T				SILENT- NONCOD ING	tm7	Human Gene Similar to SWISSPROT-ID:Q25322 TYRAMINE/OCTOPAMINE RECEPTOR 2 (TYR-LOC 2) - LOCUSTA MIGRATORIA (MIGRATORY LOCUST), 484 aa. pcls:SPTREMBL-ID:Q25322 GCR2 (G PROTEIN-COUPLED RECEPTOR) - LOCUSTA MIGRATORIA (MIGRATORY LOCUST), 484 aa.	2.90E-74	

353	cg43040271	2163	AGATAATTGCCA CTTCTCCTCTTTTC/ TTTCTCTGCTCCGC TCACGGCTTGACG	C	T			SILENT- NONCOD ING	tm7	Human Gene Similar to SWISSPROT-ID:Q25322 TYRAMINE/OCTOPAMINE RECEPTOR 2 (TYR-LOC 2) - LOCUSTA MIGRATORIA (MIGRATORY LOCUST), 484 aa, pfcl:SPTEMBL-ID:Q25322 GCR2 (G PROTEIN-COUPLED RECEPTOR) - LOCUSTA MIGRATORIA (MIGRATORY LOCUST), 484 aa.	2.90E-74	
354	cg43040273	1668	CGCAGAGCCCGC CGTGGGTCCGCC/T/ CJGCTGAGGCGCC CCAGCCAGTGCGC	T	C			SILENT- NONCOD ING	tm7	Human Gene Similar to SWISSPROT-ID:Q24563 DOPAMINE RECEPTOR 2 - DROSOPHILA MELANOGASTER (FRUIT FLY), 539 aa.	2.00E-58	5 (5q32)
355	cg43040273	1760	CAGCGCCTTCTTGC TGGCACCCCAATTA/ GJGAAGCCATGCCG CGGACCACGACGT	A	G			SILENT- NONCOD ING	tm7	Human Gene Similar to SWISSPROT-ID:Q24563 DOPAMINE RECEPTOR 2 - DROSOPHILA MELANOGASTER (FRUIT FLY), 539 aa.	2.00E-58	5 (5q32)
356	cg43040273	1793	TGCGCCGGACAC GACGTCACGACG/C /GJAAAGGACGAG GTGTGGGTGGTG G	C	G			SILENT- NONCOD ING	tm7	Human Gene Similar to SWISSPROT-ID:Q24563 DOPAMINE RECEPTOR 2 - DROSOPHILA MELANOGASTER (FRUIT FLY), 539 aa.	2.00E-58	5 (5q32)

357	cg43040273	2767	GCAGGTCTTCTTTG AAGGCTATGGIG/ CJAATGGCTACTCC AGCAACGGCAACA	G	C			SILENT- NONCOD ING	tm7	Human Gene Similar to SWISSPROT-ID:Q24563 DOPAMINE RECEPTOR 2 - DROSOPHILA MELANOGASTER (FRUIT FLY), 539 aa.	2.00E-58	5 (5q32)
358	cg43040273	2953	ATTGTAGTACAAA TGACTCACTGCTIG/ AJTAAAGCAGTTTT TCTACTTTTAAAG	G	A			SILENT- NONCOD ING	tm7	Human Gene Similar to SWISSPROT-ID:Q24563 DOPAMINE RECEPTOR 2 - DROSOPHILA MELANOGASTER (FRUIT FLY), 539 aa.	2.00E-58	5 (5q32)
359	cg43040273	3053	ATAAACTTAGAAT AAAAATTGTAAAAG ap/AJTGTATAGAG ATATGCAGAAGGA AG	gap	A			SILENT- NONCOD ING	tm7	Human Gene Similar to SWISSPROT-ID:Q24563 DOPAMINE RECEPTOR 2 - DROSOPHILA MELANOGASTER (FRUIT FLY), 539 aa.	2.00E-58	5 (5q32)
360	cg43998970	1501	AGGGGTGGAAC TG CTGATGGGATT[ga pTJCCTTCATCC TTCTGATAAAGGT A	gap	T			SILENT- NONCOD ING	transcriptfactor	Human Gene SPTREMBL- ID:Q07279 TRANSCRIPTION FACTOR NF-E2 - MUS MUSCULUS (MOUSE), 373 aa.	1.70E-177	12
361	cg43998970	249	AGCCTCCCCAGAG ACAAACCCGGGAI G/CJCCTCATCTCTC TCCTCACCCCTGCTG	G	C			SILENT- NONCOD ING	transcriptfactor	Human Gene SPTREMBL- ID:Q07279 TRANSCRIPTION FACTOR NF-E2 - MUS MUSCULUS (MOUSE), 373 aa.	1.70E-177	12

362	cg43947199	2623	GTCTCTCCGGGCC CACCCGCTGGICT TAAGGGGAAGTGG GCGAAGCTGGAGC	C	T				SILENT- NONCOD ING	transcriptfactor	Human Gene SWISSNEW- ID:P23193 TRANSCRIPTION ELONGATION FACTOR S-II (TRANSCRIPTION ELONGATION FACTOR A) - HOMO SAPIENS (HUMAN), 301 aa. pcls:SWISSPROT-ID:P23193 TRANSCRIPTION ELONGATION FACTOR S-II (TRANSCRIPTION ELONGATION FACTOR A) - HOMO SAPIENS (HUMAN), 301 aa.	4.20E-158	8
363	cg43917396	934	GGGGCCGGGCACT GCCAGGAAGGGI A/GJCTCCGGGAGA GGGAGCCGGCGGC TG	A	G				SILENT- NONCOD ING	transcriptfactor	Human Gene Similar to TREMBLNEW-ID:G2920821 TRANSCRIPTION FACTOR T- BOX 5 - HOMO SAPIENS (HUMAN), 518 aa.	6.90E-68	
364	cg40351913	2030	AGACGAAGACCCC AGGAAGTCATCCIT /CJGCAATGGGAGA GACACGAACAAAC C	T	C				SILENT- NONCOD ING	transport	Human Gene SWISSPROT- ID:Q01959 SODIUM- DEPENDENT DOPAMINE TRANSPORTER (DA TRANSPORTER) (DAT) - HOMO SAPIENS (HUMAN), 620 aa.	0	5 (5p15.3)
365	cg43921289	237	CCGAGGCTGCCA GGAGCAAGCCGAIg ap/AJGAGCCAGCCG GCCGGGCACTCC GA	gap	A				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P02545 LAMIN A (70 KD LAMIN) - Homo sapiens (Human), 664 aa.	0	1

366	cg43928515	3196	AAACAAATAAGCC CTTTTACTGAC/A/ GJATGCACCAACC TTTTCAGCTGAAG	A	G				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:Q14687 HYPOTHETICAL PROTEIN KIAA0182 - Homo sapiens (Human), 1157 aa (fragment).	0	16
367	cg43955093	1309	AGAGTCAAAAATC CAAGTTGGATTTC/ GJTAAGCAGCCTTG ACAGTAATCACTG	C	G				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene SPTREMBL- ACC:Q16084 P130 - HOMO SAPIENS (HUMAN), 1139 aa.	0	16
368	cg43955093	1336	AAGCAGCCTTGAC AGTAATCACTGA/A /GJTGATAGGAAA AAAAGACAGTTGG G	A	G				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene SPTREMBL- ACC:Q16084 P130 - HOMO SAPIENS (HUMAN), 1139 aa.	0	16
369	cg43925474	2206	AGGCAAAAGCTCA CAGTAAATGTATTA /CJCCAGAACAGGG GCCTAAGTGAAGG T	A	C				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P42566 EPIDERMAL GROWTH FACTOR RECEPTOR SUBSTRATE SUBSTRATE 15 (PROTEIN EPS15) (AF-IP PROTEIN) - Homo sapiens (Human), 896 aa.	0	1 (1p32)
370	cg44014437	4893	CTGCTCCCACTTC GCCAGCCTCCA/A/ GJGTACAACCTCC GCGTGTAGTGGGC	A	G				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P53675 CLATHRIN HEAVY CHAIN 2 (CLH-22) - Homo sapiens (Human), 1640 aa.	0	17 (17q11)

371	cg44014448	5114	CTGCTCCCAACTTC GCCAGCCTCCAIA/ GJGTACAACTTCC GGGTGTAGTGGC	A	G				SILENT- NONCOD ING	UNCLASSIFIE D	Human Gene SWISSPROT- ACC:P53675 CLATHRIN HEAVY CHAIN 2 (CLH-22) - Homo sapiens (Human), 1640 aa.	0	17 (17q11)
372	cg43973129	2242	CACTTCACTGAAA GACACCATTTATC/ ATACCCAAGGCA GAAAGTAGAACTT	C	A				SILENT- NONCOD ING	UNCLASSIFIE D	Human Gene SWISSPROT- ACC:P05060 SECRETORANIN I PRECURSOR (SGI) (CHROMOGRANIN B) - Homo sapiens (Human), 677 aa.	0	20 (20pter)
373	cg43950637	1939	GATAGGACTCAAG CTTATTTGGGATC/ TCTGATCAATTCT TTCTGATGTGT	C	T				SILENT- NONCOD ING	UNCLASSIFIE D	Human Gene SWISSNEW- ACC:Q13009 T-LYMPHOMA INVASION AND METASTASIS INDUCING PROTEIN 1 (TIAMI PROTEIN) - Homo sapiens (Human), 1591 aa.	0	21 (21q22.1)
374	cg43956384	2416	TACAGCCATCTGT ACCTACTGGAGCIC /TGCAGAAAGGAA GTCCACTCAGTCA C	C	T				SILENT- NONCOD ING	UNCLASSIFIE D	Human Gene SWISSPROT- ACC:P13866 SODIUM/GLUCOSE COTRANSPORTER 1 (NA(+)/GLUCOSE COTRANSPORTER 1) (HIGH AFFINITY SODIUM-GLUCOSE COTRANSPORTER) - Homo sapiens (Human), 664 aa.	0	22 (22q13.1)

375	cg43992229	101	AGCAGTGCAGCCC CGGCGCGGAGCAI G/AJGGAGCTCGG CCCGCGCCCGCG CC	G	A				SILENT- NONCOD ING	UNCLASSIFIE D	Human Gene SWISSPROT- ACC:P23352 KALLMANN SYNDROME PROTEIN PRECURSOR (ADHESION MOLECULE-LIKE X-LINKED) - Homo sapiens (Human), 680 aa.	0	X (Xp22.3)
376	cg44932392	260	GAGAAAAAGCATG GTACCCAAACCGAIA /TITTCACCTTTC AGCAATACTTCAC	A	T				SILENT- NONCOD ING	UNCLASSIFIE D	Human Gene TREMBLNEW- ACC:AAD23581 CULLIN 2 - HOMO SAPIENS (HUMAN), 745 aa.	0	
377	cg44932392	323	TAAAGTTTAAAG AATGTCATAATGIA /TTCATGAGCTTGA AATATCTCTAGGC A	A	T				SILENT- NONCOD ING	UNCLASSIFIE D	Human Gene TREMBLNEW- ACC:AAD23581 CULLIN 2 - HOMO SAPIENS (HUMAN), 745 aa.	0	
378	cg43981656	1121	AGCAAAGAAACAC TGCGAGAATTCIA /TGCATTGCAAA ATTCTAAGTTTGG	A	T				SILENT- NONCOD ING	UNCLASSIFIE D	Human Gene TREMBLNEW- ACC:CAA08974 GUANINE NUCLEOTIDE-EXCHANGE FACTOR - HOMO SAPIENS (HUMAN), 548 aa.	1.60E-292	10
379	cg44910613	366	AAATAAATGTTTTC ATAGTCATTACIT/A JCTTTACAAATGGGA GTGCTAAAATTC	T	A				SILENT- NONCOD ING	UNCLASSIFIE D	Human Gene SWISSPROT- ACC:P38567 HYALURONIDASE PRECURSOR (EC 3.2.1.35) (SPERM SURFACE PROTEIN PH-20) (SPERM ADHESION MOLECULE 1) - Homo sapiens (Human), 509 aa.	1.20E-280	7

380	cg44035104	189	AACTGGGTTGCTCT AAGAACTGATG(T) CCTAAACCGTCTC AGCATGGCCTGTA	T					SILENT- NONCOD ING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P37287 N- ACETYLGLUCOSAMINYL- PHOSPHATIDYLINOSITOL BIOSYNTHETIC PROTEIN (GLCNAC-P1 SYNTHESIS PROTEIN) (PHOSPHATIDYLINOSITOL GLYCAN COMPLEMENTATION CLASS A) (PIG-A) - Homo sapiens (Human), 484 aa.	4.70E-261	X (Xp22.1)
381	cg43929959	1643	CAATGCATGAATC TGTACCCCTTCGG(G) gapJAGGGCACTCAC ATGCCGCCCCAC C	G	gap				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene SPTREMBL- ACC:P78506 DIABETES MELLITUS TYPE I AUTOANTIGEN (ISLET CELL AUTOANTIGEN P69) - HOMO SAPIENS (HUMAN), 483 aa.	2.10E-258	7
382	cg43950250	1961	TTGTTTCATGATTTC TTGATGTTCTC(T) pJTAATGGAAACT AAGAGATGGAATT	C	gap				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P11926 ORNITHINE DECARBOXYLASE (EC 4.1.1.17) (ODC) - Homo sapiens (Human), 461 aa.	7.00E-251	2 (2p25)
383	cg43064090	129	GCCGAGTCCGCTG GTGGCGGACCC(A /TAGGGGAGCAGC CAGTAGGGAAGTT G	A	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P32754 4- HYDROXYPHENYLPYRUVATE DIOXYGENASE (EC 1.13.11.27) (4HPPD) (HPD) - Homo sapiens (Human), 392 aa.	4.80E-213	

384	cg43064090	130	CCGAGTCCGCTGG TGGCGGACCCCA[A /TJGGGAGCAGCC AGTAGGGAAGTTG G	A	T			SILENT- NONCOD ING	UNCLASSIFIED D	Human Gene SWISSPROT- ACC:P32754 4- DIOXYPHENYL PYRUVATE DIOXYGENASE (EC 1.13.1.1.27) (4HPPD) (HPD) - Homo sapiens (Human), 392 aa.	4.80E-213	
385	cg43064090	157	GGGAGCAGCCAGT AGGGAAGTTGGG[C/GJGAGTTCCAGA ATCAGGGGGCGTG GC	C	G			SILENT- NONCOD ING	UNCLASSIFIED D	Human Gene SWISSPROT- ACC:P32754 4- DIOXYPHENYL PYRUVATE DIOXYGENASE (EC 1.13.1.1.27) (4HPPD) (HPD) - Homo sapiens (Human), 392 aa.	4.80E-213	
386	cg43064090	61	TAATCGGGAGGGC TGGAGCAGAGGG[C/GJGGCCCCGCCG AGGGCGTGCTCA GT	C	G			SILENT- NONCOD ING	UNCLASSIFIED D	Human Gene SWISSPROT- ACC:P32754 4- DIOXYPHENYL PYRUVATE DIOXYGENASE (EC 1.13.1.1.27) (4HPPD) (HPD) - Homo sapiens (Human), 392 aa.	4.80E-213	
387	cg30490224	3296	GATGCCAAAAAA CAAAGGTGAGAA[A/CJCCACAACACA GGTCTAAACTCAG CA	A	C			SILENT- NONCOD ING	UNCLASSIFIED D	Human Gene SWISSPROT- ACC:P30968 GONADOTROPIN- RELEASING HORMONE RECEPTOR (GNRH-R) - Homo sapiens (Human), 328 aa.	1.20E-177	4 (4q21.2)
388	cg43924431	381	GTCITTTACAGATG GTTTTCAAAAT[apJAGAGTCCAGTA AAATATTCACATT	T	gap			SILENT- NONCOD ING	UNCLASSIFIED D	Human Gene SWISSNEW- ACC:Q16637 SURVIVAL MOTOR NEURON PROTEIN 1 - Homo sapiens (Human), 294 aa.	4.20E-166	5

389	cg43936047	607	CGTTGTTCTTAATG TGGATCTACCA[CT JCCCTGTGTTCAATC GAGATTCCGGTC	C	T			SILENT- NONCOD ING	UNCLASSIFIED D	Human Gene TREMBLNEW- ACC:AAD40550 P38IP - HOMO SAPIENS (HUMAN), 733 aa.	4.30E-164	13
390	cg43272443	1542	TGGGATTACAGGT GCGCACTACCACTA /GJCCAAAGCTAATTT TTGTATTTTITAG	A	G			SILENT- NONCOD ING	UNCLASSIFIED D	Human Gene SWISSNEW- ACC:P13726 TISSUE FACTOR PRECURSOR (TF) (COAGULATION FACTOR III) (THROMBOPLASTIN) (CD142 ANTIGEN) - Homo sapiens (Human), 295 aa.	7.70E-158	1 (1p22)
391	cg43966848	2065	CCTTCAGCAGCCCT GCAGCGGAAAT[CT TJAATGAGCCGCCG TAGCCGCCATCCG	C	T			SILENT- NONCOD ING	UNCLASSIFIED D	Human Gene SPTREMBL- ACC:Q92600 PROTEIN INVOLVED IN SEXUAL DEVELOPMENT, COMPLETE CDS - HOMO SAPIENS (HUMAN), 299 aa.	4.90E-156	2
392	cg43964140	176	AAAAAGCTACAGA AAAGAAATCACTTT /CJTGAATAAACACA ATGACTCAGAGGC A	T	C			SILENT- NONCOD ING	UNCLASSIFIED D	Human Gene Homologous to TREMBLNEW-ACC:AAC69899 SACM21 - MUS MUSCULUS (MOUSE), 721 aa.	1.10E-150	6
393	cg43285114	418	CAGGGACATGCGG GCACCCCGTGGGJG /gapJTCITTTGGCGGC TCACAGGACAATG G	G	gap			SILENT- NONCOD ING	UNCLASSIFIED D	Human Gene Homologous to TREMBLNEW-ACC:AAD23440 LR8 - HOMO SAPIENS (HUMAN), 270 aa.	1.90E-138	7

394	cg43948566	370	GCAGGCAGAGCAC CCTGGGACCCCAIG /gap]GGCAGAAAGGA CCCCTGCCCTCCAG T	G	gap			SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Homologous to SWISSNEW-ACC:P18582 CD81 ANTIGEN (26 KD CELL SURFACE PROTEIN TAPA-1) - Homo sapiens (Human), 236 aa.	3.30E-125	11
395	cg44003626	649	TAAACAGCTCAGT TCAGGGACTGGT[A /GJTACAAGCTGGC CACCCATCTCAGC C	A	G			SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Homologous to SPTREMBL-ACC:Q15025 MRNA (HA1652) FOR ORF, PARTIAL CDS - HOMO SAPIENS (HUMAN), 296 aa (fragment).	2.70E-123	
396	cg43917206	259	TTACAGGACATCA CCTGCCATCTT[A][A]GGTTTAATATT ACAAATGCCTAGT	T	A			SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Homologous to SWISSPROT-ACC:P22061 PROTEIN-L-ISOASPARTATE(D- ASPARTATE) O- METHYLTRANSFERASE (EC 2.1.1.77) (PROTEIN-BETA- ASPARTATE METHYLTRANSFERASE) (PMT) (PROTEIN L- ISOASPARTYL/D-ASPARTYL METHYLTRANSFERASE) (L- ISOASPARTYL PROTEIN CARBOXYL METHYLTRANSFERASE) - Homo sapiens (Human), 226 aa.	6.90E-118	6

397	cg43289666	215	GGCCGATTTTCCCA CAATTAAATC[CT]CAGTTCACCTGGT ATCCAGCTCCAG	C	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Homologous to SPTREMBL-ACC:O00559 CANCER ASSOCIATED SURFACE ANTIGEN - HOMO SAPIENS (HUMAN), 213 aa.	2.50E-111	8
398	cg43986282	840	GTTTCCACCTCCCC AGACAGGCATTTC/ TTCGAGTGGGAGGC GGGAGCACGTACC	C	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Homologous to SPTREMBL-ACC:P97314 DOUBLE LIM PROTEIN-1 - MUS MUSCULUS (MOUSE), 193 aa.	2.90E-110	12
399	cg43986282	841	TTTCCACCTCCOCA GACAGGCATTTC/ TTCGAGTGGGAGGC GGGAGCACGTACC G	C	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Homologous to SPTREMBL-ACC:P97314 DOUBLE LIM PROTEIN-1 - MUS MUSCULUS (MOUSE), 193 aa.	2.90E-110	12
400	cg43297716	1030	CTAAACCCAAATG GGGGCTGCTGGC[A /TTGACCCCGAGGG TGCCCTGGCCAGTC C	A	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Homologous to SWISSPROT-ACC:P15018 LEUKEMIA INHIBITORY FACTOR PRECURSOR (LIF) (DIFFERENTIATION- STIMULATING FACTOR) (D FACTOR) (MELANOMA- DERIVED LPL INHIBITOR) (MLPL) - Homo sapiens (Human), 202 aa.	1.20E-106 (22q12.1)	22

401	cg43980312	2160	TTTATCATTAAG TGCCAGAATGG[C/ TJCTTTAATGAAA ACAAAAACAAAG	C	T			SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Homologous to SWISSPROT-ACC:P34741 SYNDECAN-2 PRECURSOR (FIBROGLYCAN) (HEPARAN SULFATE PROTEOGLYCAN- CORE PROTEIN) (HSPG) (SYND2) - Homo sapiens (Human), 201 aa.	7.90E-101	8 (8q22)
402	cg43939240	624	GGAGGGTTGGAGT CACTGACGAATG[C /TJGAGCCGGGCCA GGCCCATGCAAAG G	C	T			SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SPTREMBL-ACC:O43399 HD54+INS2 ISOFORM - HOMO SAPIENS (HUMAN), 206 aa.	1.00E-100	
403	cg43941552	881	GCCACCTGCCCGG GCTGTGGAGGAG[C /gap]GCTCGCGCTG ACCAGGCGCTGGG GC	C	gap			SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSNEW-ACC:P11686 PULMONARY SURFACTANT- ASSOCIATED PROTEIN C PRECURSOR (SP-C) (SP5) (PULMONARY SURFACTANT- ASSOCIATED PROTEOLIPID SPL(VAL)) - Homo sapiens (Human), 197 aa.	1.60E-100	
404	cg43941552	1124	GCTTCTGCCACAC CGCAGGGACAA[A/ G/CCTGGAGAAAT GGGAGCNTGGGGA	A	G			SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSNEW-ACC:P11686 PULMONARY SURFACTANT- ASSOCIATED PROTEIN C PRECURSOR (SP-C) (SP5) (PULMONARY SURFACTANT- ASSOCIATED PROTEOLIPID SPL(VAL)) - Homo sapiens (Human), 197 aa.	1.60E-100	

405	cg42917153	914	CATTCTCTTTGTA CATAATACATTTC JACTCCCTGCCTC CTCTCCTTTCTA	C	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSPROT-ACC:P45973 HETEROCHROMATIN PROTEIN 1 HOMOLOG ALPHA (HP1 ALPHA) (ANTIGEN P25) - Homo sapiens (Human), 191 aa.	2.10E-100	12
406	cg43927693	878	CAGGGGTCAGCAG AGCTTCAGAGGTG /TGCCCCACCTGA GCCCCACCCGGG A	G	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSPROT-ACC:P30536 PERIPHERAL-TYPE BENZODIAZEPINE RECEPTOR (PBR) (PKBS) (MITOCHONDRIAL BENZODIAZEPINE RECEPTOR) - Homo sapiens (Human), 169 aa.	5.30E-95	22
407	cg43951338	507	CAGAAAGCAGCAA ATTAGTGTTTTC/ AJAGGACCGAATTC GGCTCCCCGAGCT	C	A				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSPROT-ACC:P36405 ADP- RIBOSYLATION FACTOR-LIKE PROTEIN 3 - Homo sapiens (Human), 182 aa.	3.40E-93	10
408	cg43951338	511	AAGCAGCAAAATTA GTGTTTTTCAGGJA/ CTCCGAATTCGGCT CCCGCAGCTCCTG	A	C				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSPROT-ACC:P36405 ADP- RIBOSYLATION FACTOR-LIKE PROTEIN 3 - Homo sapiens (Human), 182 aa.	3.40E-93	10
409	cg43951338	547	CTCCCGCAGCTCCT GCATCTCCATTTC/ TGCTAGATTTTAT TTCTTCTTTGCA	C	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSPROT-ACC:P36405 ADP- RIBOSYLATION FACTOR-LIKE PROTEIN 3 - Homo sapiens (Human), 182 aa.	3.40E-93	10

410	cg25236776	1234	CCCGCCAGCCCG ACGCCTACTGAGG ap/TJCCCCCGCTC GCCCCACCGCGC GC	gap	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSNEW-ACC:P01185 VASOPRESSIN-NEUROPHYSIN 2-COPEPTIN PRECURSOR [CONTAINS: ARG- VASOPRESSIN; NEUROPHYSIN 2 (NEUROPHYSIN-II); COPEPTIN] - Homo sapiens (Human), 164 aa.	7.20E-91	
411	cg25236776	1240	CCAGCCCGACGCC TACTGAGCCCCGJC /TJGCTCGCCACCC GGCGGCTCTTCG	C	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSNEW-ACC:P01185 VASOPRESSIN-NEUROPHYSIN 2-COPEPTIN PRECURSOR [CONTAINS: ARG- VASOPRESSIN; NEUROPHYSIN 2 (NEUROPHYSIN-II); COPEPTIN] - Homo sapiens (Human), 164 aa.	7.20E-91	
412	cg25236776	1242	AGCCCGACGCCTA CTGAGCCCCGCGJC /TJTCGCCCAACCGG CGCGCTCTTCGG	C	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSNEW-ACC:P01185 VASOPRESSIN-NEUROPHYSIN 2-COPEPTIN PRECURSOR [CONTAINS: ARG- VASOPRESSIN; NEUROPHYSIN 2 (NEUROPHYSIN-II); COPEPTIN] - Homo sapiens (Human), 164 aa.	7.20E-91	
413	cg25236776	1246	CGAGGCTACTGA GCCCCGCGCTCGJC /TJCCCAACCGCGC GCTCTTCGCGCCCG	C	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSNEW-ACC:P01185 VASOPRESSIN-NEUROPHYSIN 2-COPEPTIN PRECURSOR [CONTAINS: ARG- VASOPRESSIN; NEUROPHYSIN 2 (NEUROPHYSIN-II); COPEPTIN] - Homo sapiens (Human), 164 aa.	7.20E-91	

414	cg43968406	1362	GCTACGTTTACTCA CAGCCAGCGAA[ga pA]CTGACATTAAA ATAACTAACAAAC A	gap	A				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to REMTREMBL-ACC:E47283 DNA FOR ORF1 AND ORF2 FROM CHROMOSOME X - HOMO SAPIENS (HUMAN), 157 aa.	5.00E-83	X (Xp11.4)
415	cg42748886	104	CGCCTCTGATCCA AGCCACCTCCCGC /TJCAGAGAGGTGT CATGGGCTTCCAA A	C	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSNEW-ACC:P01238 CALCITONIN PRECURSOR - Homo sapiens (Human), 141 aa.	2.00E-70	11 (11p15.2)
416	cg43969533	356	CTCTGCACAAGGG AAGCCTATCCTAT/ gap/TTTTTTTCTCT TTGCGAAAACAGA	T	gap				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to TREMBLNEW-ACC:AAD39844 HSPC028 - HOMO SAPIENS (HUMAN), 419 aa.	1.60E-67	7
417	cg43976681	1119	AATGCCTCAGATC AGTGACCCAAGG[A/gap]ACCTTCCAG AATGGATGAAATA GAC	A	gap				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to TREMBLNEW-ACC:AAD29427 MYOMEGALIN - RATTUS NORVEGICUS (RAT), 2324 aa.	4.30E-66	11
418	cg43976681	1120	ATGCCTCAGATCA GTGACCCAAGGAI A/gap]CCTTCCAGA ATGGATGAAATAG ACC	A	gap				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to TREMBLNEW-ACC:AAD29427 MYOMEGALIN - RATTUS NORVEGICUS (RAT), 2324 aa.	4.30E-66	11

419	cg43984044	714	CCAAAGCGGAAGGC CATTTCCCTGC[C/ TCTTCTCAGTTG TCCGGGGCGGGGG	C	T				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SPTREMBL-ACC:O00455 TTF-1 INTERACTING PEPTIDE 20 - HOMO SAPIENS (HUMAN), 385 aa (fragment).	7.30E-66	19
420	cg43933283	398	CTAATTGTGTCGA ATTCCAGGATT[G/ A]GAGGAAAAGTT GCTCCCTTTCAGGC	G	A				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSPROT-ACC:P05062 FRUCTOSE-BISPHOSPHATE ALDOLASE B (EC 4.1.2.13) (LIVER-TYPE ALDOLASE) - Homo sapiens (Human), 363 aa.	6.60E-65	9 (9q22)
421	cg42381630	577	AAAGCAATCACAG TGTTAAAAGAAG[G /A]CAGTTGAAAT GATGCAGGCTGCT C	G	A				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SPTREMBL-ACC:O76087 GAGE-8 - HOMO SAPIENS (HUMAN), 117 aa.	5.90E-64	
422	cg41664708	423	CCAGCCAGCTCAT TTCACCTTACAC[G/ C]CTCATGGACTGA GTTTACTACTCACC	G	C				SILENT- NONCOD ING	UNCLASSIFIED	Human Gene Similar to SWISSNEW-ACC:P47992 LYMPHOTACTIN PRECURSOR (CYTOKINE SCM-1) (ATAC) (LYMPHOTAXIN) (SCM-1- ALPHA) - Homo sapiens (Human), 114 aa.	2.00E-54	1

423	cg43277632	3906	AGCCTTCCCGCAG AAAAAGATGCAG[C/TTCCCCAGACCT TCTCTGTGCTGATT	C	T	Ala	Val (652)	CONSER VATIVE	ATPase associated	Human Gene SWISSPROT- ID:P35670 COPPER- TRANSPORTING ATPASE 2 (EC 3.6.1.36) (COPPER PUMP 2) (WILSON DISEASE- ASSOCIATED PROTEIN) - HOMO SAPIENS (HUMAN), 1465 aa.	0	13 (13q14.3)
424	cg40310734	1138	TACCAGAGGCTGC ATCGGCTGCCG[C /GIAGAGCAGATGG CGTCGTATTTGGG	C	G	Ala	Gly (653)	CONSER VATIVE	cadherin	Human Gene SWISSPROT- ID:P08514 PLATELET MEMBRANE GLYCOPROTEIN IIB PRECURSOR (GPIIB) (INTEGRIN ALPHA- IIB) (CD41) - HOMO SAPIENS (HUMAN), 1039 aa.	0	17 (17q21.3 2)
425	cg40310734	1238	TGGTGGCGCTCC ACTGTATATGGA[G /CJAGCCGGGCAGA CCGAAAACCTGGCC G	G	C	Glu	Asp (654)	CONSER VATIVE	cadherin	Human Gene SWISSPROT- ID:P08514 PLATELET MEMBRANE GLYCOPROTEIN IIB PRECURSOR (GPIIB) (INTEGRIN ALPHA- IIB) (CD41) - HOMO SAPIENS (HUMAN), 1039 aa.	0	17 (17q21.3 2)
426	cg40310734	1893	CTCTCAACAGGCA GGCACCACCCCTG[A /GIACCTGGATCTG GGCGGAAAGCACA G	A	G	Asn	Asp (655)	CONSER VATIVE	cadherin	Human Gene SWISSPROT- ID:P08514 PLATELET MEMBRANE GLYCOPROTEIN IIB PRECURSOR (GPIIB) (INTEGRIN ALPHA- IIB) (CD41) - HOMO SAPIENS (HUMAN), 1039 aa.	0	17 (17q21.3 2)

427	cg43982507	1883	GGTTACAAAGTGTG AATGTAGTCGTGIG /CJCTATCAAATGG ATCTTGCTACTGGC	G	C	Gly	Ala (656)	CONSER VATIVE	eph	Human Gene SWISSPROT- ID:P98155 VERY LOW- DENSITY LIPOPROTEIN RECEPTOR PRECURSOR (VLDL RECEPTOR) - HOMO SAPIENS (HUMAN), 873 aa.	0	9 (9p24)
428	cg41534010	949	GCCGAGGACGTGC GTGGCAACCTGAIG /AJGGCAACACCG AGGGGCTGCAGAA G	G	A	Arg	Lys (657)	CONSER VATIVE	eph	Human Gene SWISSNEW- ID:P06727 APOLIPOPROTEIN A- IV PRECURSOR (APO-AIV) - HOMO SAPIENS (HUMAN), 396 aa. cds: SWISSPROT-ID:P06727 APOLIPOPROTEIN A-IV PRECURSOR (APO-AIV) - HOMO SAPIENS (HUMAN), 396 aa.	1.80E-203	11 (11q23)
429	cg43299024	1036	TACGAGGTGCCCT TGGAGACCCCGCJA /GJGTCCACAGCC GGGCACCGTCCCC A	A	G	His	Arg (658)	CONSER VATIVE	glucoamylase	Human Gene TREMBLNEW- ID:G2826521 MALTASE- GLUCOAMYLASE (EC 3.2.1.20) - HOMO SAPIENS (HUMAN), 1857 aa.	7.40E-199	17 (17q25.2)
430	cg43299024	1108	GAGGAGCCCTTCG GGTGATCGTGCIJA /GJCCGGCAGCTGG ACGGCCCGGTGCT G	A	G	His	Arg (659)	CONSER VATIVE	glucoamylase	Human Gene TREMBLNEW- ID:G2826521 MALTASE- GLUCOAMYLASE (EC 3.2.1.20) - HOMO SAPIENS (HUMAN), 1857 aa.	7.40E-199	17 (17q25.2)

431	cg43285373	12840	GACTGACTGGGA AAGGAACCTAAAI A/CITCGAGTCTGCC TGGATGAATGGAG A	A	C	Ile	Leu (660)	CONSER VATIVE	glycoprotein	Human Gene SWISSPROT- ID:P98164 LOW-DENSITY LIPOPROTEIN RECEPTOR- RELATED PROTEIN 2 (MEGALIN) (GLYCOPROTEIN 330) - HOMO SAPIENS (HUMAN), 1751 aa (fragment).	0	2
432	cg36834323	1004	AGTTATTCTAGAG GATACAGAAATC[A /GJTCGAAAGJTCOC GAGAAACTAGGGA G	A	G	His	Arg (661)	CONSER VATIVE	glycoprotein	Human Gene Similar to SWISSPROT-ID:P38159 HETEROGENEOUS NUCLEAR RIBONUCLEOPROTEIN G (HNRNP G) (GLYCOPROTEIN P43) - HOMO SAPIENS (HUMAN), 437 aa.	6.40E-91	
433	cg41568631	2101	GGACCAGGGGGCC ATGCTGCTCAAT[G/ AJTCTCAGGCCACG TCAAGGAGAGCGG	G	A	Val	Ile (662)	CONSER VATIVE	glycoprotein	Human Gene Similar to SWISSPROT-ID:P16452 ERYTHROCYTE MEMBRANE PROTEIN BAND 4.2 (P4.2) (PALLIDIN) - HOMO SAPIENS (HUMAN), 690 aa.	9.90E-70	14 (14q11.2)
434	cg42359655	666	TGCTTTTCAGGGCG GAAAACCTCTCT[A/ GJTTGTCCTGCGAG CTGAAGATATCCC	A	G	Ile	Val (663)	CONSER VATIVE	hydrolase	Human Gene SWISSPROT- ID:P09848 LACTASE- PHLORIZIN HYDROLASE PRECURSOR (EC 3.2.1.108) (EC 3.2.1.62) (LACTASE- GLYCOSYLCEAMIDASE) - HOMO SAPIENS (HUMAN), 1927 aa.	0	2 (2q21)

435	cg43998672	1331	GTGTGGGCTTGG TGAACTCTAGCA[C /AJCGGCTAATGT CTCCTGGTTTGGTC	C	A	Val	Leu (664)	CONSER VATIVE	hydroxysteroid	Human Gene SPTREMBL- ID:Q13194 11-BETA- HYDROXYSTEROID DEHYDROGENASE TYPE 2 - HOMO SAPIENS (HUMAN), 405 aa.	2.00E-220	16 (16q22)
436	cg43969028	1133	GGAGATGTGGTCA TTCCTAGTGATTJA/ TJTTTCAGATAGT GGGAGGAAGCAAC	A	T	Tyr	Phe (665)	CONSER VATIVE	immunoglob	Human Gene Homologous to SPTREMBL-ID:P91456 SIMILAR TO THE IMMUNOGLOBULIN SUPERFAMILY - CAENORHABDITIS ELEGANS, 1173 aa.	2.20E-149	18 (18q21.3)

437	cg43933479	133	AAGGAGAAAGAGAA AGCTGTTTATCCJA/ GJTTCATGGGTGA AGGTACAATAAAT	A	G	His	Arg (666)	CONSER VATIVE	interleukin	Human Gene SWISSNEW- ID:P29466 INTERLEUKIN-1 BETA CONVERTASE PRECURSOR (IL-1BC) (EC 3.4.22.36) (IL-1 BETA CONVERTING ENZYME) (ICE) (INTERLEUKIN-1 BETA CONVERTING ENZYME) (P45) (CASPASE-1) (CASP-1) - HOMO SAPIENS (HUMAN), 404 aa. pcis:SWISSPROT-ID:P29466 INTERLEUKIN-1 BETA CONVERTASE PRECURSOR (IL-1BC) (EC 3.4.22.36) (IL-1 BETA CONVERTING ENZYME) (ICE) (INTERLEUKIN-1 BETA CONVERTING ENZYME) (P45) (CASPASE-1) (CASP-1) - HOMO SAPIENS (HUMAN), 404 aa.	2.50E-206	
438	cg43942537	1163	GCCACTGTCTCTTC CAAAACCTTCA/C/ AJGCCCTGTCTTGC TTGTTCCTCGTCTA	C	A	Val	Leu (667)	CONSER VATIVE	kinesin	Human Gene SWISSNEW- ID:P33176 KINESIN HEAVY CHAIN (UBIQUITOUS KINESIN HEAVY CHAIN) (UKHC) - HOMO SAPIENS (HUMAN), 963 aa. pcis:SWISSPROT-ID:P33176 KINESIN HEAVY CHAIN (UBIQUITOUS KINESIN HEAVY CHAIN) (UKHC) - HOMO SAPIENS (HUMAN), 963 aa.	0	10

439	cg38337333	1035	TTCCAAATGCTGA GCCACAGCGTTG /AJTCTCCTGCCAT GAGCACACAGTC	G	A	Val	Ile (668)	CONSER VATIVE	MHC	Human Gene Homologous to SPTREMBL-ID:Q95368 HLA CLASS I INHIBITORY NK RECEPTOR - HOMO SAPIENS (HUMAN), 455 aa.	1.80E-113	19
440	cg38337333	271	CTGGAACAGTTTC CTCAATTAGCCTTG/ CJTGAACCCAGCAC ACGACGGGACCTA	G	C	Val	Leu (669)	CONSER VATIVE	MHC	Human Gene Homologous to SPTREMBL-ID:Q95368 HLA CLASS I INHIBITORY NK RECEPTOR - HOMO SAPIENS (HUMAN), 455 aa.	1.80E-113	19
441	cg38337333	823	TCATCGCTGGTGCT CCAAAAAATAA/A/ GJATGCTGCTGTAA TGAACCAAGAGCC	A	G	Asn	Asp (670)	CONSER VATIVE	MHC	Human Gene Homologous to SPTREMBL-ID:Q95368 HLA CLASS I INHIBITORY NK RECEPTOR - HOMO SAPIENS (HUMAN), 455 aa.	1.80E-113	19
442	cg30421838	3434	GGATGCTGTTGCTC TCCACACAGCCA/G/ TJGGGCGTTCCTAA ATGAAAGCCAAGC	G	T	Val	Leu (671)	CONSER VATIVE	nucl_recpt	Human Gene SWISSNEW- ID:P06401 PROGESTERONE RECEPTOR (PR) - HOMO SAPIENS (HUMAN), 933 aa;pcis:SWISSPROT-ID:P06401 PROGESTERONE RECEPTOR (PR) - HOMO SAPIENS (HUMAN), 933 aa.	0	11 (11q22)

443	cg43064060	1019	GCCAATGGCATCC AGAACAAAGGAGG[C/TJGGAGGTCCGC ATCTTTCACGTCTG C	C	T	Ala	Val (672)	CONSER VATIVE	nucl_reopt	Human Gene SWISSPROT- ID:Q07869 PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR ALPHA (PPAR- ALPHA) - HOMO SAPIENS (HUMAN), 468 aa.[pcis:SPTREMBL-ID:Q16241 PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR ALPHA - HOMO SAPIENS (HUMAN), 468 aa (fragment).	4.10E-254	22
444	cg43991813	1860	TCTCGACTAACAG CATTTCCAAAGA[T/ CJGGAGCGAATATT GTCCACGGTTGAG	T	C	Ile	Val (673)	CONSER VATIVE	nuclease	Human Gene SWISSPROT- ID:P40692 MUTL PROTEIN HOMOLOG 1 (DNA MISMATCH REPAIR PROTEIN MLH1) - HOMO SAPIENS (HUMAN), 756 aa.	0	3 (3p21.3)
445	cg42904626	194	GAGTGCCTTGACG ATACAGCTAATT[C/ GJAGAATCAATTTG TGGACGAATATGA	C	G	Gln	Glu (674)	CONSER VATIVE	oncogene	Human Gene Similar to SWISSPROT-ID:P01118 TRANSFORMING PROTEIN P21/K-RAS 2B - HOMO SAPIENS (HUMAN), 188 aa.	1.10E-97	12
446	cg42904626	548	AAGAAAGTTATGGA ATTCCCTTTATT[G/ CJAAACATCAGCAA AGACAAGACACAGGG	G	C	Glu	Gln (675)	CONSER VATIVE	oncogene	Human Gene Similar to SWISSPROT-ID:P01118 TRANSFORMING PROTEIN P21/K-RAS 2B - HOMO SAPIENS (HUMAN), 188 aa.	1.10E-97	12

447	cg42460457	2845	GCCGCCTCAGCCA GCAAAGCAGCGG[C/TTAGGCCAGTCC TAGCCACCACAGA G	C	T	Ala	Val (676)	CONSER VATIVE	phosphatase	Human Gene SWISSPROT- ID:P23470 PROTEIN-TYROSINE PHOSPHATASE GAMMA PRECURSOR (EC 3.1.3.48) (R- PTP- GAMMA) - HOMO SAPIENS (HUMAN), 1445 aa.	0	3 (3p14.2)
448	cg43272594	582	GGGATGTACTGCA TGGTGTTCTTGGT/ CJGCTGTATGTGCA GGCACGACTCTGT	T	C	Val (677)	Ala (677)	CONSER VATIVE	phosphatase	Human Gene Similar to SPTREMBL-ID:Q61469 PHOSPHATIDIC ACID PHOSPHATASE - MUS MUSCULUS (MOUSE), 283 aa.	1.40E-79	19
449	cg43958858	807	TCAGGTGGTGGGA ACCTACCGTTGC[C/ GJTTCCTGGAAGA AGGGAGGCTACAC	C	G	Leu (678)	Val (678)	CONSER VATIVE	polymerase	Human Gene SWISSNEW- ID:P25205 DNA REPLICATION LICENSING FACTOR MCM3 (DNA POLYMERASE ALPHA HOLOENZYME-ASSOCIATED PROTEIN P1) (RLF BETA SUBUNIT) (P102 PROTEIN) - HOMO SAPIENS (HUMAN), 808 aa. pcis:SWISSPROT-ID:P25205 DNA REPLICATION LICENSING FACTOR MCM3 (DNA POLYMERASE ALPHA HOLOENZYME-ASSOCIATED PROTEIN P1) (RLF BETA SUBUNIT) (P102 PROTEIN) - HOMO SAPIENS (HUMAN), 808 aa.	0	6 (6p12)

450	cg43916732	540	GTACAGCGGGCGG GCCACCTCGGGC/A /TCTGAGCACCAA TTTTGCGGGGGC G	A	T	Thr	Ser (679)	CONSER VATIVE	protease	Human Gene SPTREMBL- ID:Q15113 PROCOLLAGEN C- PROTEINASE ENHANCER PROTEIN PRECURSOR - HOMO SAPIENS (HUMAN), 449 aa.	1.20E-247	7 (7q21.3)
451	cg42894809	2745	GGATGCTGGAGAG TGGATCACTGTC/A/ GJATCAGACGACA ACAGCCAACCGTT A	A	G	Asn	Asp (680)	CONSER VATIVE	struct	Human Gene SWISSPROT- ID:P54296 M-PROTEIN (165 KD TTIN-ASSOCIATED PROTEIN) (165 KD CONNECTIN- ASSOCIATED PROTEIN) - HOMO SAPIENS (HUMAN), 1465 aa.	0	8
452	cg40388639	2337	GATTCTCCAGAG CTGGTGTGGAA/G /CJTCCCATCAGGC ACCCCAAGTTGA	G	C	Val	Leu (681)	CONSER VATIVE	synthase	Human Gene SWISSPROT- ID:P29475 NITRIC-OXIDE SYNTHASE, BRAIN (EC 1.14.13.39) (NOS, TYPE I) (NEURONAL NOS) (NNOS) - HOMO SAPIENS (HUMAN), 1434 aa.	0	12 (12q24.2)
453	cg40388639	2380	AAGTTTGAGTGGT TCAAGGACCTGGG /CJGCTGAAGTGGT ACGGCCTCCCGC C	G	C	Gly	Ala (682)	CONSER VATIVE	synthase	Human Gene SWISSPROT- ID:P29475 NITRIC-OXIDE SYNTHASE, BRAIN (EC 1.14.13.39) (NOS, TYPE I) (NEURONAL NOS) (NNOS) - HOMO SAPIENS (HUMAN), 1434 aa.	0	12 (12q24.2)

454	cg43124627	1524	AATTCTATATCAC TGGGGACAGAGC/ GATATATGGATAA AGATGGGTATTC	G	Ala	Gly (683)	CONSER VATIVE	synthase	Human Gene Similar to SWISSNEW-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE-COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa.lpcis:SWISSPROT-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE-COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa.	7.70E-79	16
455	cg43124627	869	TGGAACAAGTGGGA TATCCGAAATGTA /TJCTGCACACACCC ACAGCAGTTTGG	T	Thr	Ser (684)	CONSER VATIVE	synthase	Human Gene Similar to SWISSNEW-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE-COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa.lpcis:SWISSPROT-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE-COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa.	7.70E-79	16

456	cg43064068	1464	AGGAGAGGTGGTG AAGGCAATTGTG[G /AJTCCTGGCCTCGC AGTTCCTGTCCCA	G	A	Val	Ile (685)	CONSER VATIVE	synthase	Human Gene Similar to SWISSNEW-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE-COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa, pcis:SWISSPROT-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE-COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa.	7.40E-65	
457	cg2514276	1090	GTGATGGACCTC TCATATATGCC[T/ TJCGCAGCCAAGA GATCGGAAGACC	A	T	Tyr	Phe (686)	CONSER VATIVE	tm7	Human Gene SWISSPROT- ID:P33032 MELANOCORTIN-5 RECEPTOR (MC5-R) (MC-2) - HOMO SAPIENS (HUMAN), 325 aa.	7.00E-172	
458	cg32423505	964	TTCCATCTGAGGT TATAAACCCACG[A/ TJATTCAGGCAAG TGGCCAGAAATGGC	A	T	Phe	Tyr (687)	CONSER VATIVE	tm7	Human Gene Similar to SPTRMBL-ID:Q89609 G PROTEIN-COUPLED RECEPTOR - EQUINE HERPESVIRUS TYPE 2 (EHV-2), 383 aa.	1.20E-55	3 (3q21)
459	cg43335558	344	CAAGACCTAGCTC CCCAGCAGAGAGI C/TJGGCCCCACAA CAAAAGAGGTCCA GC	C	T	Ala	Val (688)	CONSER VATIVE	tnfreceptor	Human Gene Similar to TREMBLNEW-ID:G2653845 TNF RECEPTOR-RELATED RECEPTOR FOR TRAIL - HOMO SAPIENS (HUMAN), 386 aa.	5.50E-89	8

460	cg43998970	1347	GACAGAGCTGTAC CGTGACATTTTCIC/ GJAGCACCTTCGGG ATGAATCAGGCAA	C	G	Gln	Glu (689)	CONSER VATIVE	transcriptfactor	Human Gene SPTREMBL- ID:Q07279 TRANSCRIPTION FACTOR NF-E2 - MUS MUSCULUS (MOUSE), 373 aa.	1.70E-177	12
461	cg2537639	800	GAGGGCGATTCT ACTACCTGGGGGIG /CJGTTCTTCGGGGG GTGGTGCAAGAG	G	C	Gly	Ala (690)	CONSER VATIVE	transferase	Human Gene SWISSPROT- ID:P16442 FUCOSYLGLYCOPROTEIN ALPHA-N- ACETYL GALACTOSAMINYLT RANSFERASE (EC 2.4.1.40) (HISTO-BLOOD GROUP A TRANSFERASE) (A TRANSFERASE) / FUCOSYLGLYCOPROTEIN 3- ALPHA- GALACTOSYLTRANSFERASE (EC 2.4.1.37) (HISTO-BLOOD GROUP B TRANSFERASE) (B TRANSFERASE) (NAGAT) - HOMO SAPIENS (HUMAN), 354 aa.	6.50E-192	9 (9q34)

462	cg43935995	1552	ACTGTCCCTCACC ATGGTACCCCTG[A /G]TCACCCCTGCCTC TGCTTTTCCCTCT	A	G	Ile	Val (691)	CONSER VATIVE	transport	Human Gene SWISSPROT- ID:Q03518 ANTIGEN PEPTIDE TRANSPORTER 1 (APT1) (PEPTIDE TRANSPORTER TAP1) (PEPTIDE TRANSPORTER PSF1) (PEPTIDE SUPPLY FACTOR 1) (PSF-1) (PEPTIDE TRANSPORTER INVOLVED IN ANTIGEN PROCESSING 1) - HOMO SAPIENS (HUMAN), 748 aa.	0	6
463	cg43935986	1424	CCTGGAACGGCC TTGTACCTGCTC[G/ A]TAAGGAGGTG CTGCACTTGGGGG T	G	A	Val (692)	Ile (692)	CONSER VATIVE	transport	Human Gene SPTREMBL- ID:Q28437 ABC-TRANSPORTER - GORILLA GORILLA GORILLA (LOWLAND GORILLA), 703 aa.	0	6 (6p21.3)
464	cg43968274	730	GAGCAGGAGGAAG CCATGAATGCGG[C /T]CTACTCAGGCTA CGTCTACACGCAC	C	T	Ala (693)	Val (693)	CONSER VATIVE	UNCLASSIFIED D	Human Gene SPTREMBL- ACC:O14914 NEURONAL MUNC18-1 BINDING PROTEIN - HOMO SAPIENS (HUMAN), 837 aa.	0	9
465	cg44018598	3568	AGATACTTTCTATA AGCAGTTTTTA[G/C]ATTGTAGGAAGCA GCTGAATTCAAA	G	C	Leu (694)	Val (694)	CONSER VATIVE	UNCLASSIFIED D	Human Gene SWISSPROT- ACC:P29374 RETINOBLASTOMA BINDING PROTEIN 1 (RBBP-1) - Homo sapiens (Human), 1257 aa.	0	14

466	cg44926796	1825	ACACTGGAAAGCA CAACAGTTGGCA[C /GTTCTGCTAGAA AATAATAATTGCA	C	G	Thr (695)	CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSPROT- ACC:Q15046 LYSYL-TRNA SYNTHETASE (EC 6.1.1.6) (LYSINE-TRNA LIGASE) (LYSRS) (KIAA0070) - Homo sapiens (Human), 597 aa.	0	16
467	cg43055918	1622	AACGCTGCCCTGA CTGAGAAAGGCA[C C/TTGATGCTCGCTC CACTGCTGGAACC G	C	T	Arg (696)	CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P42694 HYPOTHETICAL PROTEIN KIAA0054 - Homo sapiens (Human), 1942 aa.	0	17
468	cg43966985	1381	CATCCAGGACAAC TTCTCGGTGACT[C/ GJAAGTGCCCTTCA CTGAGAGCGCCTG	C	G	Gln (697)	CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P01019 ANGIOTENSINOGEN PRECURSOR - Homo sapiens (Human), 485 aa.	3.90E-257	1 (1q42)
469	cg43918854	966	CTCAACCCCTGGTC GGAGACAACGGJA/ CTCACCATGGCCA TCAGAACAGTGCG	A	C	Ile (698)	CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P20062 TRANSCOBALAMIN II PRECURSOR - Homo sapiens (Human), 427 aa.	3.30E-228	22 (22q11.2)
470	cg43918484	1148	CTGATTCTTCGGTT CTTCTTGACTTC/G JTGCCACCTTGCCA GCCAGCTGCTCG	C	G	Glu (699)	CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P05089 ARGINASE 1 (EC 3.5.3.1) (LIVER-TYPE ARGINASE) - Homo sapiens (Human), 322 aa.	1.30E-171	6 (6q23)

471	cg43942977	1009	ACGGCCCTGGAGA ACCAGAAGAAGG[CTTGAGGAAGAAG AAAGTCTTGATTG CC	C	T	Ala	Val (700)	CONSER VATIVE	UNCLASSIFIED	Human Gene Homologous to SWISSNEW-ACC:Q12846 SYNTAXIN 4 - Homo sapiens (Human), 297 aa.	9.60E-148	
472	cg43942977	725	GGATGGTGTCTGA TGAGGAGTTGGA[G TTJCAGATGCTGGA CAGTGGGCAAAAGC G	G	T	Glu	Asp (701)	CONSER VATIVE	UNCLASSIFIED	Human Gene Homologous to SWISSNEW-ACC:Q12846 SYNTAXIN 4 - Homo sapiens (Human), 297 aa.	9.60E-148	
473	cg43943361	921	TTGGGGTTGGCTTG GTTTCAATAAG[G/ CJAACGGGGACACT TACAAATTGCTGC	G	C	Glu	Gln (702)	CONSER VATIVE	UNCLASSIFIED	Human Gene Homologous to SWISSNEW-ACC:P04179 SUPEROXIDE DISMUTASE [MN] PRECURSOR (EC 1.15.1.1) - Homo sapiens (Human), 222 aa.	5.70E-124	6 (6q25.3)
474	cg25236776	1094	GTGACCGAGCCCG AGTGCCGGAGG[G/TJCTTCACCGCC GCGCCCGGCCAG C	G	T	Gly	Val (703)	CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to SWISSNEW-ACC:P01185 VASOPRESSIN-NEUROPHYSIN 2-COPEPTIN PRECURSOR [CONTAINS: ARG- VASOPRESSIN; NEUROPHYSIN 2 (NEUROPHYSIN-ID); COPEPTIN] - Homo sapiens (Human), 164 aa.	7.20E-91	

475	cg25236776	881	CAGTGCCTCCCTG CGGCCCCGGG[G/ TCAAAGGCCGCTG CTTCGGGCCCAAC	G	T	Gly	Val (704)	CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to SWISSNEW-ACC:P01185 VASOPRESSIN-NEUROPHYSIN 2-COPEPTIN PRECURSOR [CONTAINS: ARG- VASOPRESSIN; NEUROPHYSIN 2 (NEUROPHYSIN-II); COPEPTIN] - Homo sapiens (Human), 164 aa.	7.20E-91	
476	cg38899722	30	GGCAACTCTGCT ATGGACACCAAG G/CCTACTCTGCTGT GCGGTCATCTGCT	G	C	Val	Leu (705)	CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to RENTREMBL-ACC:G292791 T- CELL RECEPTOR BETA PRECURSOR - HOMO SAPIENS (HUMAN), 145 aa (fragment).	5.70E-75	
477	cg11753818	253	GCCTGGAACACCA GGCTCTCTGCG[G/ AJTGTCATGCTTTG TCTCTGGGAGCA	G	A	Arg	His (706)	CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to RENTREMBL-ACC:G2104755 T CELL RECEPTOR V-BETA 23 - HOMO SAPIENS (HUMAN), 129 aa (fragment).	1.30E-66	7
478	cg2526759	519	AGCCACCCAGACC GGAGACTCGGCC[G /AJTCTACCTCTGTG CTGTGGAGGCCTA	G	A	Val	Ile (707)	CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to RENTREMBL-ACC:G33509 T CELL RECEPTOR - HOMO SAPIENS (HUMAN), 118 aa (fragment).	1.60E-54	
479	cg2526759	539	CGGCCGTCTACCTC TGTCCTGTGGA[G/ CIGCCTATCTAAC GACTACAAGCTCA	G	C	Glu	Asp (708)	CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to RENTREMBL-ACC:G33509 T CELL RECEPTOR - HOMO SAPIENS (HUMAN), 118 aa (fragment).	1.60E-54	

480	cg1902363	368	CAGAACAAAGCA AATGGAATTGGAIG /TJAGCATCCTGGTG GCCCTGCTGCAGA	T	Glu	Asp (709)	CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to SWISSPROT-ACC:P01286 SOMATOLIBERIN PRECURSOR (GROWTH HORMONE- RELEASING FACTOR) (GRF) (GROWTH HORMONE- RELEASING HORMONE) (GHRH) (SOMATOCRININ) - Homo sapiens (Human), 108 aa.	2.10E-52	
481	cg43277632	3110	GAAACCCGGAAGC ACTGTAATTGCGIA /GJGGTCTATAAAT GCACATGGCTCTG T	G	Arg	Gly (710)	NON- CONSER VATIVE	ATPase associated	Human Gene SWISSPROT- ID:P35670 COPPER- TRANSPORTING ATPASE 2 (EC 3.6.1.36) (COPPER PUMP 2) (WILSON DISEASE- ASSOCIATED PROTEIN) - HOMO SAPIENS (HUMAN), 1465 aa.	0	13 (13q14.3)

482	cg43252813	2306	TGTATTCTCTGTAAT GGGGCTGATGA[C/ TATATATGATGGT TATGGACCACCAC	C	T	Thr	Ile (711)	NON- CONSER VATIVE	ATPase_ associa ted	Human Gene SWISSPROT- ID:Q04656 COPPER- TRANSPORTING ATPASE 1 (EC 3.6.1.36) (COPPER PUMP 1) (MENKES DISEASE- ASSOCIATED PROTEIN) - HOMO SAPIENS (HUMAN), 1500 aa.[pcis:SWISSPROT- ID:Q04656 COPPER- TRANSPORTING ATPASE 1 (EC 3.6.1.36) (COPPER PUMP 1) (MENKES DISEASE- ASSOCIATED PROTEIN) - HOMO SAPIENS (HUMAN), 1500 aa.	0	X (Xq12)
483	cg43920913	929	GCCCCCTGAGCAGT CAGGACCCCGGCT[C /TCCGTCCTGAGT GCCACGATCCACG	C	T	Pro	Ser (712)	NON- CONSER VATIVE	biotindp	Human Gene SWISSPROT- ID:P05166 PROPIONYL-COA CARBOXYLASE BETA CHAIN PRECURSOR (EC 6.4.1.3) (PCCASE) (PROPANOYL- COA:CARBON DIOXIDE LIGASE) - HOMO SAPIENS (HUMAN), 539 aa.	8.20E-288	3 (3q21)
484	cg40310734	267	GGAGTGGGTGCTG CTGCTCTTGGGA[C/ GCTTGTGCTGCC CTCCAGCCTGGGC	C	G	Pro	Ala (713)	NON- CONSER VATIVE	cadherin	Human Gene SWISSPROT- ID:P08514 PLATELET MEMBRANE GLYCOPROTEIN IIB PRECURSOR (GPIIB) (INTEGRIN ALPHA- IIB) (CD41) - HOMO SAPIENS (HUMAN), 1039 aa.	0	17 (17q21.3 2)

485	cg40310734	3111	CGTGTCTCTCCCTCC CCTATGCGGTGIC/ GJCCCCGCTCAGCC TGCCCCGAGGGGA	C	G	Pro	Ala (714)	NON- CONSER VATIVE	cadherin	Human Gene SWISSPROT- ID:P08514 PLATELET MEMBRANE GLYCOPROTEIN IIB PRECURSOR (GPIIB) (INTEGRIN ALPHA- IIB) (CD41) - HOMO SAPIENS (HUMAN), 1039 aa.	0	17 (17q21.3 2)
486	cg43956560	777	GGGTACTATGGG CCCCAGTGTACAGT/ CJTGTGATTACAGT GTGAGCCTTTGGA	T	C	Phe	Leu (715)	NON- CONSER VATIVE	cadherin	Human Gene SWISSPROT- ID:P14151 L-SELECTIN PRECURSOR (LYMPH NODE HOMING RECEPTOR) (LEUKOCYTE ADHESION MOLECULE-1) (LAM-1) (LEUKOCYTE SURFACE ANTIGEN LEU-8) (TQ1) (GP90- MEL) (LEUKOCYTE- ENDOTHELIAL CELL ADHESION MOLECULE 1) (LECAM1) (CD62L) - HOMO SAPIENS (HUMAN), 372 aa.	1.00E-218	1 (1q23)

487	cg43956560	837	GCTGGGTACCATG GACTGTACTCAC[C] TJCTTTGGAAACT TCAGCTTCAGCTC	C	T	Pro	Ser (716)	NON- CONSER VATIVE	cadherin	Human Gene SWISSPROT- ID:P14151 L-SELECTIN PRECURSOR (LYMPH NODE HOMING RECEPTOR) (LEUKOCYTE ADHESION MOLECULE-1) (LAM-1) (LEUKOCYTE SURFACE ANTIGEN LEU-8) (TQ1) (GP90- MEL) (LEUKOCYTE- ENDOTHELIAL CELL ADHESION MOLECULE 1) (LECAM1) (CD62L) - HOMO SAPIENS (HUMAN), 372 aa.	1.00E-218	I (q23)
488	cg42388009	753	TGCAGAAAGGCACC ACAGAGACCGGAI A/GGGCAGGGCAA GGGCACCTCGAAG AC	A	G	Arg	Gly (717)	NON- CONSER VATIVE	cadherin	Human Gene SWISSPROT- ID:P21815 BONE SIALOPROTEIN II PRECURSOR (BSP II) (CELL-BINDING SIALOPROTEIN) (INTEGRIN- BINDING SIALOPROTEIN) - HOMO SAPIENS (HUMAN), 317 aa.	7.00E-172	4
489	cg43977436	1945	GTGTGTGTGTAAT GGTGTGGCTGTATC /TJGCTCCAAACAA GATCTTATTACTGA	C	T	Arg	Cys (718)	NON- CONSER VATIVE	calcium_channel	Human Gene SWISSPROT- ID:P21817 RYANODINE RECEPTOR, SKELETAL MUSCLE (SKELETAL MUSCLE CALCIUM RELEASE CHANNEL) - HOMO SAPIENS (HUMAN), 5032 aa.	0	

490	cg43280376	1130	CGGAAGCTGGTGT CCTACTGCCCCCA/ GJAAGGTGCAACA ACTGTTGCCCTC	A	G	Gln	Arg (719)	NON- CONSER VATIVE	carboxylase	Human Gene SWISSPROT- ID:P38435 VITAMIN K- DEPENDENT GAMMA- CARBOXYLASE (EC 6.4.-.-) (GAMMA-GLUTAMYL CARBOXYLASE) - HOMO SAPIENS (HUMAN), 738 aa.	0	2
491	cg42201364	1595	CCAGGGCCTCCAG GTCCAAGAGGCCA/ /CJCTCTGGAGAGC CTGGTCTTCCAGG G	A	C	Trp	Gly (720)	NON- CONSER VATIVE	collagen	Human Gene SWISSPROT- ID:Q03692 COLLAGEN ALPHA 1(X) CHAIN PRECURSOR - HOMO SAPIENS (HUMAN), 680 aa.	0	6
492	cg42201364	176	GTGTTTACGCTGA ACGATACCAAAC/ TJGCCCCACAGGCAT AAAAGGCCCACTA	C	T	Thr	Met (721)	NON- CONSER VATIVE	collagen	Human Gene SWISSPROT- ID:Q03692 COLLAGEN ALPHA 1(X) CHAIN PRECURSOR - HOMO SAPIENS (HUMAN), 680 aa.	0	6
493	cg40339378	2855	TCCAGGAATACCA GGTCTGCCCTGGT/ GJTTCCTGGAACAA GAGGATTAAAAGG	A	G	Ile	Thr (722)	NON- CONSER VATIVE	collagen	Human Gene SPTREMBL- ID:Q12823 A TYPE IV COLLAGEN - HOMO SAPIENS (HUMAN), 1690 aa (fragment).	0	X (Xq22)

494	cg43063256	606	AGACTGTGTTACC AACAGACCATGC(A /GGAAGTCAAGTG CGATGTGAAGGCT T	A	G	Arg	Gly (723)	NON- CONSER VATIVE	complement	Human Gene SWISSNEW- ID:P07358 COMPLEMENT COMPONENT C8 BETA CHAIN PRECURSOR - HOMO SAPIENS (HUMAN), 591 aa;pcis:SWISSPROT-ID:P07358 COMPLEMENT COMPONENT C8 BETA CHAIN PRECURSOR - HOMO SAPIENS (HUMAN), 591 aa.	0	1 (1p32)
495	cg44032748	414	CTCCAGTTCTACAA CTTGTTAAGG(A/ CJAAGCACAGTGTG GACAGGATTCCA	A	C	Lys	Gln (724)	NON- CONSER VATIVE	complement	Human Gene SWISSPROT- ID:P07357 COMPLEMENT COMPONENT C8 ALPHA CHAIN PRECURSOR - HOMO SAPIENS (HUMAN), 584 aa.	0	1 (1p32)
496	cg43049885	533	CAGTTTGGGGAC AGCCATGCACTG(A /CJGCCTCTGTAGC CTTTCAACCATGC	A	C	Glu	Ala (725)	NON- CONSER VATIVE	complement	Human Gene TREMBLNEW- ID:G386348 COMPLEMENT C6 - HOMO SAPIENS, 941 aa.	0	5 (5p13)
497	cg21644442	1347	CCAGGCTCTCCCA GGATCTCATCACT/ CJGCCCCCAGGG CCTCAGCAACCCC	T	C	Leu	Pro (726)	NON- CONSER VATIVE	csf	Human Gene SWISSPROT- ID:P09603 MACROPHAGE COLONY STIMULATING FACTOR-1 PRECURSOR (CSF-1) (MCSF) - HOMO SAPIENS (HUMAN), 554 aa.	5.00E-304	1 (1p21)

498	cg2733430	279	CCAAAGCTCCCATG ACCCAGACAACG[C /TJCCTTGAAAGACA AGCTGGGTTAACT G	C	T	Pro	Ser (727)	NON- CONSER VATIVE	csf	Human Gene Similar to SWISSNEW-ID:P08700 INTERLEUKIN-3 PRECURSOR (IL-3) (MULTIPOTENTIAL COLONY-STIMULATING FACTOR) (HEMATOPOIETIC GROWTH FACTOR) (P-CELL STIMULATING FACTOR) (MAST-CELL GROWTH FACTOR) (MCGF) - HOMO SAPIENS (HUMAN), 152 aa pcIs:SWISSPROT-ID:P08700 INTERLEUKIN-3 PRECURSOR (IL-3) (MULTIPOTENTIAL COLONY-STIMULATING FACTOR) (HEMATOPOIETIC GROWTH FACTOR) (P-CELL STIMULATING FACTOR) (MAST-CELL GROWTH FACTOR) (MCGF) - HOMO SAPIENS (HUMAN), 152 aa.	1.10E-77	5
499	cg43923204	1651	TCCACGTAGAAGC GGAAGCCGAGGTI A/GGGAGATGTAC GCATTGATGGAA GG	A	G	Tyr	His (728)	NON- CONSER VATIVE	cytochrome	Human Gene Similar to SWISSPROT-ID:P21592 CYTOCHROME C OXIDASE ASSEMBLY PROTEIN COX10 PRECURSOR - SACCHAROMYCES CEREVISIAE (BAKER'S YEAST), 462 aa.	1.70E-52	17

500	cg44017721	174	TTGGTAGGGACGG AACTCGGGCGCIG /TJGGCGGTGGCC GAGTGGAGATAGG A	G	T	Pro	Gln (729)	NON- CONSER VATIVE	cytochrome	Human Gene Similar to SPREMBL-ID:O00761 CYTOCHROME OXIDASE SUBUNIT VIA HEART ISOFORM PRECURSOR (EC 1.9.3.1) (CYTOCHROME-C OXIDASE) (CYTOCHROME A(3)) (CYTOCHROME AA(3)) - HOMO SAPIENS (HUMAN), 97 aa.	2.40E-32	22
501	cg41626024	279	GCTGGTTTGCTCC AGAGGCCAAG(A/ C)AGTCAGCCTACT GCCCTACAGTCA	A	C	Lys	Gln (730)	NON- CONSER VATIVE	deaminase	Human Gene Similar to SWISSPROT-ID:P32320 CYTIDINE DEAMINASE (EC 3.5.4.5) (CYTIDINE AMINOHYDROLASE) - HOMO SAPIENS (HUMAN), 146 aa. pcis:TREMBLNEW- ID:E1228801 CYTIDINE DEAMINASE (EC 3.5.4.5) - HOMO SAPIENS (HUMAN), 146 aa.	8.80E-78	1 (1p36.2)
502	cg43057018	1618	AAGCATCCGAACA ATCCTCATCTTTI/ GGAAGATGCCAG GAGCAATTCGAA T	T	G	End	Gly (731)	NON- CONSER VATIVE	dehydrogenase	Human Gene SWISSNEW- ID:P08319 ALCOHOL DEHYDROGENASE CLASS II PI CHAIN (EC 1.1.1.1) - HOMO SAPIENS (HUMAN), 391 aa. pcis:SWISSPROT-ID:P08319 ALCOHOL DEHYDROGENASE CLASS II PI CHAIN (EC 1.1.1.1) - HOMO SAPIENS (HUMAN), 391 aa.	1.30E-209	4 (4q22)

503	cg42837709	464	CCGCACCAACGCC GACATCATCGAG(A /G)CCCTGAGGAAG AAGGGCTTCAAGG G	A	G	Thr	Ala (732)	NON- CONSER VATIVE	dna_rna_bind	Human Gene Similar to TREMBL:NEW-ID:G913312 DNA BINDING PROTEIN MEF2 {CLONE XMEF2A1} - XENOPUS LAEVIS, 516 aa.	3.90E-86	1
504	cg43327954	2205	TCCACGACCGGGT AGAGAACTACAAI C/A)CCGCGGCAGC GCAAGCTCCGCAA CC	C	A	Asn	Lys (733)	NON- CONSER VATIVE	dna_rna_bind	Human Gene Similar to SPTRMBL-ID:Q61491 DNA- BINDING PROTEIN - MUS MUSCULUS (MOUSE), 546 aa.	5.50E-57	1
505	cg43971258	707	TCGTTGGAGATGA CAAGTCCGGAG(C /T)GAGCTCGGCTGT CTGGATGGGAAGG	C	T	Ala	Thr (734)	NON- CONSER VATIVE	dna_rna_bind_in hib	Human Gene Similar to SWISSNEW-ID:Q02535 DNA- BINDING PROTEIN INHIBITOR ID-3 (ID-LIKE PROTEIN INHIBITOR HLH 1R21) (HELIX- LOOP-HELIX PROTEIN HEIR-1) - HOMO SAPIENS (HUMAN), 119 aa. pcis:SWISSPROT- ID:Q02535 DNA-BINDING PROTEIN INHIBITOR ID-3 (ID- LIKE PROTEIN INHIBITOR HLH 1R21) (HELIX-LOOP- HELIX PROTEIN HEIR-1) - HOMO SAPIENS (HUMAN), 119 aa.	1.30E-60	1 (1p36.13)

506	cg41554010	1253	AGCTGGAGCAACA GCAGGAACAGCA[G/TTCAGGAGCAGC AGCAGGAGCAGGT GC	G	T	Gln	His (735)	NON- CONSER VATIVE	eph	Human Gene SWISSNEW- ID:P06727 APOLOPROTEIN A- IV PRECURSOR (APO-AIV) - HOMO SAPIENS (HUMAN), 396 aa.[pcis:SWISSPROT-ID:P06727 APOLOPROTEIN A-IV PRECURSOR (APO-AIV) - HOMO SAPIENS (HUMAN), 396 aa.	1.80E-203	11 (11q23)
507	cg43957743	1063	GTTTGGCATACTG GATATTTTAAT[CT]CAGTGGAGATAA AAGACAGCCCACT	C	T	Gly	Glu (736)	NON- CONSER VATIVE	esterase	Human Gene SWISSNEW- ID:Q15166 SERUM PARAOXONASE/ARYLESTERA SE 3 (EC 3.1.1.2) (EC 3.1.8.1) (PON 3) (SERUM ARYLDIAKYLPHOSPHATASE 3) (A-ESTERASE 3) (AROMATIC ESTERASE 3) - HOMO SAPIENS (HUMAN), 341 aa (fragment).[pcis:SWISSPROT- ID:Q15166 SERUM PARAOXONASE/ARYLESTERA SE 3 (EC 3.1.1.2) (EC 3.1.8.1) (PON 3) (SERUM ARYLDIAKYLPHOSPHATASE 3) (A-ESTERASE 3) (AROMATIC ESTERASE 3) - HOMO SAPIENS (HUMAN), 341 aa (fragment).	1.90E-178	

508	cg43957743	1079	TATTTTAATCCAGT GGAGATAAAAG(A/ C/CAGCCCACTAGG AAGTATATCAATA	A	C	Ser	Ala (737)	NON- CONSER VATIVE	esterase	Human Gene SWISSNEW- ID:Q15166 SERUM PARAOXONASE/ARYLESTERA SE 3 (EC 3.1.1.2) (EC 3.1.8.1) (PON 3) (SERUM ARYLDIARYLPHOSPHATASE 3) (A-ESTERASE 3) (AROMATIC ESTERASE 3) - HOMO SAPIENS (HUMAN), 341 aa (fragment), lpcis:SWISSPROT- ID:Q15166 SERUM PARAOXONASE/ARYLESTERA SE 3 (EC 3.1.1.2) (EC 3.1.8.1) (PON 3) (SERUM ARYLDIARYLPHOSPHATASE 3) (A-ESTERASE 3) (AROMATIC ESTERASE 3) - HOMO SAPIENS (HUMAN), 341 aa (fragment).	1.90E-178	
509	cg43248101	812	AAGTGAATTCTAT CTTGCAATGAAC(A/ G/JAGGAAGGAAAA CTCTATGCAAAAGA A	A	G	Lys	Glu (738)	NON- CONSER VATIVE	fgf	Human Gene Homologous to SWISSPROT-ID:P21781 KERATINOCYTE GROWTH FACTOR PRECURSOR (KGF) (FIBROBLAST GROWTH FACTOR-7) (FGF-7) (HBGF-7) - HOMO SAPIENS (HUMAN), 194 aa.	9.30E-106	15 (15q15)

510	cg43969014	332	GATGAGCTCTCCA ACCACGTAATTTTC/ ATGCGTTTTTGAT CCAGACCCAGATG	C	A	Arg	Ile (739)	NON- CONSER VATIVE	glucuronidase	Human Gene Similar to SWISSPROT-ID:P08236 BETA- GLUCURONIDASE PRECURSOR (EC 3.2.1.31) (BETA-G1) - HOMO SAPIENS (HUMAN), 651 aa.	7.40E-80	5
511	cg43286488	387	CACCAGCAAGATG CCCACGATCAGC/G /C/GAACCTGCCCA AGGCCTGCTCTTG	G	C	Pro	Arg (740)	NON- CONSER VATIVE	glycoprotein	Human Gene SWISSNEW- ID:P40967 MELANOCYTE PROTEIN PMEL 17 PRECURSOR (MELANOCYTE LINEAGE-SPECIFIC ANTIGEN GP100) (MELANOMA- ASSOCIATED ME20 ANTIGEN) (ME20M/ME20S) (ME20- M/ME20-S) (95 KD MELANOCYTE-SPECIFIC SECRETED GLYCOPROTEIN) - HOMO SAPIENS (HUMAN), 661 aa.lpcis:SWISSPROT-ID:P40967 MELANOCYTE PROTEIN PMEL 17 PRECURSOR (MELANOCYTE LINEAGE- SPECIFIC ANTIGEN GP100) (MELANOMA-ASSOCIATED ME20 ANTIGEN) (ME20M) (ME20-M / ME20-S) (95 KD MELANOCYTE-SPECIFIC SECRETED GLYCOPROTEIN) - HOMO SAPIENS (HUMAN), 661 aa.	0	12

512	cg44004239	663	TTTTCCCAGGGGT CACAGACTGAT/A/ GJACCCACAGAGGT CAGGGTCTTCTGT	A	G	Tyr	His (741)	NON- CONSER VATIVE	glycoprotein	Human Gene SWISSPROT- ID:Q12889 OVIDUCT-SPECIFIC GLYCOPROTEIN PRECURSOR (OVIDUCTAL GLYCOPROTEIN) (OVIDUCTIN) (ESTROGEN- DEPENDENT OVIDUCT PROTEIN) - HOMO SAPIENS (HUMAN), 678 aa.	0		
513	cg44004239	672	GGGGTCACAGACT GATAACCCACAG/A /GJGGTCAGGGTCT TCTGTCCAGTGGTC	A	G	Ser	Pro (742)	NON- CONSER VATIVE	glycoprotein	Human Gene SWISSPROT- ID:Q12889 OVIDUCT-SPECIFIC GLYCOPROTEIN PRECURSOR (OVIDUCTAL GLYCOPROTEIN) (OVIDUCTIN) (ESTROGEN- DEPENDENT OVIDUCT PROTEIN) - HOMO SAPIENS (HUMAN), 678 aa.	0		
514	cg44004239	773	CTGATGACCCACA GAAGTCATGGTC/A /GJTGGCCCCAGTG ATCTCAGTCTTCTC	A	G	Met	Thr (743)	NON- CONSER VATIVE	glycoprotein	Human Gene SWISSPROT- ID:Q12889 OVIDUCT-SPECIFIC GLYCOPROTEIN PRECURSOR (OVIDUCTAL GLYCOPROTEIN) (OVIDUCTIN) (ESTROGEN- DEPENDENT OVIDUCT PROTEIN) - HOMO SAPIENS (HUMAN), 678 aa.	0		

515	cg43932434	1504	ATATGTGTCATACT GGGAGGTGTG[G/ T]ATGTGAGGATGT ACACCCCTGTGTT	G	T	Ser	Tyr (744)	NON- CONSER VATIVE	glycoprotein	Human Gene SWISSPROT- ID:P16070 CD44 ANTIGEN PRECURSOR (PHAGOCYTTIC GLYCOPROTEIN 1) (PGP-1) (HUTCH-1) (EXTRACELLULAR MATRIX RECEPTOR-III) (ECMR-III) (GP90 LYMPHOCYTE HOMING/ADHESION RECEPTOR) (HERMES ANTIGEN) (HYALURONATE RECEPTOR) (HEPARAN SULFATE PROTEOGLYCAN) (EPICAN) (CDW44) - HOMO SAPIENS (HUMAN), 742 aa.	1.80E-195	11 (11pter)
516	cg40915005	622	AAGGAGCCTCTCT CCTTCCATGTCA[C/ T]CTGGATCGCATC CITTACAACCAT	C	T	Thr	Ile (745)	NON- CONSER VATIVE	glycoprotein	Human Gene SWISSNEW- ID:P06126 T-CELL SURFACE GLYCOPROTEIN CD1A PRECURSOR (CD1A ANTIGEN) (T-CELL SURFACE ANTIGEN T6/LEU-6) (HTA1 THYMOCYTE ANTIGEN) - HOMO SAPIENS (HUMAN), 327 aa.lpcis:SWISSPROT-ID:P06126 T-CELL SURFACE GLYCOPROTEIN CD1A PRECURSOR (CD1A ANTIGEN) (T-CELL SURFACE ANTIGEN T6/LEU-6) (HTA1 THYMOCYTE ANTIGEN) - HOMO SAPIENS (HUMAN), 327 aa.	2.00E-183	1 (1q21)

517	cg40913005	737	ATTCCAGCACCAT CGTTTCCTGTGIG/ CJCCTGTGTCAGG GGAAACTTCAGCA	G	C	Tyr	Cys (746)	NON- CONSER VATIVE	glycoprotein	Human Gene SWISSNEW- ID:P06126 T-CELL SURFACE GLYCOPROTEIN CD1A PRECURSOR (CD1A ANTIGEN) (T-CELL SURFACE ANTIGEN T6/LEU-6) (HTA1 THYMOCYTE ANTIGEN) - HOMO SAPIENS (HUMAN), 327 aa. pcis: SWISSPROT-ID:P06126 T-CELL SURFACE GLYCOPROTEIN CD1A PRECURSOR (CD1A ANTIGEN) (T-CELL SURFACE ANTIGEN T6/LEU-6) (HTA1 THYMOCYTE ANTIGEN) - HOMO SAPIENS (HUMAN), 327 aa.	2.00E-183	1 (1q21)
518	cg36834323	1529	GTGCTCCCTGATCC A TCGTGAAGCAT A/ GJTGGTAGCTCAAG TTATGTGGCATCT	A	G	Tyr	Cys (747)	NON- CONSER VATIVE	glycoprotein	Human Gene Similar to SWISSPROT-ID:P38159 HETEROGENEOUS NUCLEAR RIBONUCLEOPROTEIN G (HNRNP G) (GLYCOPROTEIN P43) - HOMO SAPIENS (HUMAN), 437 aa.	6.40E-91	
519	cg36834323	329	AATGCTGCGAAAG A ATATGAATGGA A /CJGTCITTTGCATGG AAAAGCAATAAAA	A	C	Lys	Thr (748)	NON- CONSER VATIVE	glycoprotein	Human Gene Similar to SWISSPROT-ID:P38159 HETEROGENEOUS NUCLEAR RIBONUCLEOPROTEIN G (HNRNP G) (GLYCOPROTEIN P43) - HOMO SAPIENS (HUMAN), 437 aa.	6.40E-91	

520	cg36834323	463	AAGTCGAGATCT GCAAGAGGAAGC[A/C]GTGGAGGAAC AAGAGGTGGCTT CC	A	C	Ser	Arg (749)	NON- CONSER VATIVE	glycoprotein	Human Gene Similar to SWISSPROT-ID:P38159 HETEROGENEOUS NUCLEAR RIBONUCLEOPROTEIN G (HNRNP G) (GLYCOPROTEIN P43) - HOMO SAPIENS (HUMAN), 437 aa.	6.40E-91	
521	cg44019290	1697	GCGGATAAGTAGA GGACCTTCATGT[G]GTATTGCTGGT GAAGTTGGTTCCG	T	G	Asn	His (750)	NON- CONSER VATIVE	glycoprotein	Human Gene Similar to SWISSPROT-ID:P04216 THY-1 MEMBRANE GLYCOPROTEIN PRECURSOR (THY-1 ANTIGEN) (CDW90) (CD90 ANTIGEN) - HOMO SAPIENS (HUMAN), 161 aa.	2.50E-80	11
522	cg42336656	1665	CTTAGACATACAA TATACTTACCTT[A/ G]GAGGTCACGTAT GTTTGTCGCACA	A	G	Arg	Gly (751)	NON- CONSER VATIVE	glycoprotein	Human Gene Similar to SWISSPROT-ID:Q05910 CELL SURFACE ANTIGEN MS2 PRECURSOR (EC 3.4.24.-) (MACROPHAGE CYSTEINE- RICH GLYCOPROTEIN) (CD156 ANTIGEN) - MUS MUSCULUS (MOUSE), 826 aa.	9.40E-38	
523	cg42730678	980	GGAGCGAGCGTGG ATCCAGTTCGCG[G /T]CGGGGTGTGTTG GGTCAAGTTGCTG	G	T	Ala	Asp (752)	NON- CONSER VATIVE	homeobox	Human Gene SWISSPROT- ID:P28356 HOMEBOX PROTEIN HOX-D9 (HOX-4C) (HOX-5.2) - HOMO SAPIENS (HUMAN), 342 aa.	2.60E-188	2

524	cg42714160	769	GCCCTGTGCCTGA CGGAGAGGCAGAI T/GJCAAGATATGG TTCCAGAACCGAC GC	T	G	Ile	Ser (753)	NON- CONSER VATIVE	homeobox	Human Gene Homologous to SWISSPROT-ID:P17509 HOMEBOX PROTEIN HOX-B6 (HOX-2B) (HOX-2.2) (HU-2) - HOMO SAPIENS (HUMAN), 224 aa.	1.10E-123	
525	cg42359655	3297	CTGGGCAACCATAT AGGATAGCCACIA /GJCCGTCATCAAA GCCCATGCCAGAG T	A	G	Thr	Ala (754)	NON- CONSER VATIVE	hydrolase	Human Gene SWISSPROT- ID:P09848 LACTASE- PHLORIZIN HYDROLASE PRECURSOR (EC 3.2.1.108) (EC 3.2.1.62) (LACTASE- GLYCOSYL CERAMIDASE) - HOMO SAPIENS (HUMAN), 1927 aa.	0	2 (2q21)
526	cg43925670	2172	GTGGAGGGTGCAG GTGAAGTAGCATJC /GJCACTTCCTCTT CCTCTTCTTGAT	C	G	Asp	His (755)	NON- CONSER VATIVE	interferon	Human Gene SWISSPROT- ID:Q16666 GAMMA- INTERFERON-INDUCIBLE PROTEIN IFI-16 (INTERFERON- INDUCIBLE MYELOID DIFFERENTIATION TRANSCRIPTIONAL ACTIVATOR) - HOMO SAPIENS (HUMAN), 729 aa. pels: SPTREMBL-ID:Q16666 IFI16=INTERFERON- INDUCIBLE MYELOID DIFFERENTIATION TRANSCRIPTIONAL ACTIVATOR - HOMO SAPIENS (HUMAN), 729 aa (fragment).	0	1

527	cg43090990	1083	TGCTCCATCAAAA ATGAAGCAAGGC[CT]GCCATGTTAC CGACACCGGAAAA A	C	T	Pro	Leu (756)	NON- CONSER VATIVE	kinase	Human Gene SWISSPROT- ID:Q04759 PROTEIN KINASE C, THETA TYPE (EC 2.7.1.-) (NPKC-THETA) - HOMO SAPIENS (HUMAN), 706 aa.	0	10
528	cg43969763	2663	CAAAAGCAAGAAA GTTCTTTGAGAA[G/ T]TGCCAGATGGC ACTTGGAACTTAA	G	T	Lys	Asn (757)	NON- CONSER VATIVE	kinase	Human Gene SWISSPROT- ID:Q13627 SERINE/THREONINE-SPECIFIC PROTEIN KINASE MINIBRAIN HOMOLOG (EC 2.7.1.-) (HP86) (DYRK) - HOMO SAPIENS (HUMAN), 763 aa.	0	21 (21q22.1)
529	cg43932396	1226	AGTCCACCGCGC CTCAGGCCGTC[C /T]GCTGGCCGAGT AGGAGAACTGGGG G	C	T	Gly	Ser (758)	NON- CONSER VATIVE	kinase	Human Gene SWISSPROT- ID:P31749 RAC-ALPHA SERINE/THREONINE KINASE (EC 2.7.1.-) (RAC-PK-ALPHA) (PROTEIN KINASE B) (PKB) (C- AKT) - HOMO SAPIENS (HUMAN), 480 aa.	1.40E-262	14 (14q32.3)
530	cg43917871	1429	GGCACTGAAGAAA TCCCTGACATCA[T/ C]ATTGGCGTGTCT GACGGGCGTACTG	T	C	Met	Val (759)	NON- CONSER VATIVE	kinase	Human Gene SWISSPROT- ID:P19138 CASEIN KINASE II, ALPHA CHAIN (CK II) (EC 2.7.1.37) - HOMO SAPIENS (HUMAN), AND BOS TAURUS (BOVINE), 391 aa.	2.00E-215	11 (20p13)
531	cg43917871	1621	GGGCTGACAAAGGT GCTGATTTTCAC[T/ G]GTGGACAAAGC GTTCCCATCGCTTT	T	G	Ser	Arg (760)	NON- CONSER VATIVE	kinase	Human Gene SWISSPROT- ID:P19138 CASEIN KINASE II, ALPHA CHAIN (CK II) (EC 2.7.1.37) - HOMO SAPIENS (HUMAN), AND BOS TAURUS (BOVINE), 391 aa.	2.00E-215	11 (20p13)

532	cg43917871	1713	TTCAATGTTGTATT TGTCATATAGT/C JCATATAAATCTTC TGTCGCCAGAAC	T	C	Asp	Gly (761)	NON- CONSER VATIVE	kinase	Human Gene SWISSPROT- ID:P19138 CASEIN KINASE II, ALPHA CHAIN (CK II) (EC 2.7.1.37) - HOMO SAPIENS (HUMAN), AND BOS TAURUS (BOVINE), 391 aa.	2.00E-215	11 (20p13)
533	cg43917871	2096	TGTAAATCGAAT ATCATAGTCTGT/T GJAACGTCTGGTAC AATTGCTTGAAGT	T	G	Leu	Phe (762)	NON- CONSER VATIVE	kinase	Human Gene SWISSPROT- ID:P19138 CASEIN KINASE II, ALPHA CHAIN (CK II) (EC 2.7.1.37) - HOMO SAPIENS (HUMAN), AND BOS TAURUS (BOVINE), 391 aa.	2.00E-215	11 (20p13)
534	cg43322545	1107	TGGGCTAGGCAGC CTCCATCCTCACIA/ CJCCCTTATCACA TCCGCGTGGCATG	A	C	Thr	Pro (763)	NON- CONSER VATIVE	kinasereceptor	Human Gene SWISSNEW- ID:P30530 TYROSINE-PROTEIN KINASE RECEPTOR UFO PRECURSOR (EC 2.7.1.112) (AXL ONCOGENE) - HOMO SAPIENS (HUMAN), 887 aa.lpcis:SWISSPROT-ID:P30530 TYROSINE-PROTEIN KINASE RECEPTOR UFO PRECURSOR (EC 2.7.1.112) (AXL ONCOGENE) - HOMO SAPIENS (HUMAN), 887 aa.	0	19 (19q13.1)

535	cg43322545	2116	TTCCTCCTCTATTC CCGGCTCGGGGTA/ GJCCAGCCAGTGTA CCTGCCCACTCAG	A	G	Asp	Gly (764)	NON- CONSER VATIVE	kinasereceptor	Human Gene SWISSNEW- ID:P30530 TYROSINE-PROTEIN KINASE RECEPTOR UFO PRECURSOR (EC 2.7.1.112) (AXL ONCOGENE) - HOMO SAPIENS (HUMAN), 887 aa.lpcIs:SWISSPROT-ID:P30530 TYROSINE-PROTEIN KINASE RECEPTOR UFO PRECURSOR (EC 2.7.1.112) (AXL ONCOGENE) - HOMO SAPIENS (HUMAN), 887 aa.	0	19 (19q13.1)
-----	------------	------	---	---	---	-----	--------------	--------------------------	----------------	--	---	---------------------

536	cg43958558	863	TCCGGGATAAGCT CCAGGTGCTCCA[G /TJGGTAGGCGCCT GGAGGTGCCCTGTC C	G	T	Pro	His (765)	NON- CONSER VATIVE	laminin	Human Gene Homologous to SWISSPROT-ID:P17931 GALECTIN-3 (GALACTOSE- SPECIFIC LECTIN 3) (MAC-2 ANTIGEN) (IGE-BINDING PROTEIN) (35 KD LECTIN) (CARBOHYDRATE BINDING PROTEIN 35) (CBP 35) (LAMININ-BINDING PROTEIN) (LECTIN L-29) (L-31) (GALACTOSIDE-BINDING PROTEIN) (GALBP) - HOMO SAPIENS (HUMAN), 249 aa pcsl:SWISSPROT-ID:P17931 GALECTIN-3 (GALACTOSE- SPECIFIC LECTIN 3) (MAC-2 ANTIGEN) (IGE-BINDING PROTEIN) (35 KD LECTIN) (CARBOHYDRATE BINDING PROTEIN 35) (CBP 35) (LAMININ-BINDING PROTEIN) (LECTIN L-29) (L-31) (GALACTOSIDE-BINDING PROTEIN) (GALBP) - HOMO SAPIENS (HUMAN), 249 aa.	3.90E-139	14 (14q21)
537	cg43966144	718	AAGCTTGTCATGC CTCACAGCAGTG[C /A]GCACAAGACTG CCCAGCCCAATGG A	C	A	Ala	Glu (766)	NON- CONSER VATIVE	MHC	Human Gene Homologous to SWISSPROT-ID:P28068 CLASS II HISTOCOMPATIBILITY ANTIGEN, M BETA CHAIN PRECURSOR - HOMO SAPIENS (HUMAN), 263 aa.	9.10E-147	6 (6p21.3)

538	cg43966144	823	ACTTACACCTGTGT GGTAGAGCACA/T C/TGGGGCTCCTGA GCCCATCCTTCGG	T	C	Ile	Thr (767)	NON- CONSER VATIVE	MHC	Human Gene Homologous to SWISSPROT-ID:P28068 CLASS II HISTOCOMPATIBILITY ANTIGEN, M BETA CHAIN PRECURSOR - HOMO SAPIENS (HUMAN), 263 aa.	9.10E-147	6 (6p21.3)
539	cg42686658	907	GGCCTGGTGGGCT TCCTCGTGGGCA/C TJCGCTCATCAT CATGGGCACATAT	C	T	Thr	Ile (768)	NON- CONSER VATIVE	MHC	Human Gene Homologous to SWISSPROT-ID:P06340 HLA CLASS II HISTOCOMPATIBILITY ANTIGEN, DZ ALPHA CHAIN PRECURSOR (MHC DN-ALPHA) - HOMO SAPIENS (HUMAN), 250 aa.	3.70E-134	6 (6p21.3)
540	cg38337333	1044	CTGAGCCGAGAGC GTTGTCTCTGCTG/C G/CATGAGCACCAC AGTCAGGCGCTTGA	C	G	Pro	Ala (769)	NON- CONSER VATIVE	MHC	Human Gene Homologous to SPTREMBL-ID:Q95368 HLA CLASS I INHIBITORY NK RECEPTOR - HOMO SAPIENS (HUMAN), 455 aa.	1.80E-113	19
541	cg38337333	424	AGCCCGGCGGGC CCCACGGTTCGC/A /G/CAGGAGAGAAC GTGACCTTGTCTG	A	G	Thr	Ala (770)	NON- CONSER VATIVE	MHC	Human Gene Homologous to SPTREMBL-ID:Q95368 HLA CLASS I INHIBITORY NK RECEPTOR - HOMO SAPIENS (HUMAN), 455 aa.	1.80E-113	19
542	cg42481172	340	CCGGCTGTGCTCA GGGGTGTGGGT/A /G/CGGATACAGAG GAGCGGCTGGTGG A	A	G	Thr	Ala (771)	NON- CONSER VATIVE	misc_channel	Human Gene Similar to SPTREMBL-ID:P91197 SIMILAR TO LIGAND-GATED IONIC CHANNEL PROTEIN - CAENORHABDITIS ELEGANS, 461 aa.	2.30E-71	1

543	cg3000465	238	GAAGATGCCCTCC TCAGACATGAGTGG /TJGAAAGGTTATC AGAAATGGGTCCG C	G	T	Trp	Leu (772)	NON- CONSER VATIVE	misc_channel	Human Gene Similar to SPTREMBL-ID:P91197 SIMILAR TO LIGAND-GATED IONIC CHANNEL PROTEIN - CAENORHABDITIS ELEGANS, 461 aa.	6.10E-70	8 (8p11.2)
544	cg3000465	240	AGATGCCCTCCTC AGACATGAGTGGI A/CJAAGGTTATCA GAAATGGGTCCG CC	A	C	Lys	Gln (773)	NON- CONSER VATIVE	misc_channel	Human Gene Similar to SPTREMBL-ID:P91197 SIMILAR TO LIGAND-GATED IONIC CHANNEL PROTEIN - CAENORHABDITIS ELEGANS, 461 aa.	6.10E-70	8 (8p11.2)
545	cg43249083	1067	GCCTCGGGCTTCC ACTACGGTGTGC[A /TJCGCCTGCGAGG GCTGCAAGGGGCTT T	A	T	His	Leu (774)	NON- CONSER VATIVE	nucl_recpt	Human Gene SWISSPROT- ID:P20393 V-ERBA RELATED PROTEIN EAR-1 - HOMO SAPIENS (HUMAN), 614 aa.	0	17 (17q11.2)
546	cg44928796	68	AGCGGACGGTCC GGAGCAAGCCCAI G/CJAGGCAGAGGA GGCGACAGAGGGA AA	G	C	Gln	His (775)	NON- CONSER VATIVE	nucl_recpt	Human Gene SWISSNEW- ID:P10275 ANDROGEN RECEPTOR - HOMO SAPIENS (HUMAN), 919 aa. lpcis:SWISSPROT-ID:P10275 ANDROGEN RECEPTOR - HOMO SAPIENS (HUMAN), 919 aa.	0	X (Xq11)
547	cg43323772	91	GTGCCGGGAGTGA GCGATGAGCTGG[C /TJTTCTGTCTCTGG CCCACAGAGTCCG	C	T	Leu	Phe (776)	NON- CONSER VATIVE	nuclease	Human Gene TREMBLNEW- ID:G2935442 RIBONUCLEASE H1 - HOMO SAPIENS (HUMAN), 286 aa. lpcis:TREMBLNEW- ID:G2935444 RIBONUCLEASE H1 - HOMO SAPIENS (HUMAN), 286 aa.	1.40E-157	

548	cg42732993	809	GGCTATAATCACA ATGGGAATGGTIG /TJTGAAAGCCAAA CCAAAAATGGCCA A	G	T	Cys	Phe (777)	NON- CONSER VATIVE	oncogene	Human Gene Homologous to SPTREMBL-ID:Q13692 BCR/ABL FUSION PROTEIN - HOMO SAPIENS (HUMAN), 284 aa (fragment).	6.00E-150	
549	cg42904626	155	ATATAAACTTGTG GTAGTTGGAGCTIG /TJGTGGCGTAGGC AAGAGTGCCTTGA C	G	T	Gly	Cys (778)	NON- CONSER VATIVE	oncogene	Human Gene Similar to SWISSPROT-ID:P01118 TRANSFORMING PROTEIN P21/K-RAS 2B - HOMO SAPIENS (HUMAN), 188 aa.	1.10E-97	12
550	cg42904626	304	TGGATAITCTCGAC ACAGCAGGTCAJ/ CJGAGGAGTACAGT GCAATGAGGGACC	A	C	Gln	His (779)	NON- CONSER VATIVE	oncogene	Human Gene Similar to SWISSPROT-ID:P01118 TRANSFORMING PROTEIN P21/K-RAS 2B - HOMO SAPIENS (HUMAN), 188 aa.	1.10E-97	12
551	cg42691989	706	CTGTTTCAGGATCTC CTCAITCTGACIA/T JGTTCTCCTGATGT CCAAATTGGTTG	A	T	Cys	Ser (780)	NON- CONSER VATIVE	peroxidase	Human Gene Homologous to SWISSPROT-ID:P18283 GLUTATHIONE PEROXIDASE- GASTROINTESTINAL (EC 1.11.1.9) (GSHPX-GI) (GLUTATHIONE PEROXIDASE- RELATED PROTEIN 2) (GPRP) - HOMO SAPIENS (HUMAN), 190 aa.	8.90E-101	14 (14q24.1)

552	cg43917453	4096	AGTCTCTCGCGGA GCTGGTCCGGGIA /GJCCGGGAGGGT AGGTCAGCGCAGA C	A	G	Ser	Pro (781)	NON- CONSER VATIVE	phosphatase	Human Gene TREMBLNEW- ID:Q2262075 IAR/RECEPTOR- LIKE PROTEIN-TYROSINE PHOSPHATASE PRECURSOR - HOMO SAPIENS (HUMAN), 1015 aa.	0	7
553	cg43947363	368	CTGGCGCACTACT CGGACCTGCTCCIC/ TJCTGGCGGGCCT GGGCTGATTGAG	C	T	Gly	Glu (782)	NON- CONSER VATIVE	phosphatase	Human Gene SWISSPROT- ID:P23469 PROTEIN-TYROSINE PHOSPHATASE EPSILON PRECURSOR (EC 3.1.3.48) (R- PTP- EPSILON) - HOMO SAPIENS (HUMAN), 700 aa.	0	
554	cg43928335	3187	GCACAAGGAACGG AAITGCTGTCTGJA/ GJTTCGTCTTAA CAGCAITTTGATGC	A	G	Ile	Thr (783)	NON- CONSER VATIVE	phosphatase	Human Gene SWISSPROT- ID:P54613 PROTEIN PHOSPHATASE PP2A, 65 KD REGULATORY SUBUNIT, BETA ISOFORM (PROTEIN PHOSPHATASE PP2A SUBUNIT A, BETA ISOFORM) (P65-BETA) - SUS SCROFA (FIG), 602 aa (fragment).	3.20E-302	11 (11q22)
555	cg43996195	1330	CTTCGGGGAAGT TGGGAITTCACIC/ TJTAGTCAAAGAT CTGGGCTGAGTT	C	T	Gly	Ser (784)	NON- CONSER VATIVE	phosphorylase	Human Gene SWISSPROT- ID:P00491 PURINE NUCLEOSIDE PHOSPHORYLASE (EC 2.4.2.1) (INOSINE PHOSPHORYLASE) (PNP) - HOMO SAPIENS (HUMAN), 289 aa.	2.40E-155	

556	cg44022214	3340	AGGTCCTCCTCGA ATTGGGATGGCC[A /G]AGGTGCATCAT CATCATCCCAGAG G	A	G	Trp	Arg (785)	NON- CONSER VATIVE	polymerase	Human Gene SWISSNEW- ID:P28340 DNA POLYMERASE DELTA CATALYTIC CHAIN (EC 2.7.7.7) - HOMO SAPIENS (HUMAN), 1107 aa.lpcis:SWISSPROT-ID:P28340 DNA POLYMERASE DELTA CATALYTIC CHAIN (EC 2.7.7.7) - HOMO SAPIENS (HUMAN), 1107 aa.	0	19 (19q13.3)
557	cg43958858	1593	CTCAGACCATGTC CTTCGGATGCAC[C/ G]TTACAGAGCAC CTGGGGAGCAGGA	C	G	Arg	Gly (786)	NON- CONSER VATIVE	polymerase	Human Gene SWISSNEW- ID:P25205 DNA REPLICATION LICENSING FACTOR MCM3 (DNA POLYMERASE ALPHA HOLOENZYME-ASSOCIATED PROTEIN P1) (RLF BETA SUBUNIT) (P102 PROTEIN) - HOMO SAPIENS (HUMAN), 808 aa.lpcis:SWISSPROT-ID:P25205 DNA REPLICATION LICENSING FACTOR MCM3 (DNA POLYMERASE ALPHA HOLOENZYME-ASSOCIATED PROTEIN P1) (RLF BETA SUBUNIT) (P102 PROTEIN) - HOMO SAPIENS (HUMAN), 808 aa.	0	6 (6p12)

558	cg42534568	641	CGCTTTGAGACGC AGCTGGGCACCC[A /T]GGCGCAGTCCC CAACACACTCCTG	A	T	Gln	Leu (787)	NON- CONSER VATIVE	potassium_chan nel	Human Gene SWISSPROT- ID:P22460 VOLTAGE-GATED POTASSIUM CHANNEL PROTEIN KV1.5 (HK2) (HPCN1) - HOMO SAPIENS (HUMAN), 613 aa.	0	12 (12p13)
559	cg42534568	868	GGGGACGAGGCC ATGGAGCGCTTC[C /G]GCGAGGATGAG GGCTTCATTAAAG A	C	G	Arg	Gly (788)	NON- CONSER VATIVE	potassium_chan nel	Human Gene SWISSPROT- ID:P22460 VOLTAGE-GATED POTASSIUM CHANNEL PROTEIN KV1.5 (HK2) (HPCN1) - HOMO SAPIENS (HUMAN), 613 aa.	0	12 (12p13)
560	cg42534568	910	CATTAAAGAAGAG GAGAAAGCCCTG[C /G]CCCGCAACGAG TTCCAGGCCCAGG T	C	G	Pro	Ala (789)	NON- CONSER VATIVE	potassium_chan nel	Human Gene SWISSPROT- ID:P22460 VOLTAGE-GATED POTASSIUM CHANNEL PROTEIN KV1.5 (HK2) (HPCN1) - HOMO SAPIENS (HUMAN), 613 aa.	0	12 (12p13)
561	cg43134190	898	TGGAGGGGATGCT CATTTTGATGAAIG/ CJATGAAAGGTGG ACCAACAATTCA G	G	C	Asp	His (790)	NON- CONSER VATIVE	protease	Human Gene Similar to SWISSPROT-ID:P50280 MATRILYSIN PRECURSOR (EC 3.4.24.23) (PUMP-1 PROTEASE) (UTERINE METALLOPROTEINASE) (MATRIX METALLOPROTEINASE-7) (MMP-7) (MATRIN) - RATTUS NORVEGICUS (RAT), 267 aa.	2,40E-59	11 (11q22)

562	cg43154190	923	GATGAAAGGTGGA CCAAACAATTCA[G /CIAGAGTACAAC TACATCGTGTTCG	G	C	Arg	Thr (791)	NON- CONSER VATIVE	protease	Human Gene Similar to SWISSPROT-ID:P50280 MATRILYSIN PRECURSOR (EC 3.4.24.23) (PUMP-1 PROTEASE) (UTERINE METALLOPROTEINASE) (MATRIX METALLOPROTEINASE-7) (MMP-7) (MATRIN) - RATTUS NORVEGICUS (RAT), 267 aa.	2.40E-59	11 (11q22)
563	cg43927549	694	ATTCTACGATTCCG GTTTGCTCCAG[G/T JGTAACTAGCGCT CCTTCCGTAAAC	G	T	Gly	Cys (792)	NON- CONSER VATIVE	reductase	Human Gene Homologous to SWISSPROT-ID:P16083 NAD(P)H DEHYDROGENASE (QUINONE) 2 (EC 1.6.99.2) (QUINONE REDUCTASE) (DT- DIAPHORASE) (AZOREDUCTASE) (PHYLLOQUINONE REDUCTASE) (MENADIONE REDUCTASE) - HOMO SAPIENS (HUMAN), 231 aa.	1.60E-124	6 (6pter)
564	cg43325541	1081	CGCTGCCTTCTCC GAAAGGTCTGC[C/ TJCCTTACACGCTT CGGCTTCCCGCAG	C	T	Gly	Glu (793)	NON- CONSER VATIVE	synthase	Human Gene TREMBLNEW- ID:G2725625 ACETOLACTATE SYNTHASE - HOMO SAPIENS (HUMAN), 632 aa.	0	19

565	cg43064068	1474	GTGAAGGCATTG TGGTCCTGGCCTIC/ TIGCAGTTCCTGTC CCATGACCCAGAA	C	T	Ser	Leu (794)	NON- CONSER VATIVE	synthase	Human Gene Similar to SWISSNEW-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE--COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa.lpcis:SWISSPROT-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE--COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa.	7.40E-65	
566	cg43064068	1617	GACTGTCACAGGG AAAATTCAACGA[G /AJCCAAGCTTCGA GACAAGGAGTGGA A	G	A	Ala	Thr (795)	NON- CONSER VATIVE	synthase	Human Gene Similar to SWISSNEW-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE--COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa.lpcis:SWISSPROT-ID:P39062 ACETYL-COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE--COA LIGASE) (ACYL- ACTIVATING ENZYME) (ACETYL-COA SYNTHASE) - BACILLUS SUBTILIS, 572 aa.	7.40E-65	

567	cg36988276	1119	GCAAGAAAGTTGAT TATATGACTCAG(A /G)CTAGGGTCAG AGATCCTCTCTGGC	A	G	Thr	Ala (796)	NON- CONSER VATIVE	tm7	Human Gene SWISSPROT- ID:P23945 FOLLICLE STIMULATING HORMONE RECEPTOR PRECURSOR (FSH- R) (FOLLITROPIN RECEPTOR) - HOMO SAPIENS (HUMAN), 695 aa.	0	2 (2p21)
568	cg36988276	535	AAGGCCAACCAACC TGCTCTACATCA(A/ C)CCCTGAGGCCCTT CCAGAACCTTCCC	A	C	Asn	Thr (797)	NON- CONSER VATIVE	tm7	Human Gene SWISSPROT- ID:P23945 FOLLICLE STIMULATING HORMONE RECEPTOR PRECURSOR (FSH- R) (FOLLITROPIN RECEPTOR) - HOMO SAPIENS (HUMAN), 695 aa.	0	2 (2p21)
569	cg32296848	1475	GAATGCTTTGAGA ATCCAGTGTCTC(G/ T)GCAGAAAGCAGT CTTCCAAACATGC	C	T	Arg	Cys (798)	NON- CONSER VATIVE	tm7	Human Gene SWISSPROT- ID:P35348 ALPHA-1A ADRENERGIC RECEPTOR (ALPHA 1A-ADRENOCEPTOR) (ALPHA-1C ADRENERGIC RECEPTOR) - HOMO SAPIENS (HUMAN), 466 aa.	1.60E-252	8 (8p21)
570	cg2524739	1590	TCTCTCTGGAGAA GATCCCAACCCATTC /G)ACACAAAACGG TCAGCACCCCAACC T	C	G	Ile	Met (799)	NON- CONSER VATIVE	tm7	Human Gene SWISSPROT- ID:P21728 D(1A) DOPAMINE RECEPTOR - HOMO SAPIENS (HUMAN), 446 aa.	8.30E-240	5 (5q35.1)

571	cg2320320	394	AGTGTCTGGATGA TCITTGTGGTCA(C/ TTTGCATCCGTTT CACAAATGGGCTT	C	T	Thr	Ile (800)	NON- CONSER VATIVE	tm7	Human Gene SWISSPROT- ID:P04001 GREEN-SENSITIVE OPNIN (GREEN CONE PHOTORECEPTOR PIGMENT) - HOMO SAPIENS (HUMAN), 364 aa.	8.50E-199	
572	cg43264978	519	CATCTTCTCCATCA ACCTCTTCAGCA/G JGCATTTTCTTCCTC ACGTGCATGAG	A	G	Ser	Gly (801)	NON- CONSER VATIVE	tm7	Human Gene TREMBLNEW- ID:G2736282 G PROTEIN COUPLED RECEPTOR - HOMO SAPIENS (HUMAN), 362 aa.	1.40E-196	
573	cg3003708	285	TCITTGTGGACAT CTGCTTCTCCTTT/C] CAACACCGTCCCC AAGATGCTGGCC	T	C	Phe	Ser (802)	NON- CONSER VATIVE	tm7	Human Gene TREMBLNEW- ID:E1246031 OLFACTORY RECEPTOR - HOMO SAPIENS (HUMAN), 312 aa.	2.50E-160	
574	cg38841806	68	GGCCCTGAGAGCA ACACCACGGGCAIT /C]ACAGCCTTCTC CATGCCCAAGCTGG	T	C	Ile	Thr (803)	NON- CONSER VATIVE	tm7	Human Gene Similar to SWISSPROT-ID:P30975 TACHYKININ-LIKE PEPTIDES RECEPTOR 99D (DKR) - DROSOPHILA MELANOGASTER (FRUIT FLY), 519 aa.	2.10E-67	
575	cg43336100	688	TGGAAGCGTGCA CCAGTGAGACCAJA /TTJGAGGCTTGAGT CTTTAGTGCTG	A	T	Met	Leu (804)	NON- CONSER VATIVE	tmf	Human Gene SWISSPROT- ID:P26022 PENTAXIN- RELATED PROTEIN PTX3 PRECURSOR (TUMOR NECROSIS FACTOR- INDUCIBLE PROTEIN TSG-14) - HOMO SAPIENS (HUMAN), 381 aa.	2.20E-207	3 (3q25)

576	cg43335562	234	GAGGCGCGGGGAG CCAGGCCTGGGCT /CJCCGGGTCCCA AGACCCCTTGCTC	T	C	Leu	Pro (805)	NON- CONSER VATIVE	tnfrceptor	Human Gene Similar to TREMBLNEW-ID:G2653845 TNF RECEPTOR-RELATED RECEPTOR FOR TRAIL - HOMO SAPIENS (HUMAN), 386 aa.	2.30E-55	8
577	cg43140548	2857	ACTCGCACGTGGA TCCTGAGGCTGTIA/ GJAGAGGTAAGGA AGGCTTTGCCACA G	A	G	Tyr	His (806)	NON- CONSER VATIVE	transcriptfactor	Human Gene SPTREMBL- ID:Q14872 METAL- REGULATORY TRANSCRIPTION FACTOR - HOMO SAPIENS (HUMAN), 753 aa.	0	1
578	cg43011561	1285	CATTGACAGCGAG GCCTCCTCAGCC[C/ TJCTTCATGGCGA AGAAGAAGACGCC	C	T	Leu	Phe (807)	NON- CONSER VATIVE	transcriptfactor	Human Gene SWISSPROT- ID:P35269 TRANSCRIPTION INITIATION FACTOR IIF, ALPHA SUBUNIT (TIF- ALPHA) (TRANSCRIPTION INITIATION FACTOR RAP74) - HOMO SAPIENS (HUMAN), 517 aa.	4.30E-275	19 (19p13.3)
579	cg43998970	1346	TGACAGAGCTGTA CCGTGACATTTTC/ GJCAGCACCTTCGG GATGAATCAGGCA	C	G	Phe	Leu (808)	NON- CONSER VATIVE	transcriptfactor	Human Gene SPTREMBL- ID:Q07279 TRANSCRIPTION FACTOR NF-E2 - MUS MUSCULUS (MOUSE), 373 aa.	1.70E-177	12

580	cg2337639	464	CACTACTATGTCTT CACCGACCAAGC/C/ TJGGCCCGCGGTGCC CCGCGTGACCGCTG	T	Pro	Leu (809)	NON- CONSER VATIVE	transferase	Human Gene SWISSPROT- ID:P16442 FUCOSYLGLYCOPROTEIN ALPHA-N- ACETYL GALACTOSAMINYLT RANSFERASE (EC 2.4.1.40) (HISTO-BLOOD GROUP A TRANSFERASE) (A TRANSFERASE) / FUCOSYLGLYCOPROTEIN 3- ALPHA- GALACTOSYLTRANSFERASE (EC 2.4.1.37) (HISTO-BLOOD GROUP B TRANSFERASE) (B TRANSFERASE) (NAGAT) - HOMO SAPIENS (HUMAN), 354 aa.	6.50E-192	9 (9q34)
-----	-----------	-----	--	---	-----	--------------	--------------------------	-------------	--	-----------	----------

381	cg2537639	523	TCGGCAGCTGTCA GTGCTGGAGGTG[C /GJGCGCCTACAAG CGCTGGCAGGACG T	C	G	Arg	Gly (810)	NON- CONSER VATIVE	transferase	Human Gene SWISSPROT- ID:P16442 FUCOSYLGLYCOPROTEIN ALPHA-N- ACETYLGALACTOSAMINYLT RANSFERASE (EC 2.4.1.40) (HISTO-BLOOD GROUP A TRANSFERASE) (A TRANSFERASE) / FUCOSYLGLYCOPROTEIN 3- ALPHA- GALACTOSYLTRANSFERASE (EC 2.4.1.37) (HISTO-BLOOD GROUP B TRANSFERASE) (B TRANSFERASE) (NAGAT) - HOMO SAPIENS (HUMAN), 354 aa.	6.50E-192	9 (9q34)
-----	-----------	-----	---	---	---	-----	--------------	--------------------------	-------------	---	-----------	----------

582	cg2537639	643	GGTGTGCCGTGGAC GTGGACATGGAGTT /AJTCCGGGACCAC GTGGGCGTGGAGA T	A	Phe	Ile (811)	NON- CONSER VATIVE	transferase	Human Gene SWISSPROT- ID:P16442 FUCOSYLGLYCOPROTEIN ALPHA-N- ACETYLGLALACTOSAMINYLT RANSFERASE (EC 2.4.1.40) (HISTO-BLOOD GROUP A TRANSFERASE) (A TRANSFERASE) / FUCOSYLGLYCOPROTEIN 3- ALPHA- GALACTOSYLTRANSFERASE (EC 2.4.1.37) (HISTO-BLOOD GROUP B TRANSFERASE) (B TRANSFERASE) (NAGAT) - HOMO SAPIENS (HUMAN), 354 aa.	6.50E-192	9 (934)
-----	-----------	-----	--	---	-----	--------------	--------------------------	-------------	--	-----------	---------

583	cg2537639	700	TCCGCTGTTCGGCA CCCTGCACCCC[G/ AJGCTTCTACGGAA GCAGCCGGGAGGC	G	A	Gly	Ser (812)	NON- CONSER VATIVE	transfèrase	Human Gene SWISSPROT- ID:P16442 FUCOSYLGLYCOPROTEIN ALPHA-N- ACETYL GALACTOSAMINYLT RANSFERASE (EC 2.4.1.40) (HISTO-BLOOD GROUP A TRANSFERASE) (A TRANSFERASE) / FUCOSYLGLYCOPROTEIN 3- ALPHA- GALACTOSYLTRANSFERASE (EC 2.4.1.37) (HISTO-BLOOD GROUP B TRANSFERASE) (B TRANSFERASE) (NAGAT) - HOMO SAPIENS (HUMAN), 354 aa.	6.50E-192	9 (9q34)
-----	-----------	-----	---	---	---	-----	--------------	--------------------------	-------------	--	-----------	----------

384	cg2537639	793	CAAGGACGAGGGC GATTCTACTAC[C/ AJTGGGGGGTTCT TCGGGGGGTTCGGT	C	A	Leu	Met (813)	NON- CONSER VATIVE	transferase	Human Gene SWISSPROT- ID:P16442 FUCOSYLGLYCOPROTEIN ALPHA-N- ACETYLGALACTOSAMINYLT RANSFERASE (EC 2.4.1.40) (HISTO-BLOOD GROUP A TRANSFERASE) (A TRANSFERASE) / FUCOSYLGLYCOPROTEIN 3- ALPHA- GALACTOSYLTRANSFERASE (EC 2.4.1.37) (HISTO-BLOOD GROUP B TRANSFERASE) (B TRANSFERASE) (NAGAT) - HOMO SAPIENS (HUMAN), 354 aa	6.50E-192	9 (9q34)
-----	-----------	-----	--	---	---	-----	--------------	--------------------------	-------------	--	-----------	----------

585	cg2537639	826	GTTCTTCGGGGGG TCGGTGCAAGAG[G /A]TGACGCGCTC ACGAGGCGCTGCC A	G	A	Val	Met (814)	NON- CONSER VATIVE	transferase	Human Gene SWISSPROT- ID:P16442 FUCOSYLGLYCOPROTEIN ALPHA-N- ACETYL GALACTOSAMINYLT RANSFERASE (EC 2.4.1.40) (HISTO-BLOOD GROUP A TRANSFERASE) (A TRANSFERASE) / FUCOSYLGLYCOPROTEIN 3- ALPHA- GALACTOSYLTRANSFERASE (EC 2.4.1.37) (HISTO-BLOOD GROUP B TRANSFERASE) (B TRANSFERASE) (NAGAT) - HOMO SAPIENS (HUMAN), 354 aa.	6.50E-192	9 (9q34)
586	cg42742340	3249	CACATCGTGGTGG AGCTGACCCAGG[C /A]TGACGCTTGG GCTCCAGGTGGCG G	C	A	Ala	Asp (815)	NON- CONSER VATIVE	transport	Human Gene SWISSPROT- ID:Q04671 P PROTEIN (MELANOCYTE-SPECIFIC TRANSPORTER PROTEIN) - HOMO SAPIENS (HUMAN), 838 aa.	0	15
587	cg41653463	427	GTCTGAAAGATT CCACAAGGACAT[C /G]CTGAAGCCCTC ACGAGGGAAGAGC C	C	G	Ile	Met (816)	NON- CONSER VATIVE	transport	Human Gene SWISSPROT- ID:P31641 SODIUM- AND CHLORIDE-DEPENDENT TAURINE TRANSPORTER - HOMO SAPIENS (HUMAN), 620 aa.	0	3 (3p25)

588	cg40351913	1165	CAAGTTCACCAAC AACTGCTACAGG[G /C]ACGGGATTGTC ACCACCTCCATCA A	G	C	Asp	His (817)	NON- CONSER VATIVE	transport	Human Gene SWISSPROT- ID:Q01959 SODIUM- DEPENDENT DOPAMINE TRANSPORTER (DA TRANSPORTER) (DAT) - HOMO SAPIENS (HUMAN), 620 aa.	0	5 (5p15.3)
589	cg40351913	1232	TCCTCGGGCTTCGT CGTCTTCTCCTT[C] CCTGGGTACATG GCACAGAAGCAC	T	C	Phe	Ser (818)	NON- CONSER VATIVE	transport	Human Gene SWISSPROT- ID:Q01959 SODIUM- DEPENDENT DOPAMINE TRANSPORTER (DA TRANSPORTER) (DAT) - HOMO SAPIENS (HUMAN), 620 aa.	0	5 (5p15.3)
590	cg43955093	4776	CTGCGGTAGCTGT CCCAGGCCCTCGG[C /G]CCGGCGCGCT CGTCCATGTTGAG G	C	G	Ala	Pro (819)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SPTREMBL- ACC:Q16084 P130 - HOMO SAPIENS (HUMAN), 1139 aa.	0	16
591	cg43055918	522	GCATAGGACATGG CGGGCTTGCCCC[C/ G]CGCAGAGCTCTG GGGGCTACTGCTA	C	G	Gly	Arg (820)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P42694 HYPOTHETICAL PROTEIN KIAA0054 - Homo sapiens (Human), 1942 aa.	0	17
592	cg43968854	4604	CAACCCCTAGAAG ACCTGGCTGGCTT/ G]GAAAGAGCTCTT CCAGACACCAAGTA	T	G	Leu	Trp (821)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSNEW- ACC:P46013 ANTIGEN KI-67 - Homo sapiens (Human), 3256 aa.	0	10 (10q25)

593	cg43070241	1841	CCATTGTTCAAGA CATCCTACGTTT/T/ GJGAAATGCCCTGCA AGCAAAATTGTCC	T	G	Phe	Leu (822)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P55157 MICROSOMAL TRIGLYCERIDE TRANSFER PROTEIN, LARGE SUBUNIT PRECURSOR - Homo sapiens (Human), 894 aa.	0	4 (4q22)
594	cg43262121	2001	ACAATTCAGAGAG GGAGACTGAGCA/ GTJACACCAAGCAT TGATCATGGTGCC AA	G	T	Gln	His (823)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SPTREMBL- ACC:Q15840 ZINC FINGER PROTEIN BASONUCLIN - HOMO SAPIENS (HUMAN), 994 aa.	0	
595	cg43262121	553	ATCAGGAAAGGTG TTGGATCACTGGJA /TJGCATCATGACC AGTGAGGAAGAAG T	A	T	Ser	Cys (824)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SPTREMBL- ACC:Q15840 ZINC FINGER PROTEIN BASONUCLIN - HOMO SAPIENS (HUMAN), 994 aa.	0	
596	cg43262121	937	CCCCAAACAGGAA GTCCATGGGCCCA/ /TJACCCCTGACAGC AGCTTCTTAACCTC	A	T	Asn	Tyr (825)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SPTREMBL- ACC:Q15840 ZINC FINGER PROTEIN BASONUCLIN - HOMO SAPIENS (HUMAN), 994 aa.	0	
597	cg43262121	938	CCCCAAACAGGAA TCCATGGGCCCA/ /TJCCCTGACAGCA GCCTCTTAACCTCC	A	T	Asn	Ile (826)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SPTREMBL- ACC:Q15840 ZINC FINGER PROTEIN BASONUCLIN - HOMO SAPIENS (HUMAN), 994 aa.	0	

598	cg44024279	501	CTGGAAGAACCTT GCCATGAGAAAG A/GJAATTTGGAG AAGTACGGACATT CA	A	G	Glu	Gly (827)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P02771 ALPHA- FETOPROTEIN PRECURSOR (ALPHA-FETOGLOBULIN) (ALPHA-1-FETOPROTEIN) - Homo sapiens (Human), 609 aa.	0		
599	cg44928804	1235	AATGATTAAACAC AACCTGAGACACG /A/JCGGATGAAATG TTCTGGAAACCAAG T	G	A	Ala	Thr (828)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P21589 5'-NUCLEOTIDASE PRECURSOR (EC 3.1.3.5) (ECTO-NUCLEOTIDASE) (5'- NT) (CD73 ANTIGEN) - Homo sapiens (Human), 574 aa.	9.1e-313	6 (6q14)	
600	cg43317253	367	GCCCCCAGGCGATG GCTAGCTCGTGTG/ TJCCGTGCAGGTGA AGCTGGAGCTGGG	G	T	Ala	Ser (829)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSNEW- ACC:P42568 AF-9 PROTEIN - Homo sapiens (Human), 568 aa.	2.00E-301	9	
601	cg41637661	223	CAGCTTTCATCCA TTTTTATTAT[G/A] GACATACTGCTAG TGGAAAGACCTA	G	A	Gly	Arg (830)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSNEW- ACC:O43913 ORIGIN RECOGNITION COMPLEX SUBUNIT 5 - Homo sapiens (Human), 435 aa.	6.10E-236		
602	cg42913861	3034	CAGGTGTCTCTGCG AGCCACCCGGGG A/CJTCCGGGTGGC GGGGGTGGCGGCG GC	A	C	Ser	Ala (831)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P09529 INHIBIN BETA B CHAIN PRECURSOR (ACTIVIN BETA-B CHAIN) - Homo sapiens (Human), 407 aa.	3.00E-227	2 (2cen)	

603	cg43249389	526	AGAGGAGAGAGCC GCCCTCGAGCGG[A /G]GCAAGGCGATT GAGAAAAACCTCA A	G	Ser	Gly (832)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P09471 GUANINE NUCLEOTIDE-BINDING PROTEIN G(O), ALPHA SUBUNIT 1 - Homo sapiens (Human), 353 aa.	1.40E-188	15
604	cg43919239	335	GCCAGAGTTGCAG A CATCAGGGCCAG[A /C]CTGAGCAGGAG ACCCCCAGTCCCA T	C	Ser	Arg (833)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Homologous to SWISSPROT-ACCP14207 FOLATE RECEPTOR BETA PRECURSOR (FR-BETA) (FOLATE RECEPTOR 2) (FOLATE RECEPTOR, FETAL/PLACENTAL) (PLACENTAL FOLATE- BINDING PROTEIN) (FBP) - Homo sapiens (Human), 255 aa.	4.20E-150	
605	cg41642952	787	TAGGAATGACAGC A AGTAGCAGTAAT[A /G]GGAAGGCCAAA AATCCCCCTGGAG A	G	Arg	Gly (834)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Homologous to SWISSPROT-ACC:P21583 STEM CELL FACTOR PRECURSOR (SCF) (MAST CELL GROWTH FACTOR) (MGF) (C-KIT LIGAND) - Homo sapiens (Human), 273 aa.	3.70E-142	12

606	cg43945147	221	TGTTCTGGAGCCT CAATGGTACAG[G/ C]GTGCTCGAGAAG GACAGTGTGACTC	C	Arg	Ser (835)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Homologous to SWISSNEW-ACC:P08637 LOW AFFINITY IMMUNOGLOBULIN GAMMA FC RECEPTOR III-1 PRECURSOR (FC- GAMMA RIII) (FCRIII) (IGG FC RECEPTOR III-1) (FC-GAMMA RIII-ALPHA) (CD16) (FCR-10) - Homo sapiens (Human), 254 aa.	1.60E-134	1
607	cg43926002	391	GGGCACAGAAACA CAGCAGCGGAG[C/S]AGCAACACCA GCACTGCCAACAG AT	S	Ser	??? (836)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Homologous to SWISSPROT-ACC:P50539 MAX INTERACTING PROTEIN 1 (MXI1 PROTEIN) - Homo sapiens (Human), 228 aa.	1.60E-116	10
608	cg43972311	1609	ATTGCCATTGTGGT AACTCTGGGTCT[G]CATCATCTTCAGT GCCCAATTGTG	G	Glu	Ala (837)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to TREMBLNEW-ACC:AAD38008 GLYOXALASE-1 (EC 4.4.1.5) - HOMO SAPIENS (HUMAN), 184 aa.	2.20E-98	6
609	cg42556108	521	GTGAAGCGGTGTA TGGGGACAGTGA[C /A]CCTCAACCAGG CCAGGGGCTCCTTT	A	Thr	Asn (838)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to SWISSPROT-ACC:P49913 ANTIBACTERIAL PROTEIN FALL-39 PRECURSOR (FALL-39 PEPTIDE ANTIBIOTIC) (ANTIMICROBIAL PROTEIN CAP-18) (LL-37) - Homo sapiens (Human), 170 aa.	2.90E-87	3

610	cg36842490	487	AGTGACTTCAGTA AACTCTTGGGTC[A/ C]ACTTTCCTGCCAA AAAGTACCTTGAG	A	C	Gln	Pro (839)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to SWISSPROT-ACC:P01282 VASOACTIVE INTESTINAL PEPTIDE PRECURSOR (VIP) - Homo sapiens (Human), 170 aa.	2.30E-85	
611	cg43942549	1052	CGGTATAAGGTCA AAAATCTCTGTT[G/ T]TCAGCCCAAGGTT CAGAAATTGCCTC	G	T	Val (840)	Phe (840)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to SPTREMBL-ACC:Q94218 CODED FOR BY C. ELEGANS CDNA CM10H5 - CAENORHABDITIS ELEGANS, 589 aa.	2.80E-73	4
612	cg42381630	283	AAGGCGCTATGTA CAGCCTCCTGAA[A /G]TGATTGGCCT ATGCGGCCCGAGC A	A	G	Met (841)	Val (841)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to SPTREMBL-ACC:O76087 GAGE-8 - HOMO SAPIENS (HUMAN), 117 aa.	5.90E-64	
613	cg42381630	505	TGAAGATGGTCCT GATGGCAGGAG[A/G]TGGACCCGCC AAATCCAGAGGAG GT	A	G	Met (842)	Val (842)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to SPTREMBL-ACC:O76087 GAGE-8 - HOMO SAPIENS (HUMAN), 117 aa.	5.90E-64	
614	cg3004395	260	ATTACTGAAGGGT GGAGAACAGAA[G/ G/C]GTCATGAAAA AATATCTGCTTCAT T	G	C	Gly (843)	Arg (843)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to RENTREMBL-ACC:G238693 T CELL RECEPTOR VARIABLE ALPHA CHAIN - HOMO SAPIENS (HUMAN), 143 aa (fragment).	1.00E-59	14 (14q11.2)

615	cg43960645	733	CACTTCTCTTTCT CTTTGGATGCCAT TACCCCTCTGTTG GGGGGCAGATGG	A	T	Val	Glu (844)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to SWISSNEW-ACC:O76070 GAMMA-SYNUCLEIN (PERSYN) (BREAST CANCER- SPECIFIC GENE 1 PROTEIN) - Homo sapiens (Human), 127 aa.	1.20E-38	
616	cg2526759	289	GAAGACAAGGTGG TACAAAGCCCTCTT ATCTCTGTTGT CCACGAGGGAGAC	T	A	Leu	Gln (845)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to REMTREMBL-ACC:G33509 T CELL RECEPTOR - HOMO SAPIENS (HUMAN), 118 aa (fragment).	1.60E-54	
617	cg2526759	342	TGTAACCTCTCAATT GCAGTTATGAAIG/ ATGACTAACTTTC GAAOCCTACTATG	G	A	Val	Met (846)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to REMTREMBL-ACC:G33509 T CELL RECEPTOR - HOMO SAPIENS (HUMAN), 118 aa (fragment).	1.60E-34	
618	cg2526759	364	GAACTGACTAACT TTCGAAGCCTACIT/ AJATGGTACAAGCA GGAAAAGAAAGCT	T	A	Leu	Gln (847)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to REMTREMBL-ACC:G33509 T CELL RECEPTOR - HOMO SAPIENS (HUMAN), 118 aa (fragment).	1.60E-54	
619	cg2526759	475	AGCATATTAGATA AGAAAGAACTTTT /CJGAGCATCCTGA ACATCACAGCCAC C	T	C	Phe	Ser (848)	NON- CONSER VATIVE	UNCLASSIFIED	Human Gene Similar to REMTREMBL-ACC:G33509 T CELL RECEPTOR - HOMO SAPIENS (HUMAN), 118 aa (fragment).	1.60E-54	

620	cg40310734	1067	TACAGAAATATGTC GTCGGTGCCCCC[ga p/c]ACTTGGAGCTG GACCCCTGGGAGCG G	gap	C	Thr	His (849)	FRAMES HIFT	cadherin	Human Gene SWISSPROT- ID:P08514 PLATELET MEMBRANE GLYCOPROTEIN IIB PRECURSOR (GPIIB) (INTEGRIN ALPHA- IIB) (CD41) - HOMO SAPIENS (HUMAN), 1039 aa.	0	17 (17q21.3 2)
621	cg40310734	3285	GTCGGCTTCTTCAA GCGGAACCGGC[ga p/a]CACCCCTGGAA GAAGATGATGAAG A	gap	A	Pro	His (850)	FRAMES HIFT	cadherin	Human Gene SWISSPROT- ID:P08514 PLATELET MEMBRANE GLYCOPROTEIN IIB PRECURSOR (GPIIB) (INTEGRIN ALPHA- IIB) (CD41) - HOMO SAPIENS (HUMAN), 1039 aa.	0	17 (17q21.3 2)
622	cg43956660	2521	GTCCATCACCTTCAC TTCAGTTATTC[ga p]CTAGGAGGTTG TATAGTCTTCTGA	T	gap	Arg	Glu (851)	FRAMES HIFT	cadherin	Human Gene SWISSNEW- ID:Q08722 LEUKOCYTE SURFACE ANTIGEN CD47 PRECURSOR (ANTIGENIC SURFACE DETERMINANT PROTEIN OA3) (INTEGRIN ASSOCIATED PROTEIN) (IAP) (MER6) - HOMO SAPIENS (HUMAN), 323 aa.[pcis:SWISSPROT-ID:Q08722 LEUKOCYTE SURFACE ANTIGEN CD47 PRECURSOR (ANTIGENIC SURFACE DETERMINANT PROTEIN OA3) (INTEGRIN ASSOCIATED PROTEIN) (IAP) (MER6) - HOMO SAPIENS (HUMAN), 323 aa.	1.80E-157	

623	cg43970982	2429	CTCCAGGGATAGT TGGACAGAAAGGG[ap/GJAGACCTGGC TACCCAGGACCAG CT	gap	G	Gly	Gly (852)	FRAMES HIFT	collagen	Human Gene SWISSPROT- ID:P12111 COLLAGEN ALPHA 3(VT) CHAIN PRECURSOR - HOMO SAPIENS (HUMAN), 3176 aa.	0	2
624	cg42175288	1837	GCTATGGAGGCAA AATGGGAGGAAG[ap/GJAAACGACTAC AGAAATGATCAGC GC	gap	G	Arg	Arg (853)	FRAMES HIFT	dna_rna_bind	Human Gene SPTREMBL- ID:Q92804 PUTATIVE RNA BINDING PROTEIN RBP56 - HOMO SAPIENS (HUMAN), 592 aa.	0	17
625	cg42175288	263	CGGTACTCCAGTT ATGGACAAAGT[ga p/CJTATTCACAGTC CTATGGTGGTTATG	gap	C	Tyr	Leu (854)	FRAMES HIFT	dna_rna_bind	Human Gene SPTREMBL- ID:Q92804 PUTATIVE RNA BINDING PROTEIN RBP56 - HOMO SAPIENS (HUMAN), 592 aa.	0	17
626	cg41554010	584	GGCCGAGCAGCTG CGGCGCCAGCTG[ap/GJACCCCTACG CACAGCGCATGGA GA	gap	G	Thr	Asp (855)	FRAMES HIFT	eph	Human Gene SWISSNEW- ID:P06727 APOLIPOPROTEIN A- IV PRECURSOR (APO-AIV) - HOMO SAPIENS (HUMAN), 396 aa.[pcls:SWISSPROT-ID:P06727 APOLIPOPROTEIN A-IV PRECURSOR (APO-AIV) - HOMO SAPIENS (HUMAN), 396 aa.	1.80E-203	11 (11q23)
627	cg43065549	1553	CAGACTTCACAG AGTGCTGGATGA[ap/AJCGCGGCTGC CTTGCCCCAGGGTT A	gap	A	Thr	Asn (856)	FRAMES HIFT	glycoprotein	Human Gene SWISSPROT- ID:P16452 ERYTHROCYTE MEMBRANE PROTEIN BAND 4.2 (P4.2) (PALLIDIN) - HOMO SAPIENS (HUMAN), 690 aa.	0	15 (15q15)

628	cg41568631	999	TGACCACGGGGTG CTGGATGCCCTGC[ga p/CITTTATACATCCT GGACCGGCGGGGG A	gap	C	Leu (857)	FRAMES HIIFT	glycoprotein	Human Gene Similar to SWISSPROT-ID:P16452 ERYTHROCYTE MEMBRANE PROTEIN BAND 4.2 (P4.2) (PALLIDIN) - HOMO SAPIENS (HUMAN), 690 aa.	9.90E-70	14 (14q11.2)
629	cg41637704	1220	GCGCGCGGAGACA AGGGCAGCGGAC[g ap/GJCGCCTGCGGA CTTGAGGGACAGT GA	gap	G	Pro (858)	FRAMES HIIFT	homeobox	Human Gene SWISSPROT- ID:P50219 HOMEBOX PROTEIN HB9 - HOMO SAPIENS (HUMAN), 401 aa.	1.20E-224	7
630	cg43933380	364	ATAAGTTACAATG CTTTTTTGTITTA/g ap/JAAAAAAAAAA AAGTCTGTACTTTA	A	gap	Leu (859)	FRAMES HIIFT	interferon	Human Gene SWISSPROT- ID:P15260 INTERFERON- GAMMA RECEPTOR ALPHA CHAIN PRECURSOR (CDW119) - HOMO SAPIENS (HUMAN), 489 aa.	1.40E-261	6
631	cg43072541	379	CTGTGGGGCTGGT TCTGTATCTGAT[ga p/CJATCAITTCGATT ACGAAATATAAACG T	gap	C	Ile (860)	FRAMES HIIFT	kinase	Human Gene SPTREMBL- ID:Q15802 SERINE/THREONINE PROTEIN KINASE KRS-2 - HOMO SAPIENS (HUMAN), 487 aa.	9.60E-262	20

632	cg44032168	1536	GTCAGCCGCTACC TCGACTGGATCCGga p/TJATGGGCACATC AGAGACAAGGAAG C	gap	T	His	Leu (861)	FRAMES HIIFT	protease	Human Gene Similar to SWISSPROT-ID:P25155 COAGULATION FACTOR X PRECURSOR (EC 3.4.21.6) (STUART FACTOR) (VIRUS ACTIVATING PROTEASE) (VAP) - GALLUS GALLUS (CHICKEN), 475 aa.	2.40E-82	2 (2q13)
633	cg43931248	1317	CCGGGCAGAGCTG CGTCTGCTGAGGlg ap/GJCTCAAGTTAA AAGTGGAGCAGCA CG	gap	G	Leu	Ala (862)	FRAMES HIIFT	tgf	Human Gene SWISSPROT- ID:P01137 TRANSFORMING GROWTH FACTOR BETA 1 PRECURSOR (TGF-BETA 1) - HOMO SAPIENS (HUMAN), 390 aa.	9.70E-214	19
634	cg43931248	1317	CCGGGCAGAGCTG CGTCTGCTGAGGlg ap/GJCTCAAGTTAA AAGTGGAGCAGCA CG	gap	G	Leu	Ala (863)	FRAMES HIIFT	tgf	Human Gene SWISSPROT- ID:P01137 TRANSFORMING GROWTH FACTOR BETA 1 PRECURSOR (TGF-BETA 1) - HOMO SAPIENS (HUMAN), 390 aa.	9.70E-214	19
635	cg43272560	847	AATCTCCGCACTG CAGGCCAGGGGClg ap/CJTGGCCAGCTA CAGAGAGAGGTCA CA	gap	C	Ala	Ala (864)	FRAMES HIIFT	tgfreceptor	Human Gene SWISSPROT- ID:Q03167 TGF-BETA RECEPTOR TYPE III PRECURSOR (TGFR-3) (BETAGLYCAN) - HOMO SAPIENS (HUMAN), 849 aa.	0	1 (1p33)

636	cg43266471	1067	CCAGGATCCATT GAGGATTATGG[/TGTGCTGGGACA CCATCAACTCCTCA	gap	T	Gly	Gly (865)	FRAMES HIFT	tm7	Human Gene SWISSPROT- ID:P32241 VASOACTIVE INTESTINAL POLYPEPTIDE RECEPTOR I PRECURSOR (VIP-R-1) (PITUITARY ADENYLATE CYCLASE ACTIVATING POLYPEPTIDE TYPE II RECEPTOR) (PACAP- R-2) - HOMO SAPIENS (HUMAN), 457 aa.	5.20E-254	3
637	cg43995237	625	CAAATCCCCCGTT TCTTCATCTTG[GACATGCTAAAT GAAATTACGCAGT	gap	G	Gln	Pro (866)	FRAMES HIFT	transferase	Human Gene SWISSPROT- ID:P53611 GERANYLGERANYL TRANSFERASE TYPE II BETA SUBUNIT (EC 2.5.1.-) (RAB GERANYLGERANYLTRANSFERASE BETA SUBUNIT) (RAB GERANYL- GERANYLTRANSFERASE BETA SUBUNIT) (RAB GG TRANSFERASE) (RAB GGTASE) - HOMO SAPIENS (HUMAN), 331 aa.	1.40E-182	1

638	cg43995237	638	TTCCTCATCTTGAC ATGCTAAATG[/GJAAATTACGCAG TTTCTCTATCAA	gap	G	Leu	Phe (867)	FRAMES HIFT	transferase	Human Gene SWISSPROT- ID:P53611 GERANYLGERANYL TRANSFERASE TYPE II BETA SUBUNIT (EC 2.5.1.-) (RAB GERANYLGERANYLTRANSFERASE BETA SUBUNIT) (RAB GERANYL- GERANYLTRANSFERASE BETA SUBUNIT) (RAB GG TRANSFERASE) (RAB GGTASE) - HOMO SAPIENS (HUMAN), 331 aa.	1.40E-182	1
639	cg43254094	267	CCGCCTCTGCTGCT GCTGCTGCTGCT[/GJGCGTCCCGCCC AGCCGCGAGCTTCC C	gap	G	Arg	Arg (868)	FRAMES HIFT	UNCLASSIFIED	Human Gene SWISSPROT- ACC:P78339 SUSHI REPEAT- CONTAINING PROTEIN SRPX PRECURSOR - Homo sapiens (Human), 464 aa.	6.40E-257	X
640	cg44034555	665	ATCCAGGCTGAGC TGGATCATCTGA[/gapJGGCTCCAGC CACCCGTTTCCCT T	G	gap	Pro	Leu (869)	FRAMES HIFT	UNCLASSIFIED	Human Gene SWISSNEW- ACC:Q13228 SELENIUM- BINDING PROTEIN 1 - Homo sapiens (Human), 472 aa.	3.80E-252	1
641	cg44034555	667	CCAGGCTGAGCTG GATCATCTGAGG[/gapJCTCCAGCCA CCCGTTTCCCTTG A	G	gap	Gly	Gly (870)	FRAMES HIFT	UNCLASSIFIED	Human Gene SWISSNEW- ACC:Q13228 SELENIUM- BINDING PROTEIN 1 - Homo sapiens (Human), 472 aa.	3.80E-252	1

642	cg39711096	882	AGCAGTCTCTCCG GGAGGCCACACAG[ap/GjTTACTGCCTC CAGCTGCAGCAGT GA	gap	G	Val	Gly (871)	FRAMES HIFT	UNCLASSIFIED D	Human Gene SWISSPROT- ACC:P18428 LIPOPOLYSACCHARIDE- BINDING PROTEIN PRECURSOR (LBP) - Homo sapiens (Human), 481 aa.	1.00E-251	
643	cg44128902	379	CGTTCAGAGGAG CATATCTGCTGA[ga p/CTGATGACCTGC AAGAGTCATCCAG A	gap	C	Asp	Asp (872)	FRAMES HIFT	UNCLASSIFIED D	Human Gene SWISSPROT- ACC:P18615 RD PROTEIN - Homo sapiens (Human), 380 aa.	1.00E-201	1 (1p36.2)
644	cg43946951	306	GGAACTCGAGCAC GTCGTCGGGGGA[C /gap]CCCAAGATCA CCGGGCCCTCTG GT	C	gap	Gly	Gly (873)	FRAMES HIFT	UNCLASSIFIED D	Human Gene SWISSPROT- ACC:P09467 FRUCTOSE-1,6- BISPHOSPHATASE (EC 3.1.3.11) (D-FRUCTOSE-1,6- BISPHOSPHATE 1- PHOSPHOHYDROLASE) (FBPASE) - Homo sapiens (Human), 337 aa.	3.50E-178	9 (9q22.2)
645	cg43948890	195	ATTCCCGGGGAG GGGCCCTGTAA[G /gap]GGAAACCAGA CAATCCCATGAGA CT	G	gap	Pro	Leu (874)	FRAMES HIFT	UNCLASSIFIED D	Human Gene Homologous to SPTREMBL-ACC:Q15182 SNRNP POLYPEPTIDE B - HOMO SAPIENS (HUMAN), 285 aa.	3.20E-147	20
646	cg43948890	197	TCCCGGGGAGGG GGCCTGTAAAG[G /gap]AAACCAGACA ATCCCATGAGACT CC	G	gap	Phe	Phe (875)	FRAMES HIFT	UNCLASSIFIED D	Human Gene Homologous to SPTREMBL-ACC:Q15182 SNRNP POLYPEPTIDE B - HOMO SAPIENS (HUMAN), 285 aa.	3.20E-147	20

647	cg43917524	713	GGGCTGTCTGCC CAGTGGAGGAGG[C/gap]TTCCGCTGGT GTTCTAGGGGCA TC	C	gap	Ala	Pro (876)	FRAMES HIIT	UNCLASSIFIED D	Human Gene Homologous to TREMBLNEW-ACC:AAD43025 PTD017 - HOMO SAPIENS (HUMAN), 258 aa.	3.20E-143	
648	cg43942004	373	CTCTGGCACTGGT GACTGGCGAGAG[ga p/GJCCTGGAGCGG CTTCGGAGAGGGC TA	gap	G	Asp	Glu (877)	FRAMES HIIT	UNCLASSIFIED D	Human Gene Homologous to SWISSNEW-ACC:Q99075 HEPARIN-BINDING EGF-LIKE GROWTH FACTOR PRECURSOR (HB-EGF) (HBEGF) (DIPHTERIA TOXIN RECEPTOR) (DT-R) - Homo sapiens (Human), 208 aa.	1.00E-107	5 (5q23)
649	cg43932428	681	TCGTGGCCAGGTC CTTCTGCGTAAG[C/ gap]CCCTTGCTCTG CCGACCTTGCTGG A	C	gap	Gly	Gly (878)	FRAMES HIIT	UNCLASSIFIED D	Human Gene Similar to SPTREMBL-ACC:O60869 EDF-1 PROTEIN - HOMO SAPIENS (HUMAN), 148 aa.	2.50E-72	
650	cg44010855	450	GGTCCAAATGCAA GTGCTCCCGGAA[G /gap]GGACCCCAAGA TCCGCTACAGCGA CG	G	gap	Gly	Asp (879)	FRAMES HIIT	UNCLASSIFIED D	Human Gene Similar to TREMBLNEW-ACC:AAD38944 NJAC PROTEIN - HOMO SAPIENS (HUMAN), 99 aa.	5.80E-50	5
651	cg44010855	452	TCCAAATGCAAGT GCTCCCGGAAAGG[G /gap]ACCCCAAGATC CGCTACAGOGACG TG	G	gap	Gly	Asp (880)	FRAMES HIIT	UNCLASSIFIED D	Human Gene Similar to TREMBLNEW-ACC:AAD38944 NJAC PROTEIN - HOMO SAPIENS (HUMAN), 99 aa.	5.80E-50	5

CLAIMS**WHAT IS CLAIMED IS:**

1. An isolated polynucleotide selected from the group consisting of:
 - a) a nucleotide sequence comprising one or more polymorphic sequences
(SEQ ID NOS:1 - 651);
 - b) a fragment of said nucleotide sequence, provided that the fragment
includes a polymorphic site in said polymorphic sequence;
 - c) a complementary nucleotide sequence comprising a sequence
complementary to one or more of said polymorphic sequences (SEQ ID
NOS:1 - 651); and
 - d) a fragment of said complementary nucleotide sequence, provided that the
fragment includes a polymorphic site in said polymorphic sequence.
2. The polynucleotide of claim 1, wherein said polynucleotide sequence is DNA.
3. The polynucleotide of claim 1, wherein said polynucleotide sequence is RNA.
4. The polynucleotide of claim 1, wherein said polynucleotide sequence is between
about 10 and about 100 nucleotides in length.
5. The polynucleotide of claim 1, wherein said polynucleotide sequence is between
about 10 and about 90 nucleotides in length.

6. The polynucleotide of claim 1, wherein said polynucleotide sequence is between about 10 and about 75 nucleotides in length.
7. The polynucleotide of claim 1, wherein said polynucleotide is between about 10 and about 50 bases in length.
8. The polynucleotide of claim 1, wherein said polynucleotide is between about 10 and about 40 bases in length.
9. The polynucleotide of claim 1, wherein said polynucleotide is derived from a nucleic acid encoding a polypeptide related to angiopoietin, 4-hydroxybutyrate dehydrogenase, ATP-dependent RNA helicase, MHC Class I histocompatibility antigen, or phosphoglycerate kinase.
10. The polynucleotide of claim 1, wherein said polymorphic site includes a nucleotide other than the nucleotide listed in Table 1, column 5 for said polymorphic sequence.
11. The polynucleotide of claim 1, wherein the complement of said polymorphic site includes a nucleotide other than the complement of the nucleotide listed in Table 1, column 5 for the complement of said polymorphic sequence.
12. The polynucleotide of claim 1, wherein said polymorphic site includes the nucleotide listed in Table 1, column 6 for said polymorphic sequence.

13. The polynucleotide of claim 1, wherein the complement of said polymorphic site includes the complement of the nucleotide listed in Table 1, column 6 for said polymorphic sequence.

5 14. An isolated allele-specific oligonucleotide that hybridizes to a first polynucleotide at a polymorphic site encompassed therein, wherein the first polynucleotide is chosen from the group consisting of:

10 a) a nucleotide sequence comprising one or more polymorphic sequences (SEQ ID NOS:1 - 651) provided that the polymorphic sequence includes a nucleotide other than the nucleotide recited in Table 1, column 5 for said polymorphic sequence;

b) a nucleotide sequence that is a fragment of said polymorphic sequence, provided that the fragment includes a polymorphic site in said polymorphic sequence;

15 c) a complementary nucleotide sequence comprising a sequence complementary to one or more polymorphic sequences (SEQ ID NOS:1 - 651), provided that the complementary nucleotide sequence includes a nucleotide other than the complement of the nucleotide recited in Table 1, column 5; and

20 d) a nucleotide sequence that is a fragment of said complementary sequence, provided that the fragment includes a polymorphic site in said polymorphic sequence.

25 15. The oligonucleotide of claim 14, wherein the oligonucleotide does not hybridize under stringent conditions to a second polynucleotide selected from the group consisting of:

- a) a nucleotide sequence comprising one or more polymorphic sequences (SEQ ID NOS:1 - 651), wherein said polymorphic sequence includes the nucleotide listed in Table 1, column 5 for said polymorphic sequence;
- b) a nucleotide sequence that is a fragment of any of said nucleotide sequences;
- c) a complementary nucleotide sequence comprising a sequence complementary to one or more polymorphic sequences (SEQ ID NOS:1 - 651), wherein said polymorphic sequence includes the complement of the nucleotide listed in Table 1, column 5; and
- d) a nucleotide sequence that is a fragment of said complementary sequence, provided that the fragment includes a polymorphic site in said polymorphic sequence.
16. The oligonucleotide of claim 15, wherein the oligonucleotide is between about 10 and about 51 bases in length.
17. The oligonucleotide of claim 15, wherein the oligonucleotide identifies a polypeptide related to angiopoietin, 4-hydroxybutyrate dehydrogenase, ATP-dependent RNA helicase, MHC Class I histocompatibility antigen, or phosphoglycerate kinase.
18. The oligonucleotide of claim 15, wherein the oligonucleotide is between about 15 and about 30 bases in length.
19. A method of detecting a polymorphic site in a nucleic acid, the method comprising:
- a) contacting said nucleic acid with an oligonucleotide that hybridizes to a polymorphic sequence selected from the group consisting of SEQ ID NOS: 1 - 651, or its complement, provided that the polymorphic sequence

includes a nucleotide other than the nucleotide recited in Table 1, column 5 for said polymorphic sequence, or the complement includes a nucleotide other than the complement of the nucleotide recited in Table 1, column 5; and

5 b) determining whether said nucleic acid and said oligonucleotide hybridize;

whereby hybridization of said oligonucleotide to said nucleic acid sequence indicates the presence of the polymorphic site in said nucleic acid.

20. The method of claim 19, wherein said oligonucleotide does not hybridize to said
10 polymorphic sequence when said polymorphic sequence includes the nucleotide recited in Table 1, column 5 for said polymorphic sequence, or when the complement of the polymorphic sequence includes the complement of the nucleotide recited in Table 1, column 5 for said polymorphic sequence.

15 21. The method of claim 19, wherein said oligonucleotide identifies a polypeptide related to angiopoietin, 4-hydroxybutyrate dehydrogenase, ATP-dependent RNA helicase, MHC Class I histocompatibility antigen, or phosphoglycerate kinase.

22. The method of claim 19, wherein said oligonucleotide is between about 15 and about
20 30 bases in length.

23. A method of detecting the presence of a sequence polymorphism in a subject, the method comprising:

a) providing a nucleic acid from said subject;

25 b) contacting said nucleic acid with an oligonucleotide that hybridizes to a polymorphic sequence selected from the group consisting of SEQ ID NOS:1 - 651, or its complement, provided that the polymorphic sequence

includes a nucleotide other than the nucleotide recited in for said polymorphic sequence, or the complement includes a nucleotide other than the complement of the nucleotide recited in Table 1, column 5; and

c) determining whether said nucleic acid and said oligonucleotide hybridize;

5 whereby hybridization of said oligonucleotide to said nucleic acid sequence indicates the presence of the polymorphism in said subject.

24. A method of determining the relatedness of a first and second nucleic acid, the method comprising:

- 10 a) providing a first nucleic acid and a second nucleic acid;
- b) contacting said first nucleic acid and said second nucleic acid with an oligonucleotide that hybridizes to a polymorphic sequence selected from the group consisting of SEQ ID NOS:1 - 651, or its complement, provided that the polymorphic sequence includes a nucleotide other than the
- 15 nucleotide recited in Table 1, column 5 for said polymorphic sequence, or the complement includes a nucleotide other than the complement of the nucleotide recited in Table 1, column 5;
- c) determining whether said first nucleic acid and said second nucleic acid hybridize to said oligonucleotide; and
- 20 d) comparing hybridization of said first and second nucleic acids to said oligonucleotide,

wherein hybridization of the first and second nucleic acids to said oligonucleotide indicates the first and second nucleic acids are related.

25 25. The method of claim 24, wherein said oligonucleotide does not hybridize to said polymorphic sequence when said polymorphic sequence includes the nucleotide recited in Table 1, column 5 for said polymorphic sequence, or when the complement

of the polymorphic sequence includes the complement of the nucleotide recited in Table 1, column 5 for said polymorphic sequence.

- 5 26. The method of claim 24, wherein the oligonucleotide is between about 10 and about 51 bases in length.
27. The method of claim 24, wherein the oligonucleotide is between about 10 and about 40 bases in length.
- 10 28. The method of claim 24, wherein the oligonucleotide is between about 15 and about 30 bases in length.
- 15 29. An isolated polypeptide comprising a polymorphic site at one or more amino acid residues, wherein the protein is encoded by a polynucleotide selected from the group consisting of: polymorphic sequences SEQ ID NOS:1 - 651, or their complement, provided that the polymorphic sequence includes a nucleotide other than the nucleotide recited in Table 1, column 5 for said polymorphic sequence, or the complement includes a nucleotide other than the complement of the nucleotide recited in Table 1, column 5.
- 20 30. The polypeptide of claim 29, wherein said polypeptide is translated in the same open reading frame as is a wild type protein whose amino acid sequence is identical to the amino acid sequence of the polymorphic protein except at the site of the polymorphism.
- 25 31. The polypeptide of claim 29, wherein the polypeptide encoded by said polymorphic sequence, or its complement, includes the nucleotide listed in Table 2, column 6 or

Table 3, column 5 for said polymorphic sequence, or the complement includes the complement of the nucleotide listed in Table 1, column 6.

32. An antibody that binds specifically to a polypeptide encoded by a polynucleotide comprising a nucleotide sequence encoded by a polynucleotide selected from the group consisting of polymorphic sequences SEQ ID NOS:1 - 651, or its complement, provided that the polymorphic sequence includes a nucleotide other than the nucleotide recited in Table 1, column 5 for said polymorphic sequence, or the complement includes a nucleotide other than the complement of the nucleotide recited in Table 1, column 5.

33. The antibody of claim 32, wherein said antibody binds specifically to a polypeptide encoded by a polymorphic sequence which includes the nucleotide listed in Table 1, column 6 for said polymorphic sequence.

34. The antibody of claim 32, wherein said antibody does not bind specifically to a polypeptide encoded by a polymorphic sequence which includes the nucleotide listed in Table 1, column 5 for said polymorphic sequence.

35. A method of detecting the presence of a polypeptide having one or more amino acid residue polymorphisms in a subject, the method comprising

- a) providing a protein sample from said subject;
- b) contacting said sample with the antibody of claim 34 under conditions that allow for the formation of antibody-antigen complexes; and
- c) detecting said antibody-antigen complexes,

whereby the presence of said complexes indicates the presence of said polypeptide.

36. A method of treating a subject suffering from, at risk for, or suspected of, suffering from a pathology ascribed to the presence of a sequence polymorphism in a subject, the method comprising:

- 5 a) providing a subject suffering from a pathology associated with aberrant expression of a first nucleic acid comprising a polymorphic sequence selected from the group consisting of SEQ ID NOS:1 - 651, or its complement; and
- 10 b) administering to the subject an effective therapeutic dose of a second nucleic acid comprising the polymorphic sequence, provided that the second nucleic acid comprises the nucleotide present in a wild type allele of the sequence polymorphism,

thereby treating said subject.

15 37. The method of claim 36, wherein the second nucleic acid sequence comprises a polymorphic sequence which includes the nucleotide listed in Table 1, column 5 for said polymorphic sequence.

20 38. A method of treating a subject suffering from, at risk for, or suspected of suffering from a pathology ascribed to the presence of a sequence polymorphism in a subject, the method comprising:

- 25 a) providing a subject suffering from a pathology associated with aberrant expression of a polymorphic sequence selected from the group consisting of polymorphic sequences SEQ ID NOS:1 - 651, or its complement; and
- b) administering to the subject an effective therapeutic dose of a polypeptide,

wherein said polypeptide is encoded by a polynucleotide comprising a polymorphic sequence selected from the group consisting of SEQ ID NOS:1 - 651, or by a polynucleotide comprising a nucleotide sequence that is complementary to any one of polymorphic sequences SEQ ID NOS:1 - 651, provided that said polymorphic sequence includes the nucleotide listed in Table 1, column 6 for said polymorphic sequence, thereby treating said subject.

39. A method of treating a subject suffering from, at risk for, or suspected of suffering from, a pathology ascribed to the presence of a sequence polymorphism in a subject, the method comprising:

a) providing a subject suffering from, at risk for, or suspected of suffering from, a pathology associated with aberrant expression of a first nucleic acid comprising a polymorphic sequence selected from the group consisting of SEQ ID NOS:1 - 651, or its complement; and

b) administering to the subject an effective dose of the antibody of claim 34, thereby treating said subject.

40. A method of treating a subject suffering from, at risk for, or suspected of suffering from, a pathology ascribed to the presence of a sequence polymorphism in a subject, the method comprising:

a) providing a subject suffering from, at risk for, or suspected of suffering from, a pathology associated with aberrant expression of a nucleic acid comprising a polymorphic sequence selected from the group consisting of SEQ ID NOS:1 - 651, or its complement; and

b) administering to the subject an effective dose of an oligonucleotide comprising a polymorphic sequence selected from the group consisting of SEQ ID NOS:1 - 651, or by a polynucleotide comprising a nucleotide sequence that is complementary to any one of polymorphic sequences SEQ

ID NOS:1 - 651, provided that said polymorphic sequence includes the nucleotide listed in Table 1, column 6 for said polymorphic sequence,

thereby treating said subject.

- 5 41. An oligonucleotide array, comprising one or more oligonucleotides hybridizing to a first polynucleotide at a polymorphic site encompassed therein, wherein the first polynucleotide is chosen from the group consisting of:
- a) a nucleotide sequence comprising one or more polymorphic sequences SEQ ID NOS:1 - 651;
- 10 b) a nucleotide sequence that is a fragment of any of said nucleotide sequence, provided that the fragment includes a polymorphic site in said polymorphic sequence;
- c) a complementary nucleotide sequence comprising a sequence complementary to one or more polymorphic sequences SEQ ID NOS:1 -
- 15 651; and
- d) a nucleotide sequence that is a fragment of said complementary sequence, provided that the fragment includes a polymorphic site in said polymorphic sequence.
- 20 42. The array of claim 41, wherein said array comprises 10 oligonucleotides.
43. The array of claim 41, wherein said array comprises at least 100 oligonucleotides.
44. The array of claim 41, wherein said array comprises at least 1000 oligonucleotides.

1/1




FIG. 1

SEQUENCE LISTING

<110> CuraGen Corporation
Shimkets, Richard A.
Leach, Martin D.

<120> Nucleic Acids Containing Single Nucleotide Polymorphisms and Methods of Use Thereof

<130> 15966-534C-CIP1 PCT

<140> not yet assigned

<141> 2000-12-27

<150> 09/472,688

<151> 1999-12-27

<150> 60/109,024

<151> 1998-11-17

<150> 09/443,199

<151> 1999-11-16

<150> 09/442,129

<151> 1999-11-16

<150> 09/442,849

<151> 1999-11-17

<160> 880

<170> CuraGen Patent Formatter Version 0.9

<210> 1

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43333349

<400> 1

CGCTGACAGG GGAGTCTGAG CCACAGACCC GCTCACCCGA GTGCACGCAC G

51

<210> 2

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43931765

<400> 2

ATGGATAGTC CATCTGGTTG GATGCTGTGT ACTCGTTGGC CTCGTTTCAGG T

51

<210> 3

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg44130533

<400> 3

AATACAAAGC TGAGTGGAGA GCAGTGGGTG AAGAAGTATG GCATTCCAAG T

51

<210> 4

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg34888922

<400> 4

TATTGTTATT ATGTATTCTG TTTACGTGTT TCTGTGTCAC TGCTAAGAGA A

51

<210> 5

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg34888922

<400> 5

CAAGGAGCAT TGACCGTGAG AAATATGAAC AGTTTGCGTT ATATGGCTAT G

51

<210> 6

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg40310734

<400> 6

TGGCGTCGTA TTTTGGGCAT TCAGTCGCTG TCACTGACGT CAACGGGGAT G

51

<210> 7

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg40310734

<400> 7

AGGGGGCCTA TGAAGCAGAG CTGGCGGTGC ACCTGCCCCA GGGCGCCCAC T

51

<210> 8

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg40310734

<400> 8

GTTACTGTGA AGCGGGCTTC AGCTCGGTGG TCACTCAGGC CGGAGAGCTG G

51

<210> 9

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43331935

<400> 9

GTGACAAGTA CTTCATAGAG GATGGTCGCC TGGTCATCCA CAGCCTGGAC T

51

<210> 10

<211> 51

<212> DNA

<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42388009

<400> 10
AAGGAGAAAA CAATGAAGAA CCGAATGAAG ACGAAGACTC TGAGGCTGAG A

51

<210> 11
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42388009

<400> 11
AAAACAATGA AGAACCGAAC GAAGATGAAG ACTCTGAGGC TGAGAATACC A

51

<210> 12
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44126574

<400> 12
AGAACGGCCA GCCCCTGTGG ATCCTGGGGG ATGTCTTCCT CAGGTCCTAC T

51

<210> 13
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43970983

<400> 13
GGACTCCAGT GTCCAGGGCA TCCAGTTACA TCCTATCCTG GCGGCCACTC A

51

<210> 14
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44032748

<400> 14
TAAGACGGGC AGCTACACCC GCAGCGGTTA CCTGCCAGCT GAGCAACTGG T 51

<210> 15
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41553795

<400> 15
TCCAGCCCAA GGCCAATTTT GATGCGCAGC AGTTTGCAGG GACCTGGCTC C 51

<210> 16
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43942011

<400> 16
AGGTAGGAGG GCTTGGTCTC CAAACGCCTA TTGTTTCATT CTCCACAGTG C 51

<210> 17
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg21644442

<400> 17
AACAGCCGGC AGATGTAACT GGTACCGCCT TGCCCAGGGT GGGCCCCGTG A 51

<210> 18
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41533258

<400> 18
TCCAGCGCCG GGCAGGAGGG GTCCTAGTTG CCTCCCATCT GCAGAGCTTC C 51

<210> 19
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43996714

<400> 19
ATGTGCCAC TGCAATTGGGT TGTCCGGGAG TTGATACTGG TGGGATCACA G 51

<210> 20
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43259523

<400> 20
CAGAATATGG AGGCACAGGA GCTTCTTTT TATCCACTGT GTCGTGATA G 51

<210> 21
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43057018

<400> 21
GAATAAGAAA TTCAATCTGG ATGCATTGGT GACCCATACC CTGCCTTTTG A 51

<210> 22
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg1395871

<400> 22
GTGCTCCAGA GGGGCCAGCA GGCACGGGAA AAACCGAAAC CACCAAGGAC T 51

<210> 23
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg1395871

<400> 23
CAGAGGGGCC AGCAGGCACA GGAAAGACCG AAACCACCAA GGACTTGGCT A 51

<210> 24
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg1395871

<400> 24
AGCAATGGGA AAGTTTTTTA AAGGATTGGC TTCTTCTGGT GCTTGGGCTT G 51

<210> 25
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg1395871

<400> 25
CTTCTTCTGG TGCTTGGGCT TGCTTTGATG AATTCAACCG GATTGAGTTG G 51

<210> 26
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43950268

<400> 26
AGCGGCCAC CATGGCCCTA GGGTCATCAA CAAGTCCAGC AGCAATCATG G 51

<210> 27
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43918531

<400> 27
CCGATGGCTA TGAGCAGGCT GCTCGTGTG CTATTGAACA CCTGGACAAG A 51

<210> 28
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43957743

<400> 28
CAAGTTCCTC AATAAAGTGG CAGTTTTCAG GTTCTACTGG CTCCACTTCT C 51

<210> 29

<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43319420

<400> 29
TCACCCTCAG GAGGTGGCTG TTCTGTGTCC ACGACAATA CAGAAACAAC C

51

<210> 30
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg3001932

<400> 30
TCTTCAACAT CGTCTATTGG CTTTATTATG TGAATAAAA CATGGCCTCC C

51

<210> 31
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43975899

<400> 31
GGATTTTGA CAGACTCCTA GATGGTTATG ACAATCGCCT GAGACCAGGA T

51

<210> 32
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43299024

<400> 32
GGGCCCACTT CCCCCTGGAC GTCCAGTGG ACGACCTGGA CTACATGGAC T 51

<210> 33
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43299024

<400> 33
TGAACGAGCC TTCCAACCTC ATCAGAGGCT CTGAGGACGG CTGCCCCAAC A 51

<210> 34
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43969076

<400> 34
AATTCCAAAT GAGCTCTCCA ACCACATATT TTCTGCGTTT TTGATCCAGA C 51

<210> 35
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43969014

<400> 35
AATTCCAGAT GAGCTCTCCA ACCACATATT TTCTGCGTTT TTGATCCAGA C 51

<210> 36
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43065549

<400> 36
GGACCATCTC TGTGACCACA CCTGCAGACG CTGTCATTGG CCACTACTCG C 51

<210> 37
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43065549

<400> 37
ACCCCTGGAA TAGAGAGGAT GCTGTGTTCC TGAAGAATGA GGCTCAGCGC A 51

<210> 38
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44004239

<400> 38
TGACGTCATC CATGTCCAAT GTCCATACCA TGGCCCCCCC AAAATGCTCT C 51

<210> 39
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44004239

<400> 39
GAAGGGATAT AACTGAAGCA ATAAATTTTT CACGGTTGGC AAATGTGGAC A 51

<210> 40
<211> 51
<212> DNA
<213> Homo sapiens

<220>

<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43957605

<400> 40
AGGACTGTTT TTCATT CAGC TTCAGCGTGA TTCCCATGGG CTCTTCTGTG A 51

<210> 41
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40915005

<400> 41
ATGTCTCAGG ATTCTACCCA AAGCCTGTGT GGGTGATGTG GATGCGGGGT G 51

<210> 42
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40356255

<400> 42
TGGCAATAAT AGTGCCTTCC TTGCTTCTTT TGCTATGCCT TGCATTATGG T 51

<210> 43
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44004667

<400> 43
CTGTGATATC TACATCTGGG CGCCCTTGGC CGGGACTTGT GGGGTCCTTC T 51

<210> 44
<211> 51

<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43068999

<400> 44
AGGGTCTGCG ACAGGGTTAC TTTGTAGAAG CTCAGCCCAA GATTGTCCTG G 51

<210> 45
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41568631

<400> 45
ATGGCCAGTG CTGGGTCTTT GCTGGAGTGA CCACCACAGT GCTGCGCTGC C 51

<210> 46
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41568631

<400> 46
GCTCTGTGGA GTCCATCAAG AATGGGCTGG TCTACATGAA GTACGACACG C 51

<210> 47
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41603916

<400> 47

GCATCCAGTG GGTAGGGGAC CCTCGTTGGA AGGATGGCTC CATTGTCATA C

51

<210> 48

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg41603916

<400> 48

TACACAACCT AGACTACAGT GACAACGGCA CGTTCACCTG TGACGTCAAA A

51

<210> 49

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg34317662

<400> 49

AGTCCCTTCT CCGTGGCACC TACGCCTATG GTTTTGAGAA GCCCTCTGCC A

51

<210> 50

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43983917

<400> 50

AGTCCTACTT TGCCATTAAC CACAATCCCG ACGCCAAGGA CTTGAAGCAG C

51

<210> 51

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)
<223> Accession number cg43983917

<400> 51
ACTTTGCCAT TAACCACAAC CCCGATGCCA AGGACTTGAA GCAGCTCGCG C 51

<210> 52
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42730678

<400> 52
TGGAGCGAGC GTGGATCCAG TTCGCTGCGG GGTGTTTGG GTCAAGTTGC T 51

<210> 53
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42714160

<400> 53
GTTACCAGAC GCTGGAGCTG GAGAAAGAGT TTCACTACAA TCGCTACCTG A 51

<210> 54
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43959084

<400> 54
TCAGGTAGCG ATTGTAGTGA AATTCCTTCT CCAGCTCCAG GGTCTGGTAG C 51

<210> 55
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele

<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42359655

<400> 55
GGGAAGCATT TGCCAATCAG TCCAGGGCGG AAAGGGATGC CTCCTGCAG G 51

<210> 56
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42359655

<400> 56
ACAGCCAGCG GTTTGGCCTG CACCATGTCA ACTTCAGCGA CAGCAGCAAG T 51

<210> 57
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42359655

<400> 57
ATCTGGTCAC CCTGCAGAAC CTGGGTGTGT CCCACTACCG TTTTCCATC T 51

<210> 58
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43998672

<400> 58
TGGTGTGGGC CTTGGTGAAC TCTAGAACGC GGCTAATGTC TCCTGGTTTG G 51

<210> 59
<211> 51
<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43922672

<400> 59

GGAAGCTGAC TCCAGAGGCC ATGCCTGACC TCAACTCCTC CACTGACTCT G

51

<210> 60

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42908571

<400> 60

GTAGTGAGGA ACAAGCCAGA GCTGTCCAGA TGAGTACAAA AGTCCTGATC C

51

<210> 61

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43942050

<400> 61

AGTTGGAAGT GAATGGATCG CAGCATTCAC TGACCTGTGC TTTTGAGGAC C

51

<210> 62

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43145505

<400> 62

TAAATATTCG AGACATTGAC AAGATTTATG TTCGAACAGG TATCTACCAT G

51

<210> 63
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43918241

<400> 63
AGATCTTTGA GGAAGGGGAA TCTGATGATG AGTTTGACAT GGATGAGAAT C

51

<210> 64
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43090990

<400> 64
TTCTGACGCA CATGTTTTGT ACATTTTCAGA CCAAGGAAAA CCTCTTTTTT G

51

<210> 65
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43969763

<400> 65
TTAGTATCAT TCACTGTGAT CTAAAGCCTG AAAATATCCT TCTTTGTAAC C

51

<210> 66
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)

<223> Accession number cg42879455

<400> 66

AGGTATATAC CATCATGTAC AGTTGCTGGC ATGAGAAAGC AGATGAGCGT C

51

<210> 67

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42659872

<400> 67

TGGCTCCGGC TACACCAACA TCATGCGGGT GCTAAGCATA TCCTGAGACG C

51

<210> 68

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42506800

<400> 68

GCTTGCCAAT TTCTCGTCTG TATGCCAAGT ACTTTCAAGG AGATCTGAAT C

51

<210> 69

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43966621

<400> 69

CTGTGGAGTA CATGTAGCTG AAGAGTCGCT CAATCTTCCT CAAGGGAACA C

51

<210> 70

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43917871

<400> 70

ACATCATATT GGCGCTGCTG ACGGGTGTAC TGCCCCCTGG CATGCTAGAT G

51

<210> 71

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43917871

<400> 71

CAGTGTAGAA ATAGGGGTGC TCCATGGCCT CTCTTGCACT AAGCCGTGAC T

51

<210> 72

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg44131752

<400> 72

AGCTCAATGG TGGCTCTGCG TGCTCATCCC GAAGTGACCT GCCTGGTTCC G

51

<210> 73

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43969473

<400> 73

AATTCAACCC ACTCATCTAT GGCAACGATG TGGATTCTGT GGATGTTGCA A

51

<210> 74

<211> 51

<212> DNA

<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44025829

<400> 74
AGACCCCGCC GTCCCCTGGC CAAGCCGTGG AGTGCTGCCA AGGGGACTGG T 51

<210> 75
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43318277

<400> 75
CTCACGCTTT GCAGTCATCT GGTCCACCTA GCACTCCCTC CTCTCCTCGG C 51

<210> 76
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43966144

<400> 76
TTAACACGAG GGAGCCTGTG ATGCTAGCCT GCTATGTGTG GGGCTTCTAT C 51

<210> 77
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42686658

<400> 77
CCCCTGTGAT CAATATCACC TGGCTGCGCA ACGGCCAAAC TGTCCTGAG G 51

<210> 78
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42686658

<400> 78
CACCACCAGA TGCCATGGAG ACCCTAGTCT GTGCCCTGGG CCTGGCCATC G

51

<210> 79
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42686658

<400> 79
CCATGGAGAC CCTGGTCTGT GCCCTAGGCC TGGCCATCGG CCTGGTGGGC T

51

<210> 80
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42686658

<400> 80
TGGTCTGTGC CCTGGGCCTG GCCATTGGCC TGGTGGGCTT CCTCGTGGGC A

51

<210> 81
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42686658

<400> 81
TGGGCCTGGC CATCGGCCTG GTGGGGTTCC TCGTGGGCAC CGTCCTCATC A 51

<210> 82
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42686658

<400> 82
TCGGCCTGGT GGGCTTCCTC GTGGGTACCG TCCTCATCAT CATGGGCACA T 51

<210> 83
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg38337333

<400> 83
GTTTCCTCAT TAGCCCTGTG ACCCCTGCAC ACGCAGGGAC CTACAGATGT C 51

<210> 84
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg38337333

<400> 84
TTGACATCTA CCATCTATCC AGGGAAGGGG AAGCCCATGA ACTTAGGCTC C 51

<210> 85
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg38337333

<400> 85
CTTCTAGTAG TTGGCCTTCA CCCACAGAAC CAAGCTTCAA AACTGGTATC G 51

<210> 86
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg38337333

<400> 86
GGTACTCAGT GGCCATCATC CTCTTTACCA TCCTTCCCTT CTTTCTCCTT C 51

<210> 87
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg38337333

<400> 87
TGGCCATCAT CCTCTTCACC ATCCTCCCCT TCTTTCTCCT TCATCGCTGG T 51

<210> 88
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43984759

<400> 88
AGGAGCTCAA GCGTGAGGCC GAGACTCTAC GGGAGCGGGA AGGCGAGGAG T 51

<210> 89
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg39660131

<400> 89
TCATGGGCAA CCTAAGGCAC AAGTGTGTGC GCAACTTCAC AGCGCTCAAC G 51

<210> 90
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44963814

<400> 90
CGGAATACCT GGCCATCACC TCTGAGAGCA AAGAGAACTG CACGGGCGTC C 51

<210> 91
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg21413267

<400> 91
AGAGTGGCGA GTGGGTCATC GTGGATGCCG TGGGCACCTA CAACACCAGG A 51

<210> 92
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg21413267

<400> 92
ACAACACCAG GAAGTACGAG TGCTGTGCCG AGATCTACCC GGACATCACC T 51

<210> 93

<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg3000465

<400> 93
AGAGGCTCTT TCTGCAGAAA CTTCCCAAAT TACTTTGCAT GAAAGATCAT G

51

<210> 94
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg30421838

<400> 94
GTCTAGGATG GAGATCCTAC AAACATGTCA GTGGGCAGAT GCTGTATTTT G

51

<210> 95
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg30421838

<400> 95
ATAACTTGCA TGATCTTGTC AAACAGCTTC ATCTGTACTG CTTGAATACA T

51

<210> 96
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43947341

<400> 96
TTACGTCGCC AAATTCCCAG GGCACGTTGC GCACGAACCTT CAGTACGGGA T 51

<210> 97
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43939230

<400> 97
TCCCTGTGAC CCAGGCAGGT GCATGGGTGA CACTGGTCGT GACCTGGCCA G 51

<210> 98
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42674136

<400> 98
CGGCACACAG GCCGCTCGCC GGAGCTGTGG CCCACCCCCA GCCCCTGGCC A 51

<210> 99
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41972699

<400> 99
AGAACTCGCG GGTCTCCAC TACATTATCA ACTCGCTGCC CAACCGCCGT T 51

<210> 100
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42849556

<400> 100
CTGCAACTAC CTTGAACCAG TTGAGTTGCG GATCCACCCT CAGCAGCAGC C 51

<210> 101
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43996195

<400> 101
CAGCATGACC TGGCACTGTA CTTGAGGAA AGTTGGGGAT TTCACCGTAG T 51

<210> 102
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43996195

<400> 102
TTGCAACTTG AGGTCGGTGC TTAGTATGAG ACAGAAGCCA TTCTGCAGTG T 51

<210> 103
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43948227

<400> 103
TTTACAGTTT TCTTACTGCA TCATCTATGT CAGAAATCTG TTCCTTCAGC T 51

<210> 104
<211> 51
<212> DNA
<213> Homo sapiens

<220>

<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43333426

<400> 104
AGAGCCACTA CAAGGTGGAC TACTCGCGTT TTCACAAGAC CTACGAGGTG G

51

<210> 105
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43051431

<400> 105
TCACACCTGT CCTGACCCTG GAGGACGGGT TCTACGAAGT TGACTACAAC A

51

<210> 106
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43920929

<400> 106
GCAGGATCAC CTGCACCCTC TTGGGGACCA TGATGCTCAT CCAGCTGTCT A

51

<210> 107
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43059041

<400> 107
AGTCAGACAC CAGCTTAGAA ATGACTATGG GCAATGCCTT GTTCTTGAT G

51

<210> 108
<211> 51

<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40148056

<400> 108
GGAGGACAGG CAACTCATCA CCGAATTAGT CATCAGCAAG ATGAACCAGC T

51

<210> 109
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42894986

<400> 109
ACCCGTTCTT CTGCCCACCC ACTGAAGCCC CAGACCGTGA CTTCTTGGTG G

51

<210> 110
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43961212

<400> 110
TCTGGAAGCC GGACATCCTC TGAGCGAGTC GACTGATCCG CTGGCGAACC A

51

<210> 111
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42898003

<400> 111

TCATCAGAGA TTCGATCTCC TCGTCAGTCA CGTGCTCCCC GGAGGCCCTG A 51

<210> 112
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43960684

<400> 112
GCTTTGAGGA GGAGGCGCGG TTGCGGGACG AACTGAGGC GGCCATCCGC G 51

<210> 113
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43958714

<400> 113
TTCGGAAAGG GCAAGCAGTG ACCCTCATGA TGGATGCCAC CAATATGCCA G 51

<210> 114
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43124627

<400> 114
ACACCCACAG CAGTTTGGT TTAGGTTTAT CTGTAAATGG AAGGTTCTGG C 51

<210> 115
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)
<223> Accession number cg43968419

<400> 115
TCTTCTCCAA CAGTCTGCCA CCCGCTGTCG TTGGCTGCGC CTCCAAGGCC C 51

<210> 116
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43064068

<400> 116
TTGTGGTCCT GGCCTCGCAG TTCCTATCCC ATGACCCAGA ACAGCTCACC A 51

<210> 117
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43064068

<400> 117
TCACAGGGAA AATTCAACGA GCCAAACTTC GAGACAAGGA GTGGAAGATG T 51

<210> 118
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41084924

<400> 118
TGACTCTCCC CGACCCGTCC CACCATGGTC TCCACAGCAC TCCCGACAGC C 51

<210> 119
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele

<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41084924

<400> 119
TCCGCAAGGC CTTCTGAAG ATCCTTCACT GCTGACTCTG CTGCCTGCCC G 51

<210> 120
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41084924

<400> 120
TCCTCGTCGC CACACTGGTC ATGCCATGGG TTGTCTACCT GGAGGTGGTA G 51

<210> 121
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43985000

<400> 121
TTGCTCTTTG CTGGTTCCT CTTCAATTAA GCCGTATATT GAAGAAACT G 51

<210> 122
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43985000

<400> 122
TATTGAAGAA AACTGTGTAT AACGAGATGG ACAAGAACCG ATGTGAATTA C 51

<210> 123
<211> 51
<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg44930578

<400> 123

ACGTGAACAC CGACATCTAC TCCAAAGTGC TGGTGACCGC CGTGTACCTG G

51

<210> 124

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg3003519

<400> 124

ATTCTTGAT TGCTAGGACC CTTTATAAAA GCACCCTGAA CATACCTACT G

51

<210> 125

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg3003519

<400> 125

CTTATGCTGT GATCATTTC A GTGGGTATCC TTGGAAATGC TATTCTCATC A

51

<210> 126

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43969010

<400> 126

TCCGAAAGAA GTCTTGGGAG GTGTATCAGG GAGTGTGCCA GAAAGGGGGC T

51

<210> 127
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43263108

<400> 127
AGACACCCCT TTCCCAGCTC GCCTCAGGGA GGAGGGACCC AAGGGCCCCC T 51

<210> 128
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43263108

<400> 128
GCCCTCGGC CTTCGCGGTG CTGGTGACCG GACTGGCGGC CACCGACCTG C 51

<210> 129
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43267238

<400> 129
CCAGACTGGT CCTGGTGGTG GTGGCGGTCT TCGTCGTCTG CTGGACTCCC A 51

<210> 130
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)

<223> Accession number cg43267238

<400> 130

CAGCACTCAC CATGGAATCC CCGATTGAGA TCTTCGCGG GGAGCCGGGC C

51

<210> 131

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43267238

<400> 131

CGATCCAGAT CTTCCGCGGG GAGCCTGGCC CTACCTGCGC CCGAGCGCC T

51

<210> 132

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43264978

<400> 132

TGGATCTGCA CCTCTTCGAC TACTCCGAGC CAGGGAACCT CTCGGACATC A

51

<210> 133

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg3001696

<400> 133

CGCTGCACCT GTGCATCGCG CTGGGTTACG CCAATAGCAG CCTCAACCCC G

51

<210> 134

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg3001696

<400> 134

TGGCTGTGAC CCGTCCCCGG GACGGTGCAG TGGTGTGCAT GCTCCAGTTC C

51

<210> 135

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42704646

<400> 135

TGGCCTTCCC GATCACCATG CTGCTGACTG GTTTCGTGGG CAACGCACTG G

51

<210> 136

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43326635

<400> 136

GGGATGCCAC CTTCTGCTTC ATCGTGTCGC TGGCGGTGGC TGATGTGGCC G

51

<210> 137

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg3003708

<400> 137

CCATCTCCTT CTGTGGCTGT CTCACGCAGA TGTATTTCGT TTTCATGTTC G

51

<210> 138

<211> 51

<212> DNA

<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg3003708

<400> 138
GGTGGAAAGC CTTCTCCACC TGTGGCTCTC ACCTGGCTGT GGTTCCTCTC T 51

<210> 139
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg3003708

<400> 139
ACAGCACCAT CATTGCTGTG TATTTCAACC CTCTGTCCTC CCACTCAGCT G 51

<210> 140
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg36729339

<400> 140
ACTCTCCAAT GTACTTTTTC CTCTCTAACC TCTCCTTCTT GGACCTCTGC T 51

<210> 141
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg38841806

<400> 141
GACATCAGGC GCACGGTGCC AACCTGCGCC ATCTGCAGGC CAAGAAGAAG T 51

<210> 142
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg38841806

<400> 142
AGGCGCACGG TGCCAACCTC CGCCACCTGC AGGCCAAGAA GAAGTTGTG A

51

<210> 143
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg38841806

<400> 143
CAGCCTTCTC CATGCCCAGC TGGCAACTGG CACTGTGGGC ACCAGCCTAC C

51

<210> 144
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43040273

<400> 144
CCTGTGCTGA TCTGGTCATG GGCCTAGCAG TGGTGCCCTT TGGGGCCGCC C

51

<210> 145
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43040273

<400> 145
CTTGCCCATC CAGATGCACT GGTACAGGGC CACCCACCAG GAAGCCATCA A 51

<210> 146
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43336100

<400> 146
TTGGAAGCGT GCATCCAGTG AGACCTATGA GGCTTGAGTC TTTTAGTGCC T 51

<210> 147
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg21646034

<400> 147
GTGTGAGCAG AGATGCCAGA ACCAAGGTGG ACCGAACACC ATTACATATG G 51

<210> 148
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43916882

<400> 148
TGGCAGCTAC CAGCACAATG CCTCCGCCGT CAATAAAGGC ACTGATGGTC T 51

<210> 149
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2537639

<400> 149
TGGCTCCCAT TGTCTGGGAG GGCACGTTCA ACATCGACAT CCTCAACGAG C 51

<210> 150
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2537639

<400> 150
ACGTGGACAT GGAGTTCCGC GACCATGTGG GCGTGGAGAT CCTGACTCCG C 51

<210> 151
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2537639

<400> 151
ACGTGGGCGT GGAGATCCTG ACTCCACTGT TCGGCACCCT GCACCCCGGC T 51

<210> 152
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2537639

<400> 152
GGCCCCAGTC CCAGGCCTAC ATCCCTAAGG ACGAGGGCGA TTTCTACTAC C 51

<210> 153
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2537639

<400> 153
ACGAGAGCCA CCTGAACAAG TACCTACTGC GCCACAAACC CACCAAGGTG C 51

<210> 154
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44000740

<400> 154
GGGGAGATAC TGGCTCACCC AGGAACACAG GGAACATCAC CTTATGCCAC A 51

<210> 155
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg38869466

<400> 155
ACGCAGTGGC CGTGGGCTCC CTCTGTGCTC TTTCCGCCAG TCTTCTAGGT T 51

<210> 156
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40351913

<400> 156
CCATCGCCAC GCTCCCTCTG TCCTCGGCCT GGGCCGTGGT CTTCTTCATC A 51

<210> 157

<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43964039

<400> 157
GATGGAACAG CTCCTCGGGT GTCTTATCAC TTTGGCTGGC TCCCCCCTGC C

51

<210> 158
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43992017

<400> 158
GCGGCTGCTG GTGGATGGGT GGGCGGGGGG TGCAGCCTCC ACCCCCTCCC C

51

<210> 159
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43948629

<400> 159
CGCCTGTAAT GGCTGTGAAC ATGCTTACCC AGCAGGAGGT CCCTGTCGTT A

51

<210> 160
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43955093

<400> 160
CATTGACTAG GGGCTGTGGG GGCATGCGCC CAGGTGTCCC TCCATCAGAG G 51

<210> 161
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43955093

<400> 161
AGCAGGCCAA GAGAGATCTG TGGAATGCAT CTTGTTCCAG AATACCAGAT A 51

<210> 162
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43055918

<400> 162
CGCTGGCATA GGACATGGCG GGCTTTCCCC CCGCAGAGCT CTGGGGGCTA C 51

<210> 163
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43974592

<400> 163
AAATAACAAG GCATTGAAGA ATGGCAGACG AGCGGAAAGA CGAAGGAAAG G 51

<210> 164
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956384

<400> 164
AAGGACGCAA CGCTGCCACC ATGGATAGTA GCACCTGGAG CCCCAAGACC A 51

<210> 165
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44025634

<400> 165
TGAAAGTATT CAATCCCAGA AGGAAGCTGG AATTGCCCCT TCTGTTTCTA G 51

<210> 166
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43940037

<400> 166
GCTGGCGCAC TGCTAGCCTC AGAGGAGCCA GCACCTCCTC AGCCCCCGCG C 51

<210> 167
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44024279

<400> 167
AAAACCAGCT ACCTGCCTTT CTGGAGGAAC TTTGCCATGA GAAAGAAATT T 51

<210> 168
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43926814

<400> 168
CATGAGTTT GATCCCAGCT CTCTTTTCCC TGGCTTTCTG GGCCATTCT C 51

<210> 169
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40918088

<400> 169
TTGGAGCTGG AATTACTGTG TATGAGGCCT TAGCAGCTGC TGATGAGCTT T 51

<210> 170
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43966985

<400> 170
TCAACACCTA CGTCCACTTC CAAGGTAAGA TGAAGGGCTT CTCCTGCTG G 51

<210> 171
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43924009

<400> 171
TGGCTGAC AAATTGCTTG AAGACTCGAT CCATGTAAGT GGACTGTCTT G 51

<210> 172

<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42913861

<400> 172
CTGGGCAGCT GCCCTCACAG TAGTTGCCGT AGTAGCCGGT GGGTGCTATG A

51

<210> 173
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42913861

<400> 173
GCCGAGCCTG CACCACCACA AAGGGTCGGT GCGACTCTTC GCCTGGGTCC A

51

<210> 174
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43929685

<400> 174
CATAGAAGGC CAGGAGTCAG GAGACTTGGG TTCTGTCCTG GATTATACAC C

51

<210> 175
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43929685

<400> 175
GGAGTCAGGA GACCTGGGTT CTGTCTTGGA TTATACACCA GCTCACTGAG G 51

<210> 176
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43918561

<400> 176
CCATGCCAC CCCCAGACGCC ACCACCCAC AGGCCAAGGG CTTCCGCAGG G 51

<210> 177
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42343176

<400> 177
ATTTAATGAA TTTCCTGAAG ACTGTGAGAA GTACAACTGA GAAATCCCTT T 51

<210> 178
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956382

<400> 178
AAACAATGA TATCGATGAA GTTATTATTC CCACAGCTCC CTTATACAAA C 51

<210> 179
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43984681

<400> 179
CACCATGAAG CAGTTGCTGC GGGCCTTGGA GGAGGGCCGC GTGCGGGAAG T 51

<210> 180
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43984681

<400> 180
TCCTGTACAA AGACAGGAAC CTCCATATTC CCACCATGGA AAATGGGCCT G 51

<210> 181
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43950996

<400> 181
CTGCGGTGGA GACGTCAGAG CTGCCGGGGG AGGGGGCTCC TGC GCCACAG C 51

<210> 182
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44024506

<400> 182
ACCAGTGCT CGTAGTACAC AGGCAAGCAC TTCTCCTTGC CTACCTCCAT G 51

<210> 183
<211> 51
<212> DNA
<213> Homo sapiens

<220>

<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43980381

<400> 183
CTACCGCCAA CTATGACTTT GTCCTGAAGA AGCGGACCTT CACCAAGGGA G 51

<210> 184
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42650960

<400> 184
TCCCCTGGCA GAACTACCAC CTGAATGACT GGATGGAGGA GGAATACCGC C 51

<210> 185
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43249389

<400> 185
ACATCCAGGT GGTGTTTCGAC GCCGTTACCG ACATCATCAT TGCCAACAAC C 51

<210> 186
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43946951

<400> 186
CAGTGACGGC AGGGTCAAAG TCCTTAGCGT AGCCCTCGTT AAGGCTGTAG A 51

<210> 187
<211> 51

<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43248117

<400> 187
AACCAGCCCA CTGTGAGAAG ACCACCGTGT TCAAGTCTTT GGAATGGCA G 51

<210> 188
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44027049

<400> 188
CCACAATGTT AGGAGGGTAT TTTTATATCC CTCCAGTTAA CAAATACAGC A 51

<210> 189
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43982075

<400> 189
CTGCCATCTT TCAGCCCTCT GAAACTGTGT CCAGCACAGA ATCTTCCCTG G 51

<210> 190
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43942977

<400> 190

CGCCTTCCCA GGTCCGGACA ATTCGTCAGA CTATTGTCAA ACTGGGGAAT A

51

<210> 191
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43942977

<400> 191
GGCAGCTGAA GATCACCAAT GCTGGCATGG TGTCTGATGA GGAGTTGGAG C

51

<210> 192
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43942977

<400> 192
GCGAGGTGTT TGTGTCCAAT ATCCTTAAGG ACACGCAGGT GACTCGACAG G

51

<210> 193
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43985220

<400> 193
TGTACACTGC CAGAAAAGGA AAAGGGGCCT TTTGTAATGG TCAAAAAC TA C

51

<210> 194
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)
<223> Accession number cg43985220

<400> 194
TCTTGGTGAC TGAGTTGGGC TCTTCTAGAA CACCAGAGAC TGTGAGAATG G 51

<210> 195
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43946394

<400> 195
TAGAGCGCAC ACAGGCCTCC AGCTGGGCCA TGTCGGTCTC ATCATCCCAA G 51

<210> 196
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43119818

<400> 196
CTGACAGCTA CAGGCTCTTT CAGTTTCATT TTCACTGGGG CAGTACAAAT G 51

<210> 197
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43118279

<400> 197
AGAAGTTGAA GGGGCTGGTG CCACTGGGAC CCGAATCAAG TCGACACACT A 51

<210> 198
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43325007

<400> 198

TGGGTTTCAGG GATGTAGCCC TTCTCCACAG CCAGGCGGCT CAGGGCAAAC A

51

<210> 199

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg39524111

<400> 199

GCCAAATATAG GATAGGGCAC TACAGGTTCC GGTACAGTGA CACCCTGGAG C

51

<210> 200

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43280516

<400> 200

ACGGGGAGGA GCTGCAGATG GAACCTGTGT GAGGTGTCTT CTGGGACCTG C

51

<210> 201

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43963913

<400> 201

AGAGGTTGGG GGGCGCCGAG CGCGAACGGC CCCGAAAGGG GCTGGGCTCC T

51

<210> 202

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg40262905

<400> 202

TAGTGAAAGG CCTGAAATAT ATGCTCGAGG TGGAAATTGG CAGAACTACC T

51

<210> 203

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43918168

<400> 203

CTCCATCAAC AGCATCCGGA CTGCACGGCG GCTCGCCGTG CGGCTGGGGC C

51

<210> 204

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43259701

<400> 204

CGACGAGGTG CTACGCGAGG GCGAGTTGGA GAAGCGCAGC GACAGCCTCT T

51

<210> 205

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg1527767

<400> 205

TTTTTCCAG CTTACAATGG TACAGGCAGG AGCCTGGGGA AGGTCCTGTC C

51

<210> 206
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40968986

<400> 206
AGAAGAGAGC CTGTGACACT GCCACTTGTG TGACTCATCG GCTGGCAGGC T

51

<210> 207
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42550133

<400> 207
TCATCTGAG TTCTAAGAAG CTCCTCCTCA GTGACTCTGG CTTCTATCTC T

51

<210> 208
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2526759

<400> 208
CTCTGGTTGT CCACGAGGGA GACACCGTAA CTCTCAATTG CAGTTATGAA G

51

<210> 209
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)

<223> Accession number cg41664708

<400> 209

AAGTCTGTGC TGATCCACAA GCCACGTGGG TGAGAGACGT GGTCAGGAGC A

51

<210> 210

<211> 50

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (25)...(26)

<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43300673

<400> 210

AGGGAGGCGG GGAGGGTAGC ATGGGCACAC GGCCCTCACA GGGACTCACT

50

<210> 211

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43284434

<400> 211

AGTTGAAATC AGAGAGGAAT AAAAAAGACA TTTTATATTT TATTCTGCTC C

51

<210> 212

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43132502

<400> 212

TAAGCATGAG GTGGCACGAG GCAGGCGTTG GCGATGCCAC CTGGGGGTCA C

51

<210> 213

<211> 51

<212> DNA

<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43931765

<400> 213
GGTCCCTTG CTTTATCCCA AGCTCTGAGG GACGCAGCCT GGCATGGCTC T 51

<210> 214
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43931765

<400> 214
GTCCCTTGC TTTATCCCAA GCTCGTAGGG ACGCAGCCTG GCATGGCTCT G 51

<210> 215
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43931765

<400> 215
CTTTATCCCA AGCTCGGAGG GACGCGAGCC TGGCATGGCT CTGGCCTAGC A 51

<210> 216
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43931765

<400> 216
TAGCAGCCAG GTGACATGGC CAGGCTACCT TCCTGTACAG GCACTGTGGG C 51

<210> 217
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43931765

<400> 217
GCCAGGTGAC ATGGCCAGGC ACCTTTCCTG TACAGGCACT GTGGGCTCCT G

51

<210> 218
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43931765

<400> 218
AGGTGACATG GCCAGGCACC TTCCTTGTAC AGGCACTGTG GGCTCCTGGC C

51

<210> 219
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43931765

<400> 219
AGGTGACATG GCCAGGCACC TTCCTTGTAC AGGCACTGTG GGCTCCTGGC C

51

<210> 220
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43931765

<400> 220
AGGTGACATG GCCAGGCACC TTCCTTGTA C AGGCACTGTG GGCTCCTGGC C 51

<210> 221
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43952088

<400> 221
AATCCACAAT CGGCATCAGG AAGCCCAAGT CCCAGTGGCC ATTAGGGTCC T 51

<210> 222
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44010957

<400> 222
TCCCTATGAG CCTGCAAAGG AGACATTCAG GAATGAGTTC CATGTTGAG A 51

<210> 223
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956560

<400> 223
CAGTGCATCT GGGAAGATTT CTACCCGACC AACAGTTCCT TCAGCTTCCA T 51

<210> 224
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956560

<400> 224
CAACAGTTCC TTCAGCTTCC ATTTACCCCC TCATTATCC CTCAACCCCC A 51

<210> 225
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956560

<400> 225
TGCTCTCCTT TCCCCTGCCC CCAGAACTTT TATCCACTTA CCTAGATTCT A 51

<210> 226
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43264626

<400> 226
TGGCCACAGT GAAAAAGGTC ATGGGAGGAG AGAAGCAAAG TAGGAAGGAT C 51

<210> 227
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43011543

<400> 227
ACCGCACCTT TTCCACCGGT GGGGGGCCCA GTGAAGTTTA ACAAACTGCT G 51

<210> 228
<211> 51
<212> DNA
<213> Homo sapiens

<220>
 <221> allele
 <222> (26)...(0)
 <223> single nucleotide polymorphism

 <221> misc_feature
 <222> (0)...(0)
 <223> Accession number cg43011543

 <400> 228
 CATACCACGT TCACTGCAAG GGGGGGAACG TGTGGGTTC TCTATTCAAG A 51

 <210> 229
 <211> 51
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> allele
 <222> (26)...(0)
 <223> single nucleotide polymorphism

 <221> misc_feature
 <222> (0)...(0)
 <223> Accession number cg43933757

 <400> 229
 GAAACCCAGT AGGCTCCTGG AGGCCCTGGT CAGCTTGCTT GGAATCCAGC A 51

 <210> 230
 <211> 51
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> allele
 <222> (26)...(0)
 <223> single nucleotide polymorphism

 <221> misc_feature
 <222> (0)...(0)
 <223> Accession number cg41553795

 <400> 230
 TGGTGGTGCT ACCCTTGGCC TCCCAGAGTC CTGCCACCCT GCTGCCGCCA C 51

 <210> 231
 <211> 51
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> allele
 <222> (26)...(0)
 <223> single nucleotide polymorphism

 <221> misc_feature
 <222> (0)...(0)
 <223> Accession number cg42542496

 <400> 231
 AGCCCTTCTC CACCCGATA GATTCTTCAC CCTTGGCCCG CCTTTGCCCC A 51

 <210> 232

<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42542496

<400> 232
ACCCGATAG ATTCTCACC CTTGGTCCGC CTTTGCCCCA CCCTACTCTG C 51

<210> 233
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41533258

<400> 233
GTGCCTGGAC ATTTGCCTTG CTGGATGGGG ACTGGGGATG TGGGAGGGAG C 51

<210> 234
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2753430

<400> 234
ACGACTTTGA GCCTCGCGAT CTTTGTGAGTC CAACGTCCAG CTCGTTCTCT G 51

<210> 235
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44036323

<400> 235
TGCGGCTTAA AAGGGCAACC CGCGCCGGAC CCTTCCTCCC TAGTCGCGGG G 51

<210> 236
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43918671

<400> 236
GAGAGACCAT TTAATTACAT CAGTTTGGTT TATAGACATT TGAATCATAT C 51

<210> 237
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43057018

<400> 237
AGTTTCATTA TACTTTTCTC TCCACGTTTT GTCTATGTTG AAAATTTTCT G 51

<210> 238
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44005808

<400> 238
ACAAGACAGA AGCTGAAGTG CATCCCAAAG GTGCTCAGAG AGCCGGCCCG C 51

<210> 239
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44005808

<400> 239
TCTTCCCTTCT CCAGCCGGCA GGCCCGCGCC GCTTAGGAGG GAGAGCCCAC C 51

<210> 240
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956159

<400> 240
TGGCGAGTCC AGGGTCACCC ACATAACCAT GCACCACGGG TGCTATGCCG C 51

<210> 241
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956159

<400> 241
GAGTCCAGGG TCACCCACAT ACCATTGCAC CACGGGTGCT ATGCCGCTTC T 51

<210> 242
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956159

<400> 242
TACCATGCAC CACGGGTGCT ATGCCACTTC TTACAGGACC TTTTtagccc T 51

<210> 243
<211> 51
<212> DNA
<213> Homo sapiens

<220>

<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956159

<400> 243
CCTGGAGGCA ACTGGGTAGG GTGCACAACG GCATGCTTTG GCTGGAACAC G

51

<210> 244
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956159

<400> 244
CAGAACGGCA TGCTTTGGCT GGAACCACGC ATCCCTCCTT CCACGGCCGG C

51

<210> 245
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43971258

<400> 245
CAGAGCTAGC TCTGGCTCTT CAGGCTACAA GTTCACAGTC CTTGCTCCT G

51

<210> 246
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43971258

<400> 246
GTCTTCGCT CCTGAGCACC AGGTTCACTC TCCAGGAAGG GATTGCTGA A

51

<210> 247
<211> 51

<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43982507

<400> 247
GATACCTTTG CGTGGATCAA GCTTGCTGTA CTTGACCGTT TTTATATTAC T

51

<210> 248
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43982507

<400> 248
CAAAAAAATT TATAAACTAA TTTTGGTACG TATGAATGAT ATCTTTGACC T

51

<210> 249
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43982507

<400> 249
CCTCCTTCTC CCCCTTTCCC CTCCCCGCCC CCACCTTCTT CCTCCTTTCG G

51

<210> 250
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41554010

<400> 250

CTGCCCTGCC ACCTGTCTGT CTGTCTCCAA AGAAGTTCTG GTATGAACTT G 51

<210> 251
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41554010

<400> 251
CTGCCCTGCC ACCTGTCTGT CTGTCTCCAA AGAAGTTCTG GTATGAACTT G 51

<210> 252
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43984905

<400> 252
TCCCTCCAGG ACTAGGCTGG AGGAACCCAG TGGGGTCCCC CCTGAGTGGG C 51

<210> 253
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43984905

<400> 253
CACATGTGGG GACAGGGCTG GTGTGCCTGC TCCAGCCTC TTGCTCAGAG C 51

<210> 254
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)
<223> Accession number cg43992911

<400> 254
CTAAAGTCGG AGTATCTTCT TCCAAAATTT CACGTCTTGG CGGCCGTTC A 51

<210> 255
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43932434

<400> 255
TTTCTAGAGG GGGTCTGTTG AAGATATGTA ACTAGTACAC CCCAACCCCC A 51

<210> 256
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43932434

<400> 256
CCCCAACCCC CAACCTCAGT GGAAAGCAAT GCCCAGGGAT TAGGCTATGG A 51

<210> 257
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43318219

<400> 257
GCGCAGGTCA GAGGGCGGCC GCAGCGGGCC TCCGCGAGGT CCCCACGCC G 51

<210> 258
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele

<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43967861

<400> 258
CTCTATACTG TACACTCACC CATAATCAAA CAATTACACC ATGGTATAAA 50

<210> 259
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43967861

<400> 259
TCTATACTGT ACACTCACCC ATAATCAAAC AATTACACCA TGGTATAAAG 50

<210> 260
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43965366

<400> 260
GCCGAATAGC CTGGGTTTGG AAAAGTATGT TTTTGAAATA TGTGGGATCT C 51

<210> 261
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)

<223> Accession number cg43965366

<400> 261

TACTGACCTA AATCACACCC TAGACTTATC AGAGGGAAAT TCTGACCATA A

51

<210> 262

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43322513

<400> 262

TGTCCTTGAA GAACATGCAC TTGGCGCGGA TGGCACAAGC AAAATGGTAG A

51

<210> 263

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg41637704

<400> 263

CCGCGCCCC AGTAGGAGCC CCGCGGCCCA GCAGGTGCGG CGCGCACGGA G

51

<210> 264

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg41637704

<400> 264

CCAGCAGGTG CGGCGCGCAC GGAGCGCGCC GGCCGGCGGC TTCTCCCGGA G

51

<210> 265

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg41637704

<400> 265

TGAAACTTGA AACCGCCTCT GGAGCTGCCA TTCTGCAGAG TATTTGGAAA A

51

<210> 266

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43980506

<400> 266

TCCAAGAAAG GGTCATGGAA GCTTACTGGG AATAATCCTC TCAATTAGAA A

51

<210> 267

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43961305

<400> 267

GGGGGGTTTT TTTTTTTTTT CTCTGTTTTT TTTTTTTTTT TTTTTTTTTT T

51

<210> 268

<211> 50

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (25)...(26)

<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43998672

<400> 268

CTGGGGGTTT TCGGGGAGGA ACCAAGGCTC ACGGAGCCTC CTGTGCTGCA

50

<210> 269
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43998672

<400> 269
GGGGGTTTTC GGGGAGGAAC CAAGGCTCAC GGAGCCTCCT GTGCTGCAGT

50

<210> 270
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42908571

<400> 270
GAGTTAATTT ATGTAAGTCA TATTTTATAT TTTTAAGAAG TACCACTTGA A

51

<210> 271
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42908571

<400> 271
CTTACCTCAA ATAAATGGCT AACTTTTATAC ATATTTTAA AGAAATATTT A

51

<210> 272
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42164914

<400> 272
CAGCCCCCAT TGTGGTCACA GGAAGCAGAG GAGGCCACGT TCTTACTAGT T 51

<210> 273
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43958501

<400> 273
CCCAACCTGG GTTTGGCAGA CATCAGAATG ATGGAGTACA TTTTGCAGAT A 51

<210> 274
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43958501

<400> 274
CACCCCCAGG TTCTCCTAGT TCAGAAAAAA GCTGTGAAAG TGAAGAAGG A 51

<210> 275
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43090990

<400> 275
TTTATTCTAT TCCTATCTGT GGATGGGTAA ATGGCTGGGG GGCCAGCCCT G 51

<210> 276
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42879455

<400> 276
AGCCTTTGTG CTCCCACTCA ATACACAAAG GCCCCTCTCT ACATCTGGGA A 51

<210> 277
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42879455

<400> 277
AAAAAGGCCC CTCTCTACAT CTGGGGATGC ACCTCTTCTT TGATTCCCTG G 51

<210> 278
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43971741

<400> 278
AGCAACTTGG CTGAGCCCCA CTACATACAG AGAAATCATC AACCTGACTT A 51

<210> 279
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43971741

<400> 279
TAAGAGTTTT CAAGATGTCA AACTTAAGGC TGATCAGCAG ATGGGATGTG A 51

<210> 280

<211> 51
 <212> DNA
 <213> Homo sapiens

<220>
 <221> allele
 <222> (26)...(0)
 <223> single nucleotide polymorphism

<221> misc_feature
 <222> (0)...(0)
 <223> Accession number cg43971741

<400> 280
 TTTTAAAAA TCCATCCACA CACATTGGTA AATTAAGTAT AAATTCCTTT G

51

<210> 281
 <211> 50
 <212> DNA
 <213> Homo sapiens

<220>
 <221> allele
 <222> (26)...(0)
 <223> single nucleotide polymorphism

<221> misc_feature
 <222> (25)...(26)
 <223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
 <222> (0)...(0)
 <223> Accession number cg43947749

<400> 281
 AACGTCGATT CGCACCCTCC AACCTGCCCC GCCCTCCTA CAGCTGTAAC

50

<210> 282
 <211> 50
 <212> DNA
 <213> Homo sapiens

<220>
 <221> allele
 <222> (26)...(0)
 <223> single nucleotide polymorphism

<221> misc_feature
 <222> (25)...(26)
 <223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
 <222> (0)...(0)
 <223> Accession number cg43947749

<400> 282
 ACGTCGATTC GCACCGTCCA ACCTGCCCCG CCCCTCCTAC AGCTGTAAC

50

<210> 283
 <211> 51
 <212> DNA
 <213> Homo sapiens

<220>

<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44131752

<400> 283
CACTTAATAC CAGAGACCCC CCCCCCTTCC CCTCCCCCTT CCCCTCCCCC T 51

<210> 284
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43917718

<400> 284
AGACGTGTCT GCCACAGGTC TCAGGGTAAC AGATGCCCTG TCCACTGAGA G 51

<210> 285
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43928048

<400> 285
TTTGATGGAA AGGTTGTCCA CACTGAGAAT TATCACACAC TTGATCAGGA A 51

<210> 286
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42714751

<400> 286
CCCTCCGGAT TCGGCGCGCG TGCGGMCCGC CGCGAGTGAG GGTTTTTCGTG G 51

<210> 287
<211> 51

<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43322545

<400> 287
TCCAAGCTAA GCACTGCCAC TGGGGGAAAC TCCACCTTCC CACTTTCCCA C

51

<210> 288
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43322545

<400> 288
CCACCTCCAT CCCAGACAGG TCCCTGCCCT TCTCTGTGCA GTAGCATCAC C

51

<210> 289
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43322545

<400> 289
CACCTCCATC CCAGACAGGT CCCTCGCCTT CTCTGTGCAG TAGCATCACC T

51

<210> 290
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43980494

<400> 290

GTCTGATAGA AGAGGAGCAG GAGAAGCAAA TCGTTAAAC CTAGCGAATT C 51

<210> 291
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43925424

<400> 291
TCAGGAGCAA GGCGAATGTA TGACACCATG TCCACAATGG TGTACATAAA G 51

<210> 292
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42479188

<400> 292
TTCTGAAGAG GCTGACGATT TTAGTGCTC ATTTTTTTCC TTTCTCCAGA A 51

<210> 293
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42686658

<400> 293
CTAGCTTCCC TTCCCATTCA ACACACACAC ACATTCTTGC TCTACCCAAA G 51

<210> 294
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)
 <223> Accession number cg38337333

<400> 294
 TGTCTCAAAC CCAGCTTGCC AGCTCCAATG TACCAGCAGC TGGAACTCTGA A 51

<210> 295
 <211> 51
 <212> DNA
 <213> Homo sapiens

<220>
 <221> allele
 <222> (26)...(0)
 <223> single nucleotide polymorphism

<221> misc_feature
 <222> (0)...(0)
 <223> Accession number cg27803682

<400> 295
 GTGGCTGGGC TATTCCATCC ATCTGGAAGC ACATTTGAGC CTCCAGGCTT C 51

<210> 296
 <211> 51
 <212> DNA
 <213> Homo sapiens

<220>
 <221> allele
 <222> (26)...(0)
 <223> single nucleotide polymorphism

<221> misc_feature
 <222> (0)...(0)
 <223> Accession number cg21413267

<400> 296
 CGAGCGGCAC CCAGAGCCTG CACCCGCCCT CACCGTCCTT CTGCGTCCCC C 51

<210> 297
 <211> 51
 <212> DNA
 <213> Homo sapiens

<220>
 <221> allele
 <222> (26)...(0)
 <223> single nucleotide polymorphism

<221> misc_feature
 <222> (0)...(0)
 <223> Accession number cg21413267

<400> 297
 AGGAGCCCTC TTCGGTGTC CCGAGCGCCA CGGTCAAGAC CCGCAGCACC A 51

<210> 298
 <211> 51
 <212> DNA
 <213> Homo sapiens

<220>
 <221> allele

<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg21413267

<400> 298
CGGTCAAGAC CCGCAGCACC AAAGCGCCGC CCCCACACCT GCCCCTGTCG C 51

<210> 299
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42481172

<400> 299
GAGCCGTGTG GCTGTGGCCT CCGGGCGGCG GTGGACGGCG TCGCCTTCAT C 51

<210> 300
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg39518465

<400> 300
GGGTGCACGG CCGGCCCTGG GCAGGCGTAG CCATGGAGCT GTGGCGCCAA T 51

<210> 301
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41972699

<400> 301
ATGGGGCCCG TGTCTCGCCA GGAGGCGCAG ACCCGGCTCC AGGGCCAGCG C 51

<210> 302
<211> 51
<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg40333812

<400> 302

AGCATTTGAG GAAGCATAAC TGACGTGTGA AGGGGGTGTG GGGTACTTGC C

51

<210> 303

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43280482

<400> 303

AGCATCTGCA GACGACCCCC GCAGCCTTTC CCTCGGACCC CCCTCGAAGC C

51

<210> 304

<211> 47

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (22)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg44014837

<400> 304

CACTGCTGTG CAGGGCAGGG ATGCTCCAGG CAGACAGCCC AGCAAAG

47

<210> 305

<211> 50

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (25)...(26)

<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg41626506

<400> 305

CAGCACAGCG AGCGCTCTCA TTCTGCCTTT TTCTCTCTTC TCAGCCAACT

50

<210> 306

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43918944

<400> 306

TCTGTAGAGC TCTGAAAAGG TTGACGATAT AGAGGTCTTG TATGTTTTTA C

51

<210> 307

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43988365

<400> 307

GACAGACGAG ACAGTGAGGT ATGTGGGGCT GCTCCGGAAT GGTCCGGAGG C

51

<210> 308

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43969460

<400> 308

TAAGTATGCA AGACAAGACT TGGTCGTCAC GTTCGCGTCT CTAGTTGATT T

51

<210> 309

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43933809

<400> 309

AATTAAAACT CTAGGTGTAT ACTTACATGG AACTAGTTTA TTCCTATTT A

51

<210> 310

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43931444

<400> 310

TGCTCGCGCC GTGCCACTAA GGTCATTCCC GCCTCCGAGA GCCCAGAGCC G

51

<210> 311

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42937321

<400> 311

CTTTTCCCTC TTACCCTCTC TCTCTTAACA TCGTAAACAA CAGACTTACG T

51

<210> 312

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42937321

<400> 312

CCTCTTACCC TCTCTCTCTG AACATTGTAA ACAACAGACT TACGTTAAAC T

51

<210> 313

<211> 51

<212> DNA

<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40991963

<400> 313
CAAAATGTAA CAGTGGCTTT TCAACGGGAG TAAAGCAAAG TCTCTAAAGC T 51

<210> 314
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43951366

<400> 314
AAAGATGTTT GAATACTTAA ACACTATCAC AAGATGGCAA AATGCTGAAA G 51

<210> 315
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43951366

<400> 315
TGGTGGAGCC ACTGCAGTGT TATCTCAAAA TAAGAATATT TTGTTGAGAT A 51

<210> 316
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43306254

<400> 316
CACTTAACTT GCATGTGCAC AGCTTCTGGT AACAAATATC GCTAAACCTT A 51

<210> 317
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43306254

<400> 317
ATGTGATTAA TTATGTGATG AAAACTTTTT TTATAAATGA TCTTGGTCTA T 51

<210> 318
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42918089

<400> 318
CAATCAGAAT TGATAAGCAC TGTTCTTCCA CTCCATTAG CAATTATGTC A 51

<210> 319
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44032168

<400> 319
TCCATCCCTC TTTTGGGCTC TTCTGCAGGG AAGTAACATT TACTGAGCAC C 51

<210> 320
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43154190

<400> 320
TACCCGGAAG TTGAGCTCAA TTTCATTCT GTTTCTGGC CACAAGTGGC A 51

<210> 321
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43927549

<400> 321
CCCAGTCCTG CGGCTCCTAC TGGGGCGTGC GCTGGTCGGA AGATTGCTGG A 51

<210> 322
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43927549

<400> 322
TACTGGGGAG TCGCTGGTC GGAAGGATTG CTGGACTCGC TGAAGAGAGA C 51

<210> 323
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43927549

<400> 323
CGGTCCGTGG TCCCCGGGGG CGCAGGTCGC AGCGTCCCG CCTCCAGGC G 51

<210> 324
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43947066

<400> 324
TTCTCAAAAG GCTGGGGGTA TTTATGTAAG AACTTATTCC AAAGTGACTC T 51

<210> 325
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43923264

<400> 325
AGGAAAGCCG GAGAATTGGG GCACGAAGAG GGGGGGCTTT GATGACCCGC 50

<210> 326
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43942332

<400> 326
AGATTCATCA GAATAGGATT TTTGCCAAAT CCCACCCATA TGCTGTTGAG C 51

<210> 327
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele.
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43274705

<400> 327
CCGCTGTCTC TGTCTTCGCT TTTTATTCAA GAAGAATAAT GCGACGAAAA T 51

<210> 328

<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42207316

<400> 328
CCACTTCTCT GGGACACATT GCCTTTTGTT TTCTCCAGCA TGGCCTTGCT C 51

<210> 329
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43927885

<400> 329
CATCATCATC ATAGTTTACT TCAGCTCTTA AATCCCCGAG GAGTCTGCCC T 51

<210> 330
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40388639

<400> 330
CTCTTGCCCA GCCGGCTGCA AGTTTGTAA GCGCGGGACA GACACTGCTG A 51

<210> 331
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43949316

<400> 331
AGGTTACCAA ACAGGAATAC AACACTTCTC TCCCTTTTCT GCTCTAGAAG G 51

<210> 332
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43958714

<400> 332
TGGGTGATGA TCACTGTGCT GCTTGCGGCT CATGGCAGAG CATTCAAGTGC C 51

<210> 333
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43275028

<400> 333
ACAGACTGGC TGCAGCATTA GGAATTAGGT CATTCCGAAA CTCATCATTG A 51

<210> 334
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43275028

<400> 334
GGTCATTCCG AAACATCATCA TTGAACCAGG AAGAAGAAGA GTTCAATCTT A 51

<210> 335
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43275028

<400> 335
AGAATGCAC TGAATTCGT TCTTCGAACA CAGATATAAT TGTGGTTCA A 51

<210> 336
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43275028

<400> 336
CTTTCACTTG GTGCTGGAGA ATTCAGAAGT CAAGAACATG CTAAGCATAA G 51

<210> 337
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43275028

<400> 337
TTCAAAAGTC AAGAACATGC TAAGCGTAAG GGACCCAAGG TAGAAAGAGA T 51

<210> 338
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43275028

<400> 338
TTCTCCTTCC AGAATGAGGC CCTGGGAGGA CCCTCCTAGT GATCTGTTAC T 51

<210> 339
<211> 51
<212> DNA
<213> Homo sapiens

<220>

<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43275028

<400> 339
ACTACATAAG GACAGCAACA TGCCTGTGGA CATGAGAGAA TTTGTCTTAC T 51

<210> 340
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43985000

<400> 340
GAAAAAATC ACAAGGCAAC TGTGAGTCCG GGAATCTCTT CTCTGATCCT T 51

<210> 341
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg39565524

<400> 341
TCCGACCCA CACACCCTGA GGGAGGCCTA CCCTAGCCTC AGCCGCTCCT G 51

<210> 342
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43306266

<400> 342
ATAATCCATG CCTCTGAATA TTAGAGTGGT TTCTTGGATG GGATTTTGAA T 51

<210> 343
<211> 51

<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43306266

<400> 343
GGGATTTTGA ATATGCATTT AAGAACGTTG GGAAGAATTT CACAGATGAT G 51

<210> 344
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43306266

<400> 344
GAATATGCAT TTAAGAAGTT GGAACAATT TCACAGATGA TGATTGGAGG A 51

<210> 345
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43329467

<400> 345
TCGGCAAATC TTGAAAGCTG CAGGGTGCAG AGACATGGAT GTGACTTCCC A 51

<210> 346
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2751286

<400> 346

AAGGCATAAG AACTAGGAGC TGCTGGACAT TTCAATATGA AGGGCAACTC C 51

<210> 347
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg11751407

<400> 347
GAATGTGGGG ATAAGGCATT GGGACTCTAT CAGGTATCCT GAGGAGAGAC T 51

<210> 348
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43326635

<400> 348
CAGCCGGGAG CTCTGCCAGC TTTGGTGAAG GAGGGTGCTT GCCTCGTGCC C 51

<210> 349
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43326635

<400> 349
CGGGAGCTCT GCCAGCTTTG GCGAACGAGG GTGCTTGCCT CGTGCCCCTT G 51

<210> 350
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43993798

<400> 350

TGCTCTTGCT GCTGATGGAG GAGGAGGGGG TGGATCCCGT GGAGCCTCCA A

51

<210> 351

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43040271

<400> 351

ATGCTTCCCC CAACCCTAGG GAATCCACAC TTAAGATAAT TCGCCACTTC T

51

<210> 352

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43040271

<400> 352

CCAACCCTAG GGAATCAACA CTTAATATAA TCGCCACTT CTCCTCTTTC T

51

<210> 353

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43040271

<400> 353

AGATAATTCG CCACTTCTCC TCTTTTCTC TGCTCCGCTC ACGGCTTGCA G

51

<210> 354

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43040273

<400> 354
CGCAGAGCCC CGCCGTGGGT CCGCCCGCTG AGGCGCCCC AGCCAGTGCG C 51

<210> 355
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43040273

<400> 355
CAGCGCCTTC TTGCTGGCAC CCAATGGAAG CCATGCGCCG GACCACGACG T 51

<210> 356
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43040273

<400> 356
TGCGCCGGAC CACGACGTCA CGCAGGAAAG GGACGAGGTG TGGGTGGTGG G 51

<210> 357
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43040273

<400> 357
GCAGGTCTTC TTTGAAGGCC TATGGCAATG GCTACTCCAG CAACGGCAAC A 51

<210> 358
<211> 51
<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43040273

<400> 358

ATTGTAGTAC AAATGACTCA CTGCTATAAA GCAGTTTTTC TACTTTTAAA G

51

<210> 359

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43040273

<400> 359

ATAAACTTAG AATAAAATTG TAAAAATTGT ATAGAGATAT GCAGAAGGAA G

51

<210> 360

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43998970

<400> 360

AGGGGTGGAA CTGCTGATGG GATTTTCCTT CATTCCCTTC TGATAAAGGT A

51

<210> 361

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43998970

<400> 361

AGCCTCCCA GAGACAACAC CGGGACCCTC ATCTCTCTCC TCACCCTGCT G

51

<210> 362
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43947199

<400> 362
GTCTTCTCCG CGCCCACCCC GCTGGTAAGG GGAAGTGGGC GAAGCTGGAG C

51

<210> 363
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43917396

<400> 363
GGGGCCGGGC ACTGCCCAGG AAGGGGCTCC GGGAGAGGGA GCCGGCGGCT G

51

<210> 364
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40351913

<400> 364
AGACGAAGAC CCCAGGAAGT CATCCCGCAA TGGGAGAGAC ACGAACAAAC C

51

<210> 365
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)

<223> Accession number cg43921289

<400> 365

CCCACGCCCTG CCAGGAGCAA GCCGAAGAGC CAGCCGGCCG GCGCACTCCG A

51

<210> 366

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43928515

<400> 366

AAACAAATAA GCCCTTTTCTA CTGACGATGC ACCCAACCTT TTCAGCTGAA G

51

<210> 367

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43955093

<400> 367

AGAGTCAAAA ATCCAAGTTT GGATTGTAAG CAGCCTTGAC AGTAATCACT G

51

<210> 368

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43955093

<400> 368

AAGCAGCCTT GACAGTAATC ACTGAGTGGT AGGGAAAAAA AGACAGTTGG G

51

<210> 369

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43925474

<400> 369

AGGCAAAAGC TCACAGTAAA TGTATCCCAG AACAGGGGCC TAAGTGAAGG T

51

<210> 370

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg44014437

<400> 370

CTGCTCCCAN CTTGCCAGC CTCCAGTGTA CAACTTCCGC GTGTAGTGGG C

51

<210> 371

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg44014448

<400> 371

CTGCTCCCAA CTTGCCAGC CTCCAGTGTA CAACTTCCGC GTGTAGTGGG C

51

<210> 372

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43973129

<400> 372

CACTTCACTG AAAGACACCA TTTATATACC CAAGGGCAGA AAGTAGAACT T

51

<210> 373

<211> 51

<212> DNA

<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43950657

<400> 373
GATAGGACTC AAGCTTATTT GGGATTCTGA TCAATTCTTT CTGATGTTGT T 51

<210> 374
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956384

<400> 374
TACAGCCATC TGTACCTACT GGAGCTGCAG AAGGGAAGTC CACTCAGTCA C 51

<210> 375
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43992229

<400> 375
AGCAGTGCAG CCCC GGCGCG GAGCAAGGAG CCTCGGCCCG CGCCCGGCGC C 51

<210> 376
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44932392

<400> 376
GAGAAAAAGC ATGGTACCCA ACCGATTTTC CACTTTTCAG CAATACTTCA C 51

<210> 377

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg44932392

<400> 377

TAAAGTTTTA AGAAATGTCA TAATGTCATG AGCTTGAAAT ATCTCTAGGC A

51

<210> 378

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43981656

<400> 378

AGCAAAGAAA CACTGGCAGA ATTCCTGCAT TTGCAAAT CTAAGTTTTG G

51

<210> 379

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg44910613

<400> 379

AAATAAATGT TTCATAGTC ATTACACTTT ACAATGGGAG TGCTAAAATT C

51

<210> 380

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg44035104

<400> 380
AACTGGGTTG CTCTAAGAAC TGATGCCTAA ACCGTCTCAG CATGGCCTGT A 51

<210> 381
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43929959

<400> 381
CAATGCATGA ATCTGTACCC TTCGGAGGGC ACTCACATGC CGCCCCCAGC 50

<210> 382
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43950250

<400> 382
TTGTTTCATGA TTTCTTGATG TTCCTTAATG GAAAACTAAG AGATGGAATT 50

<210> 383
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43064090

<400> 383
GCCGAGTCCG CTGGTGGGCG GACCCTAGGG GAGCAGCCAG TAGGGAAGTT G 51

<210> 384

<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43064090

<400> 384
CCGAGTCCGC TGGTGGGCGG ACCCATGGGG AGCAGCCAGT AGGGAAGTTG G

51

<210> 385
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43064090

<400> 385
GGGAGCAGCC AGTAGGGAAG TTGGGGGAGT TCCAGAATCA GGGGGCGTGG C

51

<210> 386
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43064090

<400> 386
TAATCGGGAG GGCTGGAGCA GAGGGGGGCC CCGCCGAGGG GCGTGGTCAG T

51

<210> 387
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg30490224

<400> 387
GATGCCAAAA AAACAAAGGT GAGAACCCAC AACACAGGTC TAAACTCAGC A 51

<210> 388
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43924431

<400> 388
GTCTTTTACA GATGGTTTTT CAAAAAGAGT CCAGTAAAT ATTTCACATT 50

<210> 389
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43936047

<400> 389
CGTTGTTTCCT AATGTGGATC TACCATCCCT GTGTTTCATCG AGATTCCGGT C 51

<210> 390
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43272443

<400> 390
TGGGATTACA GGTGCGCACT ACCACGCCAA GCTAATTTTT GTATTTTTTA G 51

<210> 391
<211> 51
<212> DNA
<213> Homo sapiens

<220>

<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43966848

<400> 391
CCTTCAGCAC CCCTGCAGCG GAAAATAATG AGCCGCCGTA GCCGCCATCC G

51

<210> 392
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43964140

<400> 392
AAAAAGCTAC AGAAAAGAAA TCACTCTGAA AAACACAATG ACTCAGAGGC A

51

<210> 393
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43285114

<400> 393
CAGGGACATG CGGGCACCCC GTGGGTCTTT GCGGGCTCAC AGGACAATGG

50

<210> 394
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43948566

<400> 394

GCAGGCAGAG CACCCTGGGA CCCAGGCAG AAGGACCCCT GCCCTCCAGT

50

<210> 395

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg44003626

<400> 395

TAAACAGCTC AGTTCAGGA CTGGTGTACA AGCTGGCCAC CCATCTCAGC C

51

<210> 396

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43917206

<400> 396

TTACAGGACA TCACCTGCCA TCTTAAGGTT TAATATTTAC AAATGCCTAG T

51

<210> 397

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43289666

<400> 397

GGCCGATTTT TCCACAATTT AAATCTCAGT TCACCTGGTA TCCAGCTCCA G

51

<210> 398

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43986282

<400> 398
GTTTCCACCT CCCAGACAG GCATTTCGAG TGGGAGGCGG GAGCACGTAC C 51

<210> 399
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43986282

<400> 399
TTTCCACCTC CCCAGACAGG CATTCTGAGT GGGAGGCGGG AGCACGTACC G 51

<210> 400
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43297716

<400> 400
CTAAACCCAA ATGGGGGCTG CTGGCTGACC CCGAGGGTGC CTGGCCAGTC C 51

<210> 401
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43980312

<400> 401
TTTTATCATT AAAGTGCCAG AATGGTTCTT TAATGAAAAC AAAAAACAAA G 51

<210> 402
<211> 51
<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43939240

<400> 402

GGAGGGTTGG AGTCACTGAC GAATGTGAGC CGGGCCAGGC CCATGCAAAG G

51

<210> 403

<211> 50

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (25)...(26)

<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43941552

<400> 403

GCCACCTGCC CGGGCTGTGG AGGAGGCTCG CGCTGACCAG GCGCTGGGGC

50

<210> 404

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43941552

<400> 404

GCTTCTGCCC ACACCGCAGG GACAAGCCCT GGAGAAATGG GAGCNTGGGG A

51

<210> 405

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42917153

<400> 405

CATTTCCTTT TGTACATAAT ACATTACCT CCCTGCCTCC TCTCCTTTCT A

51

<210> 406

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43927693

<400> 406

CAGGGGTCAG CAGAGCTTCA GAGGTGCCC CACCTGAGCC CCCACCCGGG A

51

<210> 407

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43951338

<400> 407

CAGAAAGCAG CAAATTAGTG TTTTAAAGGA CCGAATTCGG CTCCCGCAGC T

51

<210> 408

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43951338

<400> 408

AAGCAGCAAA TTAGTGTTTT TCAGGCCCGA ATTCGGCTCC CGCAGCTCCT G

51

<210> 409

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43951338

<400> 409

CTCCCGCAGC TCCTGCATCT CCATTTGTCT AGATTTTATT TCTTCITTGC A

51

<210> 410

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg25236776

<400> 410

CCCGCCGAGC CCGACGCCTA CTGAGTCCCC GCGCTCGCCC CACCGGCGCG C

51

<210> 411

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg25236776

<400> 411

CCAGCCCGAC GCCTACTGAG CCCCCTGCTC GCCCCACCGG CGCGCTCTTC G

51

<210> 412

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg25236776

<400> 412

AGCCCGACGC CTACTGAGCC CCGCGTTCGC CCCACCGGCG CGCTCTTCGC G

51

<210> 413

<211> 51

<212> DNA

<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg25236776

<400> 413
CGACGCCTAC TGAGCCCCGC GCTCGTCCCA CCGGCGCGCT CTCGCGCCC G 51

<210> 414
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43968406

<400> 414
GCTACGTTTA CTCACAGCCA GCGAACTGA CATTAAATA ACTAACAAAC A 51

<210> 415
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42748886

<400> 415
CGCCTCTGAT CCAAGCCACC TCCCGTCAGA GAGGTGTCAT GGGCTTCCAA A 51

<210> 416
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43969533

<400> 416
CTCTGCACAA GGAAGCCTA TCCTATTTTT TTTTCCTTG CGAAAACAGA 50

<210> 417
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43976681

<400> 417
AATGCCTCAG ATCAGTGACC CAAGGACCTT CCAGAATGGA TGAAATAGAC 50

<210> 418
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43976681

<400> 418
ATGCCTCAGA TCAGTGACCC AAGGACCTTC CAGAATGGAT GAAATAGACC 50

<210> 419
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43984044

<400> 419
CCAAGCGGAA GGCCATTTTC CTGCTCTTC CTCAGTTGTC CGGGGCGGG G 51

<210> 420

<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43933283

<400> 420
CTAATTGTGT CGAATTTCCA GGATTAGAGG AAAAGTTGCT CCCTTTCAGC C 51

<210> 421
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42381630

<400> 421
AAAGCAATCA CAGTGTAAAA AGAAGACACG TTGAAATGAT GCAGGCTGCT C 51

<210> 422
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41664708

<400> 422
CCAGCCAGCT CATTTCACTT TACACCCTCA TGGACTGAGT TTATACTCAC C 51

<210> 423
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43277632

<400> 423
AGCCTTCCCG CAGAAAAAGA TGCAGTCCCC CAGACCTTCT CTGTGCTGAT T 51

<210> 424
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40310734

<400> 424
TACCAGAGGC TGCATCGGCT GCGCGGAGAG CAGATGGCGT CGTATTTTGG G 51

<210> 425
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40310734

<400> 425
TGGTGGGCGC TCCACTGTAT ATGGACAGCC GGGCAGACCG AAAAÇTGGCC G 51

<210> 426
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40310734

<400> 426
CTCTCAACAG GCAGGCACCA CCCTGGACCT GGATCTGGGC GGAAAGCACA G 51

<210> 427
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43982507

<400> 427
GGTTACAAGT GTGAATGTAG TCGTGCCTAT CAAATGGATC TTGCTACTGG C 51

<210> 428
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41554010

<400> 428
GCCGAGGACG TCGGTGGCAA CCTGAAGGGC AACACCGAGG GGCTGCAGAA G 51

<210> 429
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43299024

<400> 429
TACGAGGTGC CCTTGGAGAC CCCGCGTGTC CACAGCCGGG CACCGTCCCC A 51

<210> 430
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43299024

<400> 430
GAGGAGCCCT TCGGGGTGAT CGTGCCCGG CAGCTGGACG GCCGCGTGCT G 51

<210> 431
<211> 51
<212> DNA
<213> Homo sapiens

<220>

<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43285373

<400> 431
GACTGACTGG GGAAAGGAAC CTAAACTCGA GTCTGCCTGG ATGAATGGAG A

51

<210> 432
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg36834323

<400> 432
AGTTATCTTA GAGGATACAG AAATCGTCGA AGTTCCCGAG AAAGTAGGGA G

51

<210> 433
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41568631

<400> 433
GGACCAGGGG GCCATGCTGC TCAATATCTC AGGCCACGTC AAGGAGAGCG G

51

<210> 434
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42359655

<400> 434
TGCTTTTCAG GCGGAAAC TCTCTGTTGT CCTGCGAGCT GAAGATATCC C

51

<210> 435
<211> 51

<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43998672

<400> 435
GTGTGGGCCT TGGTGAAGCTC TAGCAAGCGG CTAATGTCTC CTGGTTTGGT C 51

<210> 436
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43969028

<400> 436
GGAGATGTGG TCATTCCTAG TGATTTTTTT CAGATAGTGG GAGGAAGCAA C 51

<210> 437
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43933479

<400> 437
AAGGAGAAGA GAAAGCTGTT TATCCGTTCC ATGGGTGAAG GTACAATAAA T 51

<210> 438
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43942537

<400> 438

GCCACTGTCT CTTCCAAACC CTTCAAGCCT TGTCTTGCTT GTTCTCGTCT A

51

<210> 439

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg38337333

<400> 439

TTCCAAATGC TGAGCCCAGA GCGTTATCTC CTGCCCATGA GCACCACAGT C

51

<210> 440

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg38337333

<400> 440

CTGGAACAGT TTCCTCATTG GCCCTCTGAC CCCAGCACAC GCAGGGACCT A

51

<210> 441

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg38337333

<400> 441

TCATCGCTGG TGCTCCAAAA AAAAAGATGC TGCTGTAATG AACCAAGAGC C

51

<210> 442

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg30421838

<400> 442

GGATGCTGTT GCTCTCCAC AGCCATTGGG CGTTCCAAAT GAAAGCCAAG C

51

<210> 443

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43064060

<400> 443

GCCAATGGCA TCCAGAACAA GGAGGTGGAG GTCCGCATCT TTCCTGCTG C

51

<210> 444

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43991813

<400> 444

TCTCGACTAA CAGCATTTC AAAGACGGAG CGAATATTGT CCACGGTTGA G

51

<210> 445

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42904626

<400> 445

GAGTGCCTTG ACGATACAGC TAATTGAGAA TCATTTTGTG GACGAATATG A

51

<210> 446

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42904626

<400> 446
AAGAAGTTAT GGAATTCCTT TTATTCAAAC ATCAGCAAAG ACAAGACAGG G 51

<210> 447
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42460457

<400> 447
GCCGCCTCAG CCAGCAAGCA GCGGTTAGG CCAGTCCTAG CCACCACAGA G 51

<210> 448
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43272594

<400> 448
GGGATGTACT GCATGGTGT CTTGGCGCTG TATGTGCAGG CACGACTCTG T 51

<210> 449
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43958858

<400> 449
TCAGGTGGTG GGAACCTACC GTTGCGTTCC TGGAAAGAAG GGAGGCTACA C 51

<210> 450
<211> 51
<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43916732

<400> 450

GTACAGCGGG CGGGCCACCT CGGGCTCTGA GCACCAATTT TGCGGGGGGC G

51

<210> 451

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42894809

<400> 451

GGATGCTGGA GAGTGGATCA CTGTCGATCA GACGACAACA GCCAACCGTT A

51

<210> 452

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg40388639

<400> 452

GATTCTCCA GAGCTGGTGT TGGAACCTCC CATCAGGCAC CCCAAGTTTG A

51

<210> 453

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg40388639

<400> 453

AAGTTTGAGT GGTTCAGGA CCTGGCGCTG AAGTGGTACG GCCTCCCCGC C

51

<210> 454
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43124627

<400> 454
AATTTCTATA TCACTGGGGA CAGAGGATAT ATGGATAAAG ATGGGTATTT C

51

<210> 455
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43124627

<400> 455
TGGAACAAGT GGATATCCGA AAATGTCTGC ACACACCCAC AGCAGTTTTG G

51

<210> 456
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43064068

<400> 456
AGGAGAGGTG GTGAAGGCAT TTGTGATCCT GGCCTCGCAG TTCCTGTCCC A

51

<210> 457
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)

<223> Accession number cg2514276

<400> 457

GTGATGGACC CTCTCATATA TGCCTTCCGC AGCCAAGAGA TCGGAAGAC C

51

<210> 458

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg32423505

<400> 458

TTCCATCTGA GGTTTATAAA CCACGTATTC AGGCAAAGTG GCCAGAATGG C

51

<210> 459

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43335558

<400> 459

CAAGACCTAG CTCCCCAGCA GAGAGTGGCC CCACAACAAA AGAGGTCCAG C

51

<210> 460

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43998970

<400> 460

GACAGAGCTG TACCGTGACA TTTTCGAGCA CCTTCGGGAT GAATCAGGCA A

51

<210> 461

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg2537639

<400> 461

GAGGGCGATT TCTACTACCT GGGGGCGTTC TTCGGGGGGT CGGTGCAAGA G

51

<210> 462

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43935995

<400> 462

AGTGTCCTC ACCATGGTCA CCCTGGTCAC CCTGCCTCTG CTTTTCCTTC T

51

<210> 463

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43935986

<400> 463

CCTGGAACGC GCCTTGTAAC TGCTCATAAG GAGGGTGCTG CACTTGGGGG T

51

<210> 464

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43968274

<400> 464

GAGCAGAGG AAGCCATGAA TGCGGTCTAC TCAGGCTACG TCTACACGCA C

51

<210> 465

<211> 51

<212> DNA

<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44018598

<400> 465
AGATACTTTC TATAAGCAGT TTTTACATTG TAGGAAGCAG CTGAATTCAA A 51

<210> 466
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44926796

<400> 466
ACACTGGAAA GCACAACAGT TGGCAGTTCT GTCTAGAAAA TAATAATTGC A 51

<210> 467
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43055918

<400> 467
AACGCTGCCC TGACTGAGAA AGGCATGATG CTCGCTCCAC TGCTGGAACC G 51

<210> 468
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43966985

<400> 468
CATCCAGGAC AACTTCTCGG TGACTGAAGT GCCCTTCACT GAGAGCGCCT G 51

<210> 469
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43918854

<400> 469
CTTCAACCCCT GGTCGGAGAC AACGGCTCAC CATGGCCATC AGAACAGTGC G

51

<210> 470
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43918484

<400> 470
CTGATTCTTC CGTTCTTCTT GACTTGTGCC ACCTTGCCAG CCAGCTGCTC G

51

<210> 471
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43942977

<400> 471
ACGGCCCTGG AGAACCAGAA GAAGGTGAGG AAGAAGAAAG TCTTGATTGC C

51

<210> 472
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43942977

<400> 472
GGATGGTGTC TGATGAGGAG TTGGATCAGA TGCTGGACAG TGGGCAAAGC G 51

<210> 473
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43943361

<400> 473
TTGGGGTTGG CTTGGTTTCA ATAAGCAACG GGGACACTTA CAAATTGCTG C 51

<210> 474
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg25236776

<400> 474
GTGACCGAGC CCGAGTGCCG CGAGGTCTTT CACCGCCGCG CCCGCGCCAG C 51

<210> 475
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg25236776

<400> 475
CAGTGCCTCC CCTGCGGCCC CGGGGTCAAA GGCCGCTGCT TCGGGCCCAG C 51

<210> 476
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg38899722

<400> 476
GGCCAACTCT GCTATGGACA CCAGACTACT CTGCTGTGCG GTCATCTGTC T 51

<210> 477
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg11753818

<400> 477
GCCTGGAACA CCAGGCTCCT CTGCCATGTC ATGCTTTGTC TCCTGGGAGC A 51

<210> 478
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2526759

<400> 478
AGCCACCCAG ACCGGAGACT CGGCCATCTA CCTCTGTGCT GTGGAGGCCT A 51

<210> 479
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2526759

<400> 479
CGGCCGTCTA CCTCTGTGCT GTGGACGCCT ATTCTAACGA CTACAAGCTC A 51

<210> 480
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg1902363

<400> 480
CAGAACAAAA GCAAATGGAA TTGGATAGCA TCCTGGTGGC CCTGCTGCAG A 51

<210> 481
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43277632

<400> 481
GAAACCCGGA AGCACTGTAA TTGCGGGGTC TATAAATGCA CATGGCTCTG T 51

<210> 482
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43252813

<400> 482
TGTATTCCTG TAATGGGGCT GATGATATAT ATGATGGTTA TGGACCACCA C 51

<210> 483
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43920913

<400> 483
GCCCCTGAGC AGTCAGGACC CGGCTTCCGT CCGTGAGTGC CACGATCCCA G 51

<210> 484

<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40310734

<400> 484
GGAGTGGGTG CTGCTGCTCT TGGGAGCTTG TGCTGCCCCCT CCAGCCTGGG C

51

<210> 485
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40310734

<400> 485
CGTGTCTTCC CTCCCCATG CGGTGGCCCC GCTCAGCCTG CCCCAGGGG A

51

<210> 486
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956560

<400> 486
GGGTACTAT GGGCCCCAGT GTCAGCTTGT GATTCAGTGT GAGCCTTTGG A

51

<210> 487
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956560

<400> 487
GCTGGGTACC ATGGACTGTA CTCACTCTTT GGGAACTTC AGCTTCAGCT C 51

<210> 488
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42388009

<400> 488
TGCAGAAGGC ACCACAGAGA CCGGAGGGCA GGGCAAGGC ACCTCGAAGA C 51

<210> 489
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43977436

<400> 489
GTGTGTGTGT AATGGTGTGG CTGTATGCTC CAACCAAGAT CTTATTACTG A 51

<210> 490
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43280376

<400> 490
CGGAAGCTGG TGTCCTACTG CCCCCGAAGG TTGCAACAAC TGTTGCCCCCT C 51

<210> 491
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42201364

<400> 491
CCAGGGCCTC CAGGTCCAAG AGGCCCCTCT GGAGAGCCTG GTCTTCCAGG G

51

<210> 492
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42201364

<400> 492
GTGTTTACG CTGAACGATA CCAAATGCCC ACAGGCATAA AAGGCCCACT A

51

<210> 493
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40339378

<400> 493
TCCAGGAATA CCAGGTCTGC CTGGTGTTC TGAACAAGA GGATTAAAAG G

51

<210> 494
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43063256

<400> 494
AGACTGTGTT ACCAACAGAC CATGCGGAAG TCAAGTGCGA TGTGAAGGCT T

51

<210> 495
<211> 51
<212> DNA
<213> Homo sapiens

<220>

<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44032748

<400> 495
CTCCAGTTCT ACAACTTGTG TAAGGCAAGC ACAGTGTGGA CAGGATTTCC A

51

<210> 496
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43049885

<400> 496
CAGTTTGGGG GACAGCCATG CACTGCGCCT CTGGTAGCCT TTCAACCATG C

51

<210> 497
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg21644442

<400> 497
CCAGGCTCTC CCAGGATCTC ATCACC GCGC CCCAGGGCC TCAGCAACCC C

51

<210> 498
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2753430

<400> 498
CCAAGCTCCC ATGACCCAGA CAACGTCCTT GAAGACAAGC TGGGTAACT G

51

<210> 499
<211> 51

<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43923204

<400> 499
TCCACGTAGA AGCGGAAGCC GAGGTGGGAG ATGTACGCAT TGATGGGAAG G

51

<210> 500
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44017721

<400> 500
TTGGTAGGGA CGGAACCTCGG GCGCTGGCG GTGGCCCCGAG TGGAGATAGG A

51

<210> 501
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41626024

<400> 501
GCTGGTTTGC TCCCAGGAGG CCAAGCAGTC AGCCTACTGC CCCTACAGTC A

51

<210> 502
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43057018

<400> 502

AAGCATCCGA ACAATCCTCA TCTTTGGAAG ATGCCAGGAG CAATTCGGAA T

51

<210> 503
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42837709

<400> 503
CCGCACCAAC GCCGACATCA TCGAGGCCCT GAGGAAGAAG GGCTTCAAGG G

51

<210> 504
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43327954

<400> 504
TCCACGACCG GGTAGAGAAC TACAAACCGC GGCAGCGCAA GCTCCGCAAC C

51

<210> 505
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43971258

<400> 505
TCGTTGGAGA TGACAAGTTC CGGAGTGAGC TCGGCTGTCT GGATGGGAAG G

51

<210> 506
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg41554010

<400> 506

AGCTGGAGCA ACAGCAGGAA CAGCATCAGG AGCAGCAGCA GGAGCAGGTG C

51

<210> 507

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43957743

<400> 507

GTTTGGCATA CCTGGATATT TTAATTCAGT GGAGATAAAA GACAGCCCAC T

51

<210> 508

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43957743

<400> 508

TATTTTAATC CAGTGGAGAT AAAAGCCAGC CCACTAGGAA GTATATCAAT A

51

<210> 509

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43248101

<400> 509

AAGTGAATTC TATCTTGCAA TGAACGAGGA AGGAAACTC TATGCAAAGA A

51

<210> 510

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43969014

<400> 510
GATGAGCTCT CCAACCACGT ATTTTATGCG TTTTGTATCC AGACCCAGAT G 51

<210> 511
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43286488

<400> 511
CACCAGCAAG ATGCCACGA TCAGCCGAAC CTGCCCAAGG CCTGCTTCTT G 51

<210> 512
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44004239

<400> 512
TTTTCCCAG GGGTCACAGA CTGATGACCC ACAGAGGTCA GGGTCTTCTG T 51

<210> 513
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44004239

<400> 513
GGGGTCACAG ACTGATAACC CACAGGGGTC AGGGTCTTCT GTCCAGTGGT C 51

<210> 514
<211> 51
<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg44004239

<400> 514

CTGATGACCC ACAGAAGTCA TGGTCGTTGC CCCAGTGATC TCAGTCTTCT C

51

<210> 515

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43932434

<400> 515

ATATGTGTCA TACTGGGAGG TGTTGTATGT GAGGATGTAC ACCCCTGTGT T

51

<210> 516

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg40915005

<400> 516

AAGGAGCCTC TCTCCTTCCA TGTCATCTGG ATCGCATCCT TTTACAACCA T

51

<210> 517

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg40915005

<400> 517

ATTCCAGCAC CATCGTTTTC CTGTGCCCCT GGTCCAGGGG AAACCTCAGC A

51

<210> 518
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg36834323

<400> 518
GTGCTCCCTG ATCCTCGTGA AGCATGTGGT AGCTCAAGTT ATGTGGCATC T

51

<210> 519
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg36834323

<400> 519
AATGCTGCGA AAGATATGAA TGGAACGTCT TTGCATGGAA AAGCAATAAA A

51

<210> 520
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg36834323

<400> 520
AAGTCTGAGA TCTGCAAGAG GAAGCCGTGG AGGAACAAGA GGGTGGCTTC C

51

<210> 521
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)

<223> Accession number cg44019290

<400> 521

GCGGATAAGT AGAGGACCTT CATGTGGTAT TTGCTGGTGA AGTTGGTTCG G

51

<210> 522

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42336656

<400> 522

CTTAGACATA CAATATACTT ACCTTGGAGG TCACGTATGT TTGTCCGCAC A

51

<210> 523

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42730678

<400> 523

GGAGCGAGCG TGGATCCAGT TCGCGTCGGG GTTGTTTGGG TCAAGTTGCT G

51

<210> 524

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42714160

<400> 524

GGCCCTGTGCC TGACGGAGAG GCAGAGCAAG ATATGGTTCC AGAACCGACG C

51

<210> 525

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42359655

<400> 525

CTGGGCACCA TATAGGATAG CCCACGCCGT CATCAAAGCC CATGCCAGAG T

51

<210> 526

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43925670

<400> 526

GTGGAGGGTG CAGGTGAAGT AGCATGCACT TCCTTCTTCC TCTTTCTTGA T

51

<210> 527

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43090990

<400> 527

TGCTCCATCA AAAATGAAGC AAGGCTGCCA TGTTTACCGA CACCGGAAA A

51

<210> 528

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43969763

<400> 528

CAAAAGCAAG AAAGTTCTTT GAGAATTTC CAGATGGCAC TTGGAACCTA A

51

<210> 529

<211> 51

<212> DNA

<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43932396

<400> 529
AGTCCACCGC CGCCTCAGGC CGTGCTGCTG GCCGAGTAGG AGAACTGGGG G 51

<210> 530
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43917871

<400> 530
GGCACTGAAG AAATCCCTGA CATCACATTG GCGCTGCTGA CGGGCGTACT G 51

<210> 531
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43917871

<400> 531
GGGCTGACAA GGTGCTGATT TTCACGGTGG ACAAAGCGTT CCCATCGCTT T 51

<210> 532
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43917871

<400> 532
TTCAATGTTG TATTTGTCAA TATAGCCATA TAAATCTTCT GTCCCCAGAA C 51

<210> 533
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43917871

<400> 533
TGTAATCG AATATCATAG TCTGTGAACG TCTGGTACAA TTGCTTGAAG T

51

<210> 534
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43322545

<400> 534
TCGGCTAGGC AGCCTCCATC CTCACCCCCC TTATCACATC CGCGTGGCAT G

51

<210> 535
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43322545

<400> 535
TTCCTCCTCT ATTCCCGGCT CGGGGGCCAG CCAGTGATCC TGCCCACTCA G

51

<210> 536
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43958558

<400> 536
TCCGGGATAA GCTCCAGGTG CTCCATGGTA GGCGCCTGGA GGTGCCTGTC C 51

<210> 537
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43966144

<400> 537
AAGCTTGTC TGCCTCACAG CAGTGAGCAC AAGACTGCCC AGCCCAATGG A 51

<210> 538
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43966144

<400> 538
ACTTACACCT GTGTGGTAGA GCACACTGGG GCTCCTGAGC CCATCCTTCG G 51

<210> 539
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42686658

<400> 539
GGCTGGTGG GCTTCCTCGT GGGCATCGTC CTCATCATCA TGGGCACATA T 51

<210> 540
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg38337333

<400> 540
CTGAGCCCAG AGCGTTGTCT CCTGCGCATG AGCACCACAG TCAGGCCTTG A 51

<210> 541
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg38337333

<400> 541
AGCCCCGCCG GGCCCCACGG TTCGCGCAGG AGAGAACGTG ACCTTGTCCT G 51

<210> 542
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42481172

<400> 542
CCGGCTGTGC TCAGGGGTGT GGGGTGCGGA TACAGAGGAG CGGCTGGTGG A 51

<210> 543
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg3000465

<400> 543
GAAGATGCCC TCCTCAGACA TGAGTTGAAA GGTTATCAGA AATGGGTCCG C 51

<210> 544
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg3000465

<400> 544
AGATGCCCTC CTCAGACATG AGTGGCAAGG TTATCAGAAA TGGGTCCGCC C 51

<210> 545
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43249083

<400> 545
GCCTCGGGCT TCCACTACGG TGTGCTCGCC TGCAGGGCT GCAAGGGCTT T 51

<210> 546
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44928796

<400> 546
AGCGGACGG TCCGGAGCAA GCCCACAGGC AGAGGAGGCG ACAGAGGGAA A 51

<210> 547
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43323772

<400> 547
GTGCCGGGAG TGAGCGATGA GCTGGTTTCT GTTCCTGGCC CACAGAGTCG C 51

<210> 548

<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42732993

<400> 548
GGCTATAATC ACAATGGGGA ATGGTTTGAA GCCCAAACCA AAAATGGCCA A

51

<210> 549
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42904626

<400> 549
ATATAAACTT GTGGTAGTTG GAGCTTGTGG CGTAGGCAAG AGTGCCTTGA C

51

<210> 550
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42904626

<400> 550
TGGATATTCT CGACACAGCA GGTCACGAGG AGTACAGTGC AATGAGGGAC C

51

<210> 551
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42691989

<400> 551
CTGTTTCAGGA TCTCCTCATT CTGACTGTTC TCCTGATGTC CAAATTGGTT G 51

<210> 552
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43917453

<400> 552
AGGTCCTCGC GGAGCTGGGT CCGGGGCCCCG GGAGGGTAGG TCAGCGCAGA C 51

<210> 553
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43947363

<400> 553
CTGGCGCACT ACTCGGACCT GCTCCTCCTG GCGGGCCTGG GGCTGATTGA G 51

<210> 554
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43928335

<400> 554
GCACAAGGAA CGGAATTGCT GTCTGGTTTC TGCTTTAACA GCATTTGATG C 51

<210> 555
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43996195

<400> 555
CTTCGGGGAA AGTTGGGGAT TTCACTGTAG TCAAAGATCT GGGCCTGAGT T 51

<210> 556
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44022214

<400> 556
AGGTCTCTCT CGAATTGGGA TGGCCGAGGT GCATCATCAT CATCCCAGAG G 51

<210> 557
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43958858

<400> 557
CTCAGACCAT GTCC^{TT}CGGA TGCACGGTTA CAGAGCACCT GGGGAGCAGG A 51

<210> 558
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42534568

<400> 558
CGCTTTGAGA CGCAGCTGGG CACCC^TTGGCG CAGTTCCCCA ACACACTCCT G 51

<210> 559
<211> 51
<212> DNA
<213> Homo sapiens

<220>

<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42534568

<400> 559
GGGGGACGAG GCCATGGAGC GCTTCGGCGA GGATGAGGGC TTCATTAAAG A 51

<210> 560
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42534568

<400> 560
CATTAAAGAA GAGGAGAAGC CCCTGGCCCG CAACGAGTTC CAGCGCCAGG T 51

<210> 561
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43154190

<400> 561
TGGAGGGGAT GCTCATTITG ATGAACATGA AAGGTGGACC AACAAITTTCA G 51

<210> 562
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43154190

<400> 562
GATGAAAGGT GGACCAACAA TTTCACAGAG TACAACITAC ATCGTGTTC G 51

<210> 563
<211> 51

<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43927549

<400> 563
ATTCTACGAT TCCGGTTTGC TCCAGTGTA ACTAGCGCTC CTTCCGTAA C 51

<210> 564
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43325541

<400> 564
CGCTGCCTTC TCCCGAAAGG TCTGCTCCTT CACGCGTTTC GCTTCCCGCA G 51

<210> 565
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43064068

<400> 565
GTGAAGGCAT TTGTGGTCCT GGCCTTGCAG TTCCTGTCCC ATGACCCAGA A 51

<210> 566
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43064068

<400> 566

GACTGTCACA GGGAAAATTC AACGAACCAA GCTTCGAGAC AAGGAGTGA A

51

<210> 567

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg36988276

<400> 567

GCAAGAAGTT GATTATATGA CTCAGGCTAG GGGTCAGAGA TCCTCTCTGG C

51

<210> 568

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg36988276

<400> 568

AAGGCCAACA ACCTGCTCTA CATCACCCCT GAGGCCTTCC AGAACCTTCC C

51

<210> 569

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg32296848

<400> 569

GAATGTCTTG AGAATCCAGT GTCTCTGCAG AAAGCAGTCT TCCAAACATG C

51

<210> 570

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)
<223> Accession number cg2524739

<400> 570
TCTCTCTGGA GAAGATCCAA CCCATGACAC AAAACGGTCA GCACCCAACC T 51

<210> 571
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2320320

<400> 571
AGTGTCTGGA TGATCTTTGT GGTCAATGCA TCCGTTTTCA CAAATGGGCT T 51

<210> 572
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43264978

<400> 572
CATCTTCTCC ATCAACCTCT TCAGCGGCAT TTTCTTCCTC ACGTGCATGA G 51

<210> 573
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg3003708

<400> 573
TCCTTTGTGG ACATCTGCTT CTCCTCCACC ACCGTCCCCA AGATGCTGGC C 51

<210> 574
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele

<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg38841806

<400> 574
GGCCCTGAGA GCAACACCAC GGGCACCACA GCCTTCTCCA TGCCCAGCTG G 51

<210> 575
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43336100

<400> 575
TGGAAGCGTG CATCCAGTGA GACCATTGAG GCTTGAGTCT TTTAGTGCCT G 51

<210> 576
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43335562

<400> 576
GAGGCGCGGG GAGCCAGGCC TGGGCCCCGG GTCCCCAAGA CCCTTGCTGCT C 51

<210> 577
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43140548

<400> 577
ACTCGCACGT GGATCCTGAG GCTGTGAGAG GTAAGGAAGG CTTTGCCACA G 51

<210> 578
<211> 51
<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43011561

<400> 578

CATTGACAGC GAGGCCTCCT CAGCCTTCTT CATGGCGAAG AAGAAGACGC C

51

<210> 579

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43998970

<400> 579

TGACAGAGCT GTACCGTGAC ATTTTGCAGC ACCTTCGGGA TGAATCAGGC A

51

<210> 580

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg2537639

<400> 580

CACTACTATG TCCTCACCGA CCAGCTGGCC GCGGTGCCCC GCGTGACGCT G

51

<210> 581

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg2537639

<400> 581

TCGGCAGCTG TCAGTGCTGG AGGTGGGCGC CTACAAGCGC TGGCAGGACG T

51

<210> 582
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2537639

<400> 582
GGTGTGCGTG GACGTGGACA TGGAGATCCG CGACCACGTG GCGGTGGAGA T

51

<210> 583
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2537639

<400> 583
TCCGCTGTTC GGCACCTGTC ACCCCAGCTT CTACGGAAGC AGCCGGGAGG C

51

<210> 584
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2537639

<400> 584
CAAGGACGAG GGCGATTCTT ACTACATGGG GGGGTTCTTC GGGGGGTCGG T

51

<210> 585
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)

<223> Accession number cg2537639

<400> 585

GTTCTTCGGG GGGTCGGTGC AAGAGATGCA GCGGCTCACC AGGGCCTGCC A

51

<210> 586

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg42742340

<400> 586

CACATCGTGG TGGAGCTGAC CCAGGATGAC GCTTTGGGCT CCAGGTGGCG G

51

<210> 587

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg41653463

<400> 587

GTCTGAAAGA TTTCCACAAG GACATGCTGA AGCCCTCACC AGGGAAGAGC C

51

<210> 588

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg40351913

<400> 588

CAAGTTCACC AACAACTGCT ACAGGCACGC GATTGTCACC ACCTCCATCA A

51

<210> 589

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg40351913

<400> 589

TCCTCCGGCT TCGTCGTCTT CTCCTCCCTG GGGTACATGG CACAGAAGCA C

51

<210> 590

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43955093

<400> 590

CTGCGGTAGC TGTCCCAGGC CTCGGGCCGC GCCGCCTCGT CCATGTTGAG G

51

<210> 591

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43055918

<400> 591

GCATAGGACA TGGCGGGCTT GCGCCGCGCA GAGCTCTGGG GGCTACTGCT A

51

<210> 592

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43968854

<400> 592

CAACCCCTAG AAGACCTGGC TGGCTGGAAA GAGCTCTTCC AGACACCACT A

51

<210> 593

<211> 51

<212> DNA

<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43070241

<400> 593
CCATTGTTCA AGACATCCTA CGTTTGGAAA TGCCTGCAAG CAAAATTGTC C 51

<210> 594
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43262121

<400> 594
ACAATTCAGA GAGGGAGACT GAGCATAAC CAGCATTGAT CATGGTGCCA A 51

<210> 595
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43262121

<400> 595
ATCAGGAAAG GTGTTGGATC ACTGGTGCAT CATGACCACT GAGGAAGAAG T 51

<210> 596
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43262121

<400> 596
CCCCAAACAG GAAGTCCATG GGCCCTACCC TGACAGCAGC TTCTTAACTT C 51

<210> 597
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43262121

<400> 597
CCCAAACAGG AAGTCCATGG GCCCATCCCT GACAGCAGCT TCTTAAC TTC C

51

<210> 598
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44024279

<400> 598
CTGGAAGAAC TTTGCCATGA GAAAGGAATT TTGGAGAAGT ACGGACATTC A

51

<210> 599
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44928804

<400> 599
AATGATTAAAC AACAACTGA GACACACGGA TGAAATGTTC TGAACCACG T

51

<210> 600
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43317253

<400> 600
GCCCCCAGGC ATGGCTAGCT CGTGTTCCTG GCAGGTGAAG CTGGAGCTGG G 51

<210> 601
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41637661

<400> 601
CAGCTTTCCA TCCATTTTAA TTTATAGACA TACTGCTAGT GGAAAGACCT A 51

<210> 602
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42913861

<400> 602
CAGGTGTCCT GCGAGCCACC CGGGGCTCCG GGTGGCGGGG GTGGCGGCCG C 51

<210> 603
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43249389

<400> 603
AGAGGAGAGA GCCGCCCTCG AGCGGGGCAA GGCGATTGAG AAAAACCTCA A 51

<210> 604
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43919239

<400> 604
GCCAGAGTTG CAGCATCAGG GCCAGCCTGA GCAGGAGACC CCCAGTCCCA T 51

<210> 605
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41642952

<400> 605
TAGGAATGAC AGCAGTAGCA GTAATGGGAA GGCCAAAAT CCCCTGGAG A 51

<210> 606
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43945147

<400> 606
TGTTCTGGA GCCTCAATGG TACAGCGTGC TCGAGAAGGA CAGTGTGACT C 51

<210> 607
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43926002

<400> 607
GGGCACAGAA ACACAGCAGC GGGAGSAGCA ACACCAGCAC TGCCAACAGA T 51

<210> 608
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43972311

<400> 608
ATTGCCATTG TGGTAACTCT GGGTCGCATC ATCTTCAGTG CCCCAATTGT G 51

<210> 609
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42556108

<400> 609
GTGAAGCGGT GTATGGGGAC AGTGAACCTC AACCAGGCCA GGGGCTCCTT T 51

<210> 610
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg36842490

<400> 610
AGTGACTTCA GTAAACTCTT GGGTCCACTT TCTGCCAAAA AGTACCTTGA G 51

<210> 611
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43942549

<400> 611
CGGTATAACG TCAAAAATCC TGTTTTTCAG CCAAGGTCA GAAATTGCCT C 51

<210> 612

<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42381630

<400> 612
AAGGCGCTAT GTACAGCCTC CTGAAGTGAT TGGGCCTATG CGGCCCGAGC A

51

<210> 613
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42381630

<400> 613
TGAAGATGGT CCTGATGGGC AGGAGGTGGA CCCGCCAAAT CCAGAGGAGG T

51

<210> 614
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg3004395

<400> 614
ATTACTGAAG GGTGGAGAAC AGAAGCGTCA TGAAAAAATA TCTGCTTCAT T

51

<210> 615
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43960645

<400> 615
CACTTCCTCT TTCTCTTTGG ATGCCTCACC CTCCTGTTGG GGGGCAGATG G 51

<210> 616
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2526759

<400> 616
GAAGACAAGG TGGTACAAAG CCCTCAATCT CTGTTGTCC ACGAGGGAGA C 51

<210> 617
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2526759

<400> 617
TGTAACCTCTC AATTGCAGTT ATGAAATGAC TAACTTTCGA AGCCTACTAT G 51

<210> 618
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2526759

<400> 618
GAAGTGACTA ACTTTCGAAG CCTACAATGG TACAAGCAGG AAAAGAAAGC T 51

<210> 619
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg2526759

<400> 619
AGCATATTAG ATAAGAAAGA ACTTTCCAGC ATCCTGAACA TCACAGCCAC C 51

<210> 620
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40310734

<400> 620
TACAGAATAT GTCGTCGGTG CCCCCCACTT GGAGCTGGAC CCTGGGAGCG G 51

<210> 621
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg40310734

<400> 621
GTCGGCTTCT TCAAGCGGAA CCGGCACACC CCTGGAAGAA GATGATGAAG A 51

<210> 622
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43956660

<400> 622
GTCCATCACT TCACTTCAGT TATTCCTAG GAGGTTGTAT AGTCTTCTGA 50

<210> 623
<211> 51

<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43970982

<400> 623
CTCCAGGGAT AGTTGGACAG AAGGGGAGAC CCTGGCTACC CAGGACCAGC T 51

<210> 624
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42175288

<400> 624
GCTATGGAGG CAAAATGGGA GGAAGGAAAC GACTACAGAA ATGATCAGCG C 51

<210> 625
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg42175288

<400> 625
CGGTACTCC AGTTATGGAC AAAGTCTATT CACAGTCCTA TGGTGGTTAT G 51

<210> 626
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg41554010

<400> 626

GGCCGAGCAG CTGCGGCGCC AGCTGGACCC CCTACGCACA GCGCATGGAG A

51

<210> 627

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43065549

<400> 627

CAGACTTCCA CAGAGTGCTG GATGAACGCG GCCTGCCTTG CCCAGGGTT A

51

<210> 628

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg41568631

<400> 628

TGACCACGGG GTGCTGGATG CCTGCCTTAT ACATCCTGGA CCGCGGGGG A

51

<210> 629

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg41637704

<400> 629

GCGCCGCGAG ACAAGGGCAG CGGACGCGCC TCGGGACTTG AGGGACAGTG A

51

<210> 630

<211> 50

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43933380

<400> 630
ATAAGTTACA ATGCTTTTTT TGTTTAAAAA AAAAAAAGT CTGTACTTTA 50

<210> 631
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43072541

<400> 631
CTGTGGGGCT GGTTCGTAT CTGATCATCA TTCGATTACG AAATAAACG T 51

<210> 632
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44032168

<400> 632
GTCAGCCGCT ACCTCGACTG GATCCTATGG GCACATCAGA GACAAGGAAG C 51

<210> 633
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43931248

<400> 633
CCGGGCAGAG CTGCGTCTGC TGAGGGCTCA AGTTAAAAGT GGAGCAGCAC G 51

<210> 634
<211> 51
<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43931248

<400> 634

CCGGGCAGAG CTGCGTCTGC TGAGGGCTCA AGTTAAAAGT GGAGCAGCAC G

51

<210> 635

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43272560

<400> 635

AATCTCCGCA CTGCAGGCCA GGGGCCTGGC CAGCTACAGA GAGAGGTCAC A

51

<210> 636

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43266471

<400> 636

CCAGGATCCA TTTTGAGGAT TATGGTGTGC TGGGACACCA TCAACTCCTC A

51

<210> 637

<211> 51

<212> DNA

<213> Homo sapiens

<220>

<221> allele

<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature

<222> (0)...(0)

<223> Accession number cg43995237

<400> 637

CAAAATCCCC CGTTTCTTCA TCTTGGACAT GCTAAATGA AATTACGCAG T

51

<210> 638
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43995237

<400> 638
TTCTTCATCT TGACATGCTA AAATGGAAAT TACGCAGTTT CTCTCTATCA A

51

<210> 639
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43254094

<400> 639
CCGCCTCTGC TGCTGCTGCT GCTGCGGCGT CCCGCCAGC CGCAGCTTCC C

51

<210> 640
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44034555

<400> 640
ATCCAGGCTG AGCTGGATCA TCTGAGGCCT CCAGCCACCC GTTTTCCCTT

50

<210> 641
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)

<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44034555

<400> 641
CCAGGCTGAG CTGGATCATC TGAGGCCTCC AGCCACCCGT TTTCCCTTGA 50

<210> 642
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg39711096

<400> 642
AGCGAGTCCT CCGGGAGGCC CACAGGTTAC TGCCTCCAGC TGCAGCAGTG A 51

<210> 643
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44128902

<400> 643
CGTTCCAGAG GAGCATATCT GCTGACTGAT GACCTGCAAG AGTCATCCAG A 51

<210> 644
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43946951

<400> 644
GGAAC TCGAG CACGTCGTCG GGGGACCCAA GATCACCGGC GCCCTCTGGT 50

<210> 645
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43948890

<400> 645
ATCCCCGGGG GAGGGGGCCC TGTAAGGAAA CCAGACAATC CCATGAGACT 50

<210> 646
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43948890

<400> 646
TCCCCGGGGA GGGGGCCCTG TAAGGAAACC AGACAATCCC ATGAGACTCC 50

<210> 647
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43917524

<400> 647
GGGCCTGTCT GCCCAGTGGG GGAGGTTCCG CTGGTGTCTT AGGGGGCATC 50

<210> 648
<211> 51
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43942004

<400> 648
CTCTCGGCAC TGGTGACTGG CGAGAGCCTG GAGCGGCTTC GGAGAGGGCT A 51

<210> 649
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg43932428

<400> 649
TCGTGGCCAG GTCCTTCTGC GTAAGCCCTT GCTCTGCCGA CCTTGCTGGA 50

<210> 650
<211> 50
<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44010855

<400> 650
GGTCCAAATG CAAGTGCTCC CGGAAGGACC CAAGATCCGC TACAGCGACG 50

<210> 651
<211> 50

<212> DNA
<213> Homo sapiens

<220>
<221> allele
<222> (26)...(0)
<223> single nucleotide polymorphism

<221> misc_feature
<222> (25)...(26)
<223> Nucleotide deleted between bases 25 and 26

<221> misc_feature
<222> (0)...(0)
<223> Accession number cg44010855

<400> 651
TCCAAATGCA AGTGCTCCCG GAAGGACCCA AGATCCGCTA CAGCGACGTG

50

<210> 652
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 652
Pro Ala Glu Lys Asp Ala Val Pro Gln Thr Phe Ser Val Leu
1 5 10

<210> 653
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 653
Arg Leu His Arg Leu Arg Gly Glu Gln Met Ala Ser Tyr Phe
1 5 10

<210> 654
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 654
Gly Ala Pro Leu Tyr Met Asp Ser Arg Ala Asp Arg Lys Leu
1 5 10

<210> 655
<211> 14
<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 655

Gln Ala Gly Thr Thr Leu Asp Leu Asp Leu Gly Gly Lys His
1 5 10

<210> 656

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 656

Lys Cys Glu Cys Ser Arg Ala Tyr Gln Met Asp Leu Ala Thr
1 5 10

<210> 657

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 657

Asp Val Arg Gly Asn Leu Lys Gly Asn Thr Glu Gly Leu Gln
1 5 10

<210> 658

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 658

Val Pro Leu Glu Thr Pro Arg Val His Ser Arg Ala Pro Ser
1 5 10

<210> 659

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 659

Pro Phe Gly Val Ile Val Arg Arg Gln Leu Asp Gly Arg Val
1 5 10

<210> 660
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 660
Trp Gly Lys Glu Pro Lys Leu Glu Ser Ala Trp Met Asn Gly
1 5 10

<210> 661
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 661
Ser Arg Gly Tyr Arg Asn Arg Arg Ser Ser Arg Glu Thr Arg
1 5 10

<210> 662
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 662
Gly Ala Met Leu Leu Asn Ile Ser Gly His Val Lys Glu Ser
1 5 10

<210> 663
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 663
Gln Gly Gly Lys Leu Ser Val Val Leu Arg Ala Glu Asp Ile
1 5 10

<210> 664
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 664
Pro Gly Asp Ile Ser Arg Leu Leu Glu Phe Thr Lys Ala His
1 5 10

<210> 665
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 665
Val Val Ile Pro Ser Asp Phe Phe Gln Ile Val Gly Gly Ser
1 5 10

<210> 666
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 666
Lys Arg Lys Leu Phe Ile Arg Ser Met Gly Glu Gly Thr Ile
1 5 10

<210> 667
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 667
Glu Gln Ala Arg Gln Gly Leu Lys Gly Leu Glu Glu Thr Val
1 5 10

<210> 668
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 668
Cys
1

<210> 669
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 669
Ser Phe Leu Ile Ser Pro Leu Thr Pro Ala His Ala Gly Thr
1 5 10

<210> 670
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 670
Trp Cys Ser Lys Lys Lys Asp Ala Ala Val Met Asn Gln Glu
1 5 10

<210> 671
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 671
Val Ala Leu Pro Gln Pro Leu Gly Val Pro Asn Glu Ser Gln
1 5 10

<210> 672
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 672
Gly Ile Gln Asn Lys Glu Val Glu Val Arg Ile Phe His Cys
1 5 10

<210> 673
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 673

Val Asp Asn Ile Arg Ser Val Phe Gly Asn Ala Val Ser Arg
1 5 10

<210> 674

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 674

Leu Thr Ile Gln Leu Ile Glu Asn His Phe Val Asp Glu Tyr
1 5 10

<210> 675

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 675

Tyr Gly Ile Pro Phe Ile Gln Thr Ser Ala Lys Thr Arg Gln
1 5 10

<210> 676

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 676

Ser Ala Ser Lys Gln Ala Val Arg Pro Val Leu Ala Thr Thr
1 5 10

<210> 677

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 677

Tyr Cys Met Val Phe Leu Ala Leu Tyr Val Gln Ala Arg Leu
1 5 10

<210> 678
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 678
Val Gly Thr Tyr Arg Cys Val Pro Gly Lys Lys Gly Gly Tyr
1 5 10

<210> 679
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 679
Gly Arg Ala Thr Ser Gly Ser Glu His Gln Phe Cys Gly Gly
1 5 10

<210> 680
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 680
Gly Glu Trp Ile Thr Val Asp Gln Thr Thr Thr Ala Asn Arg
1 5 10

<210> 681
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 681
Pro Glu Leu Val Leu Glu Leu Pro Ile Arg His Pro Lys Phe
1 5 10

<210> 682
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)

<223> cSNP translation

<400> 682

Glu Trp Phe Lys Asp Leu Ala Leu Lys Trp Tyr Gly Leu Pro
1 5 10

<210> 683

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 683

Tyr Ile Thr Gly Asp Arg Gly Tyr Met Asp Lys Asp Gly Tyr
1 5 10

<210> 684

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 684

Ser Gly Tyr Pro Lys Met Ser Ala His Thr His Ser Ser Phe
1 5 10

<210> 685

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 685

Val Val Lys Ala Phe Val Ile Leu Ala Ser Gln Phe Leu Ser
1 5 10

<210> 686

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 686

Asp Pro Leu Ile Tyr Ala Phe Arg Ser Gln Glu Met Arg Lys
1 5 10

<210> 687

<211> 14

<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 687
Leu Ala Thr Leu Pro Glu Tyr Val Val Tyr Lys Pro Gln Met
1 5 10

<210> 688
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 688
Leu Ala Pro Gln Gln Arg Val Ala Pro Gln Gln Lys Arg Ser
1 5 10

<210> 689
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 689
Leu Tyr Arg Asp Ile Phe Glu His Leu Arg Asp Glu Ser Gly
1 5 10

<210> 690
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 690
Asp Phe Tyr Tyr Leu Gly Ala Phe Phe Gly Gly Ser Val Gln
1 5 10

<210> 691
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 691

Leu Thr Met Val Thr Leu Val Thr Leu Pro Leu Leu Phe Leu
1 5 10

<210> 692

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 692

Arg Ala Leu Tyr Leu Leu Ile Arg Arg Val Leu His Leu Gly
1 5 10

<210> 693

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 693

Glu Glu Ala Met Asn Ala Val Tyr Ser Gly Tyr Val Tyr Thr
1 5 10

<210> 694

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 694

Ser Ala Ala Ser Tyr Asn Val Lys Thr Ala Tyr Arg Lys Tyr
1 5 10

<210> 695

<211> 8

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 695

Glu Ser Thr Thr Val Gly Ser Ser
1 5

<210> 696

<211> 14

<212> PRT

<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 696
Gln Gln Trp Ser Glu His His Ala Phe Leu Ser Gln Gly Ser
1 5 10

<210> 697
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 697
Asp Asn Phe Ser Val Thr Glu Val Pro Phe Thr Glu Ser Ala
1 5 10

<210> 698
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 698
Pro Gly Arg Arg Gln Arg Leu Thr Met Ala Ile Arg Thr Val
1 5 10

<210> 699
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 699
Leu Ala Gly Lys Val Ala Gln Val Lys Lys Asn Gly Arg Ile
1 5 10

<210> 700
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 700
Leu Glu Asn Gln Lys Lys Val Arg Lys Lys Val Leu Ile

1 5 10

<210> 701
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 701
Val Ser Asp Glu Glu Leu Asp Gln Met Leu Asp Ser Gly Gln
1 5 10

<210> 702
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 702
Trp Leu Gly Phe Asn Lys Gln Arg Gly His Leu Gln Ile Ala
1 5 10

<210> 703
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 703
Glu Pro Glu Cys Arg Glu Val Phe His Arg Arg Ala Arg Ala
1 5 10

<210> 704
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 704
Leu Pro Cys Gly Pro Gly Val Lys Gly Arg Cys Phe Gly Pro
1 5 10

<210> 705
<211> 14
<212> PRT
<213> Homo sapiens

<220>

<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 705

Ser Ala Met Asp Thr Arg Leu Leu Cys Cys Ala Val Ile Cys
1 5 10

<210> 706
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 706

Asn Thr Arg Leu Leu Cys His Val Met Leu Cys Leu Leu Gly
1 5 10

<210> 707
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 707

Gln Thr Gly Asp Ser Ala Ile Tyr Leu Cys Ala Val Glu Ala
1 5 10

<210> 708
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 708

Val Tyr Leu Cys Ala Val Asp Ala Tyr Ser Asn Asp Tyr Lys
1 5 10

<210> 709
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 709

Gln Lys Gln Met Glu Leu Asp Ser Ile Leu Val Ala Leu Leu
1 5 10

<210> 710
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 710
Gly Ser Thr Val Ile Ala Gly Ser Ile Asn Ala His Gly Ser
1 5 10

<210> 711
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 711
Pro Val Met Gly Leu Met Ile Tyr Met Met Val Met Asp His
1 5 10

<210> 712
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 712
Ser Ser Gln Asp Pro Ala Ser Val Arg Glu Cys His Asp Pro
1 5 10

<210> 713
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 713
Val Leu Leu Leu Leu Gly Ala Cys Ala Ala Pro Pro Ala Trp
1 5 10

<210> 714
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)

<223> cSNP translation

<400> 714

Ser Leu Pro Tyr Ala Val Ala Pro Leu Ser Leu Pro Arg Gly
1 5 10

<210> 715

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 715

Tyr Gly Pro Gln Cys Gln Leu Val Ile Gln Cys Glu Pro Leu
1 5 10

<210> 716

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 716

Thr Met Asp Cys Thr His Ser Leu Gly Asn Phe Ser Phe Ser
1 5 10

<210> 717

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 717

Gly Thr Thr Glu Thr Gly Gly Gln Gly Lys Gly Thr Ser Lys
1 5 10

<210> 718

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 718

Cys Asn Gly Val Ala Val Cys Ser Asn Gln Asp Leu Ile Thr
1 5 10

<210> 719

<211> 14

<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 719
Leu Val Ser Tyr Cys Pro Arg Arg Leu Gln Gln Leu Leu Pro
1 5 10

<210> 720
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 720
Arg Pro Gly Ser Pro Glu Gly Pro Leu Gly Pro Gly Gly Pro
1 5 10

<210> 721
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 721
Tyr Ala Glu Arg Tyr Gln Met Pro Thr Gly Ile Lys Gly Pro
1 5 10

<210> 722
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 722
Asn Pro Leu Val Pro Gly Thr Pro Gly Arg Pro Gly Ile Pro
1 5 10

<210> 723
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 723

Val Thr Asn Arg Pro Cys Gly Ser Gln Val Arg Cys Glu Gly
1 5 10

<210> 724

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 724

Ser Thr Thr Cys Val Arg Gln Ala Gln Cys Gly Gln Asp Phe
1 5 10

<210> 725

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 725

Gly Gly Gln Pro Cys Thr Ala Pro Leu Val Ala Phe Gln Pro
1 5 10

<210> 726

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 726

Ser Pro Arg Ile Ser Ser Pro Arg Pro Gln Gly Leu Ser Asn
1 5 10

<210> 727

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 727

Pro Met Thr Gln Thr Thr Ser Leu Lys Thr Ser Trp Val Asn
1 5 10

<210> 728

<211> 12

<212> PRT

<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 728
Ile Asn Ala Tyr Ile Ser His Leu Gly Phe Arg Phe
1 5 10

<210> 729
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 729
Leu His Ser Gly His Arg Gln Arg Pro Glu Phe Arg Pro Tyr
1 5 10

<210> 730
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 730
Cys Ser Gln Glu Ala Lys Gln Ser Ala Tyr Cys Pro Tyr Ser
1 5 10

<210> 731
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 731
Arg Thr Ile Leu Ile Phe Gly Arg Cys Gln Glu Gln Phe Gly
1 5 10

<210> 732
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 732
Asn Ala Asp Ile Ile Glu Ala Leu Arg Lys Lys Gly Phe Lys

1 5 10

<210> 733
<211> 9
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 733
Asp Arg Val Glu Asn Tyr Lys Pro Arg
1 5

<210> 734
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 734
Ile Gln Thr Ala Glu Leu Thr Pro Glu Leu Val Ile Ser Asn
1 5 10

<210> 735
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 735
Glu Gln Gln Gln Glu Gln His Gln Glu Gln Gln Glu Gln
1 5 10

<210> 736
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 736
Leu Ser Phe Ile Ser Thr Glu Leu Lys Tyr Pro Gly Met Pro
1 5 10

<210> 737
<211> 14
<212> PRT
<213> Homo sapiens

<220>

<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 737
Ile Leu Pro Ser Gly Leu Ala Phe Ile Ser Thr Gly Leu Lys
1 5 10

<210> 738
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 738
Phe Tyr Leu Ala Met Asn Glu Glu Gly Lys Leu Tyr Ala Lys
1 5 10

<210> 739
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 739
Gly Leu Asp Gln Lys Arg Ile Lys Tyr Val Val Gly Glu Leu
1 5 10

<210> 740
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 740
Ala Gly Leu Gly Gln Val Arg Leu Ile Val Gly Ile Leu Leu
1 5 10

<210> 741
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 741
Thr Leu Thr Ser Val Gly His Gln Ser Val Thr Pro Gly Glu
1 5 10

<210> 742
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 742
Gly Gln Lys Thr Leu Thr Pro Val Gly Tyr Gln Ser Val Thr
1 5 10

<210> 743
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 743
Thr Glu Ile Thr Gly Ala Thr Thr Met Thr Ser Val Gly His
1 5 10

<210> 744
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 744
Gly Val Tyr Ile Leu Thr Tyr Asn Thr Ser Gln Tyr Asp Thr
1 5 10

<210> 745
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 745
Pro Leu Ser Phe His Val Ile Trp Ile Ala Ser Phe Tyr Asn
1 5 10

<210> 746
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 746

Ser Thr Ile Val Phe Leu Cys Pro Trp Ser Arg Gly Asn Phe
1 5 10

<210> 747

<211> 13

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 747

Pro Asp Pro Arg Glu Ala Cys Gly Ser Ser Ser Tyr Val
1 5 10

<210> 748

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 748

Ala Lys Asp Met Asn Gly Thr Ser Leu His Gly Lys Ala Ile
1 5 10

<210> 749

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 749

Arg Ser Ala Arg Gly Ser Arg Gly Gly Thr Arg Gly Trp Leu
1 5 10

<210> 750

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 750

Asn Phe Thr Ser Lys Tyr His Met Lys Val Leu Tyr Leu Ser
1 5 10

<210> 751

<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 751
Ile Gln Tyr Thr Tyr Leu Gly Gly His Val Cys Leu Ser Ala
1 5 10

<210> 752
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 752
Leu Asp Pro Asn Asn Pro Asp Ala Asn Trp Ile His Ala Arg
1 5 10

<210> 753
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 753
Cys Leu Thr Glu Arg Gln Ser Lys Ile Trp Phe Gln Asn Arg
1 5 10

<210> 754
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 754
Pro Tyr Arg Ile Ala His Ala Val Ile Lys Ala His Ala Arg
1 5 10

<210> 755
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 755

Lys Arg Lys Lys Glu Val His Ala Thr Ser Pro Ala Pro Ser
1 5 10

<210> 756

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 756

Ile Lys Asn Glu Ala Arg Leu Pro Cys Leu Pro Thr Pro Gly
1 5 10

<210> 757

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 757

Ala Arg Lys Phe Phe Glu Asn Leu Pro Asp Gly Thr Trp Asn
1 5 10

<210> 758

<211> 8

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 758

Phe Ser Tyr Ser Ala Ser Ser Thr
1 5

<210> 759

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 759

Pro Val Ser Ser Ala Asn Val Met Ser Gly Ile Ser Ser Val
1 5 10

<210> 760

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 760

Trp Glu Arg Phe Val His Arg Glu Asn Gln His Leu Val Ser
1 5 10

<210> 761

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 761

Gly Thr Glu Asp Leu Tyr Gly Tyr Ile Asp Lys Tyr Asn Ile
1 5 10

<210> 762

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 762

Lys Gln Leu Tyr Gln Thr Phe Thr Asp Tyr Asp Ile Arg Phe
1 5 10

<210> 763

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 763

Gly Ser Leu His Pro His Pro Pro Tyr His Ile Arg Val Ala
1 5 10

<210> 764

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 764

Leu Tyr Ser Arg Leu Gly Gly Gln Pro Val Tyr Leu Pro Thr
1 5 10

<210> 765
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 765
Ala Pro Pro Gly Ala Tyr His Gly Ala Pro Gly Ala Tyr Pro
1 5 10

<210> 766
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 766
Val Met Pro His Ser Ser Glu His Lys Thr Ala Gln Pro Asn
1 5 10

<210> 767
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 767
Thr Cys Val Val Glu His Thr Gly Ala Pro Glu Pro Ile Leu
1 5 10

<210> 768
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 768
Val Gly Phe Leu Val Gly Ile Val Leu Ile Ile Met Gly Thr
1 5 10

<210> 769
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 769
Gln Ser Val Val Ser Cys Ala
1 5

<210> 770
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 770
Pro Gly Pro Thr Val Arg Ala Gly Glu Asn Val Thr Leu Ser
1 5 10

<210> 771
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 771
Cys Ser Gly Val Trp Gly Ala Asp Thr Glu Glu Arg Leu Val
1 5 10

<210> 772
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 772
Ala Leu Leu Arg His Glu Leu Lys Gly Tyr Gln Lys Trp Val
1 5 10

<210> 773
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 773
Leu Leu Arg His Glu Trp Gln Gly Tyr Gln Lys Trp Val Arg
1 5 10

<210> 774
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 774
Gly Phe His Tyr Gly Val Leu Ala Cys Glu Gly Cys Lys Gly
1 5 10

<210> 775
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 775
Asp Gly Pro Glu Gln Ala His Arg Gln Arg Arg Gln Arg
1 5 10

<210> 776
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 776
Glu
1

<210> 777
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 777
Asn His Asn Gly Glu Trp Phe Glu Ala Gln Thr Lys Asn Gly
1 5 10

<210> 778
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 778

Leu Val Val Val Gly Ala Cys Gly Val Gly Lys Ser Ala Leu
1 5 10

<210> 779

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 779

Ile Leu Asp Thr Ala Gly His Glu Glu Tyr Ser Ala Met Arg
1 5 10

<210> 780

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 780

Phe Gly His Gln Glu Asn Ser Gln Asn Glu Glu Ile Leu Asn
1 5 10

<210> 781

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 781

Leu Thr Tyr Pro Pro Gly Pro Arg Thr Gln Leu Arg Glu Asp
1 5 10

<210> 782

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 782

Gln Pro Gln Ala Arg Gln Glu Glu Gln Val Arg Val Val Arg
1 5 10

<210> 783

<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 783
Asn Ala Val Lys Ala Glu Thr Arg Gln Gln Phe Arg Ser Leu
1 5 10

<210> 784
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 784
Ala Gln Ile Phe Asp Tyr Ser Glu Ile Pro Asn Phe Pro Arg
1 5 10

<210> 785
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 785
Asp Asp Asp Asp Ala Pro Arg Pro Ser Gln Phe Glu Glu Asp
1 5 10

<210> 786
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 786
His Val Leu Arg Met His Gly Tyr Arg Ala Pro Gly Glu Gln
1 5 10

<210> 787
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 787
Glu Thr Gln Leu Gly Thr Leu Ala Gln Phe Pro Asn Thr Leu
1 5 10

<210> 788
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 788
Glu Ala Met Glu Arg Phe Gly Glu Asp Glu Gly Phe Ile Lys
1 5 10

<210> 789
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 789
Glu Glu Glu Lys Pro Leu Ala Arg Asn Glu Phe Gln Arg Gln
1 5 10

<210> 790
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 790
Asp Ala His Phe Asp Glu His Glu Arg Trp Thr Asn Asn Phe
1 5 10

<210> 791
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 791
Arg Trp Thr Asn Asn Phe Thr Glu Tyr Asn Leu His Arg Val
1 5 10

<210> 792
<211> 14
<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 792

Asp Ser Gly Leu Leu Gln Cys Lys Leu Ala Leu Leu Ser Val

1

5

10

<210> 793

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 793

Glu Ala Glu Arg Val Lys Glu Gln Thr Phe Arg Glu Lys Ala

1

5

10

<210> 794

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 794

Ala Phe Val Val Leu Ala Leu Gln Phe Leu Ser His Asp Pro

1

5

10

<210> 795

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 795

Thr Gly Lys Ile Gln Arg Thr Lys Leu Arg Asp Lys Glu Trp

1

5

10

<210> 796

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 796

Val Asp Tyr Met Thr Gln Ala Arg Gly Gln Arg Ser Ser Leu
1 5 10

<210> 797
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 797
Asn Asn Leu Leu Tyr Ile Thr Pro Glu Ala Phe Gln Asn Leu
1 5 10

<210> 798
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 798
Leu Arg Ile Gln Cys Leu Cys Arg Lys Gln Ser Ser Lys His
1 5 10

<210> 799
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 799
Leu Glu Lys Ile Gln Pro Met Thr Gln Asn Gly Gln His Pro
1 5 10

<210> 800
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 800
Trp Met Ile Phe Val Val Ile Ala Ser Val Phe Thr Asn Gly
1 5 10

<210> 801
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 801
Ser Ile Asn Leu Phe Ser Gly Ile Phe Phe Leu Thr Cys Met
1 5 10

<210> 802
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 802
Val Asp Ile Cys Phe Ser Ser Thr Thr Val Pro Lys Met Leu
1 5 10

<210> 803
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 803
Glu Ser Asn Thr Thr Gly Thr Thr Ala Phe Ser Met Pro Ser
1 5 10

<210> 804
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 804
Val His Pro Val Arg Pro Leu Arg Leu Glu Ser Phe Ser Ala
1 5 10

<210> 805
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 805
Arg Gly Ala Arg Pro Gly Pro Arg Val Pro Lys Thr Leu Val
1 5 10

<210> 806
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 806
Lys Ala Phe Leu Thr Ser His Ser Leu Arg Ile His Val Arg
1 5 10

<210> 807
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 807
Ser Glu Ala Ser Ser Ala Phe Phe Met Ala Lys Lys Lys Thr
1 5 10

<210> 808
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 808
Glu Leu Tyr Arg Asp Ile Leu Gln His Leu Arg Asp Glu Ser
1 5 10

<210> 809
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 809
Tyr Val Phe Thr Asp Gln Leu Ala Ala Val Pro Arg Val Thr
1 5 10

<210> 810
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 810

Leu Ser Val Leu Glu Val Gly Ala Tyr Lys Arg Trp Gln Asp
1 5 10

<210> 811

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 811

Val Asp Val Asp Met Glu Ile Arg Asp His Val Gly Val Glu
1 5 10

<210> 812

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 812

Phe Gly Thr Leu His Pro Ser Phe Tyr Gly Ser Ser Arg Glu
1 5 10

<210> 813

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 813

Glu Gly Asp Phe Tyr Tyr Met Gly Gly Phe Phe Gly Gly Ser
1 5 10

<210> 814

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 814

Gly Gly Ser Val Gln Glu Met Gln Arg Leu Thr Arg Ala Cys
1 5 10

<210> 815

<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 815
Val Val Glu Leu Thr Gln Asp Asp Ala Leu Gly Ser Arg Trp
1 5 10

<210> 816
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 816
Lys Asp Phe His Lys Asp Met Leu Lys Pro Ser Pro Gly Lys
1 5 10

<210> 817
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 817
Thr Asn Asn Cys Tyr Arg His Ala Ile Val Thr Thr Ser Ile
1 5 10

<210> 818
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 818
Gly Phe Val Val Phe Ser Ser Leu Gly Tyr Met Ala Gln Lys
1 5 10

<210> 819
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 819
Met Asp Glu Ala Ala Arg Pro Glu Ala Trp Asp Ser Tyr Arg
1 5 10

<210> 820
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 820
Ser Pro Gln Ser Ser Ala Arg Gly Lys Pro Ala Met Ser Tyr
1 5 10

<210> 821
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 821
Leu Glu Asp Leu Ala Gly Trp Lys Glu Leu Phe Gln Thr Pro
1 5 10

<210> 822
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 822
Val Gln Asp Ile Leu Arg Leu Glu Met Pro Ala Ser Lys Ile
1 5 10

<210> 823
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 823
Ser Glu Arg Glu Thr Glu His Thr Pro Ala Leu Ile Met Val
1 5 10

<210> 824
<211> 14
<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 824

Lys Val Leu Asp His Trp Cys Ile Met Thr Ser Glu Glu Glu
1 5 10

<210> 825

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 825

Gln Glu Val His Gly Pro Tyr Pro Asp Ser Ser Phe Leu Thr
1 5 10

<210> 826

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 826

Gln Glu Val His Gly Pro Ile Pro Asp Ser Ser Phe Leu Thr
1 5 10

<210> 827

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 827

Glu Leu Cys His Glu Lys Gly Ile Leu Glu Lys Tyr Gly His
1 5 10

<210> 828

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 828

Asn Asn Asn Leu Arg His Thr Asp Glu Met Phe Trp Asn His
1 5 10

<210> 829
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 829
Gly Met Ala Ser Ser Cys Ser Val Gln Val Lys Leu Glu Leu
1 5 10

<210> 830
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 830
Pro Ser Ile Phe Ile Tyr Arg His Thr Ala Ser Gly Lys Thr
1 5 10

<210> 831
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 831
Pro Pro Pro Pro Gly Ala Pro Gly Gly Ser Gln Asp Thr
1 5 10

<210> 832
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 832
Arg Ala Ala Leu Glu Arg Gly Lys Ala Ile Glu Lys Asn Leu
1 5 10

<210> 833
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 833
Thr Gly Gly Leu Leu Leu Arg Leu Ala Leu Met Leu Gln Leu
1 5 10

<210> 834
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 834
Asp Ser Ser Ser Ser Asn Gly Lys Ala Lys Asn Pro Pro Gly
1 5 10

<210> 835
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 835
Leu Glu Pro Gln Trp Tyr Ser Val Leu Glu Lys Asp Ser Val
1 5 10

<210> 836
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 836
Gln Lys His Ser Ser Gly Xaa Ser Asn Thr Ser Thr Ala Asn
1 5 10

<210> 837
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 837
Trp Gly Thr Glu Asp Asp Ala Thr Gln Ser Tyr His Asn Gly
1 5 10

<210> 838
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 838
Arg Cys Met Gly Thr Val Asn Leu Asn Gln Ala Arg Gly Ser
1 5 10

<210> 839
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 839
Phe Ser Lys Leu Leu Gly Pro Leu Ser Ala Lys Lys Tyr Leu
1 5 10

<210> 840
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 840
Thr Ser Lys Ile Leu Phe Phe Ser Gln Gly Ser Glu Ile Ala
1 5 10

<210> 841
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 841
Tyr Val Gln Pro Pro Glu Val Ile Gly Pro Met Arg Pro Glu
1 5 10

<210> 842
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 842

Gly Pro Asp Gly Gln Glu Val Asp Pro Pro Asn Pro Glu Glu

1

5

10

<210> 843

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 843

Lys Gly Gly Glu Gln Lys Arg His Glu Lys Ile Ser Ala Ser

1

5

10

<210> 844

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 844

Ala Pro Gln Gln Glu Gly Glu Ala Ser Lys Glu Lys Glu Glu

1

5

10

<210> 845

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 845

Lys Val Val Gln Ser Pro Gln Ser Leu Val Val His Glu Gly

1

5

10

<210> 846

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 846

Leu Asn Cys Ser Tyr Glu Met Thr Asn Phe Arg Ser Leu Leu

1

5

10

<210> 847

<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 847
Thr Asn Phe Arg Ser Leu Gln Trp Tyr Lys Gln Glu Lys Lys
1 5 10

<210> 848
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 848
Leu Asp Lys Lys Glu Leu Ser Ser Ile Leu Asn Ile Thr Ala
1 5 10

<210> 849
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (8)...(0)
<223> cSNP translation

<400> 849
Glu Tyr Val Val Gly Ala Pro His Leu Glu Leu Asp Pro Gly
1 5 10

<210> 850
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 850
Phe Phe Lys Arg Asn Arg His Thr Pro Gly Arg Arg
1 5 10

<210> 851
<211> 9
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 851

Thr Ile Gln Pro Pro Arg Glu

1

5

<210> 852

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (8)...(0)

<223> cSNP translation

<400> 852

Gly Ile Val Gly Gln Lys Gly Arg Pro Trp Leu Pro Arg Thr

1

5

10

<210> 853

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (8)...(0)

<223> cSNP translation

<400> 853

Gly Gly Lys Met Gly Gly Arg Lys Arg Leu Gln Lys

1

5

10

<210> 854

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (8)...(0)

<223> cSNP translation

<400> 854

Tyr Ser Ser Tyr Gly Gln Ser Leu Phe Thr Val Leu Trp Trp

1

5

10

<210> 855

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (8)...(0)

<223> cSNP translation

<400> 855

Glu Gln Leu Arg Arg Gln Leu Asp Pro Leu Arg Thr Ala His

1

5

10

<210> 856

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 856

Ser Thr Glu Cys Trp Met Asn Ala Ala Cys Leu Ala Pro Gly
1 5 10

<210> 857

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (9)...(0)

<223> cSNP translation

<400> 857

His Gly Val Leu Asp Ala Cys Leu Ile His Pro Gly Pro Ala
1 5 10

<210> 858

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 858

Arg Asp Lys Gly Ser Gly Arg Ala Cys Gly Leu Glu Gly Gln
1 5 10

<210> 859

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 859

Thr Asp Phe Phe Phe Phe
1 5

<210> 860

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (8)...(0)

<223> cSNP translation

<400> 860

Gly Ala Gly Ser Val Ser Asp His His Ser Ile Thr Lys
1 5 10

<210> 861
<211> 12
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 861
Arg Tyr Leu Asp Trp Ile Leu Trp Ala His Gln Arg
1 5 10

<210> 862
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (8)...(0)
<223> cSNP translation

<400> 862
Ala Glu Leu Arg Leu Leu Arg Ala Gln Val Lys Ser Gly Ala
1 5 10

<210> 863
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (8)...(0)
<223> cSNP translation

<400> 863
Ala Glu Leu Arg Leu Leu Arg Ala Gln Val Lys Ser Gly Ala
1 5 10

<210> 864
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (8)...(0)
<223> cSNP translation

<400> 864
Pro His Cys Arg Pro Gly Ala Trp Pro Ala Thr Glu Arg Gly
1 5 10

<210> 865
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (8)...(0)
<223> cSNP translation

<400> 865
Ile His Phe Glu Asp Tyr Gly Val Leu Gly His His Gln Leu
1 5 10

<210> 866
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 866
Asn Phe Ile Leu Ala Cys Pro Arg
1 5

<210> 867
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 867
Arg Glu Lys Leu Arg Asn Phe His Phe Ser Met Ser Arg
1 5 10

<210> 868
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (8)...(0)
<223> cSNP translation

<400> 868
Leu Leu Leu Leu Leu Arg Arg Pro Ala Gln Pro Gln Leu
1 5 10

<210> 869
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 869
Lys Arg Val Ala Gly Gly Leu Arg
1 5

<210> 870
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (8)...(0)
<223> cSNP translation

<400> 870
Gly Lys Arg Val Ala Gly Gly Leu Arg
1 5

<210> 871
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 871
Ser Ser Gly Arg Pro Thr Gly Tyr Cys Leu Gln Leu Gln Gln
1 5 10

<210> 872
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (8)...(0)
<223> cSNP translation

<400> 872
Gln Arg Ser Ile Ser Ala Asp
1 5

<210> 873
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (8)...(0)
<223> cSNP translation

<400> 873
Arg Ala Pro Val Ile Leu Gly Pro Pro Thr Cys Ser Ser
1 5 10

<210> 874
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 874

Met Gly Leu Ser Gly Phe Leu Thr Gly Pro Pro Pro Gly
1 5 10

<210> 875

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (8)...(0)

<223> cSNP translation

<400> 875

Leu Met Gly Leu Ser Gly Phe Leu Thr Gly Pro Pro Pro Pro
1 5 10

<210> 876

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 876

Pro Arg Thr Pro Ala Glu Pro Pro Pro Leu Gly Arg Gln Ala
1 5 10

<210> 877

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (7)...(0)

<223> cSNP translation

<400> 877

Gly Thr Gly Asp Trp Arg Glu Pro Gly Ala Ala Ser Glu Arg
1 5 10

<210> 878

<211> 14

<212> PRT

<213> Homo sapiens

<220>

<221> VARIANT

<222> (9)...(0)

<223> cSNP translation

<400> 878

Gln Gly Arg Gln Ser Lys Gly Leu Arg Arg Arg Thr Trp Pro
1 5 10

<210> 879

<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (8)...(0)
<223> cSNP translation

<400> 879
Lys Cys Lys Cys Ser Arg Lys Asp Pro Arg Ser Ala Thr Ala
1 5 10

<210> 880
<211> 14
<212> PRT
<213> Homo sapiens

<220>
<221> VARIANT
<222> (7)...(0)
<223> cSNP translation

<400> 880
Cys Lys Cys Ser Arg Lys Asp Pro Arg Ser Ala Thr Ala Thr
1 5 10

15